

STUDIA GEOBOTANICA

An international journal

Vol. 16 1998

EDITORS

G. Cristofolini - Bologna
Lj. Ilijanić - Zagreb
E. Mayer - Ljubljana
P.L. Nimis - Trieste
L. Orlóci - London, On.
F. Pedrotti - Camerino
S. Pignatti - Roma
A. Pirola - Pavia
J. Poelt - Graz
L. Poldini - Trieste
E. Wilkus Pignatti - Trieste

EDITOR IN CHIEF

† D. Lausi - Trieste

SECRETARIES

M. Codogno - Trieste
M. Tretiach - Trieste



Dipartimento di Biologia
Sezione di Geobotanica ed Ecologia vegetale

Università di Trieste

INSTRUCTIONS TO AUTHORS

STUDIA GEOBOTANICA publishes original articles in the field of geobotany. It also publishes reviews of published works, short communications, notices of scientific meetings and summaries of proceedings. MANUSCRIPTS should be submitted in triplicate including tables and figures to the Secretary of the Editorial Board. The Secretary will confirm receipt of the manuscript. The suitability of submitted manuscripts will be judged by appropriate members of the Editorial Board. Appropriate referees will be consulted. The Secretary will inform authors of the acceptance or rejection of manuscripts or the need for substantial revisions or reduction. Manuscript should be typed, double-spaced, on white bond paper A4 format with 35 mm margin on all side. All pages should be numbered. Send also one copy on 3.5" disk based on Microsoft Word5.5 program.

LANGUAGE should be preferably English. Italian, German, Spanish and French are also accepted.

THE TITLE should be informative and brief. The name(s) (given name-surname) of the author(s) should be placed below the title. References to research projects, institutional publication numbers, etc. should be placed in a footnote in the first page.

KEYWORDS, no more than 10, in English, should be placed on the title page in alphabetical order.

ABSTRACT. An English abstract must be included. For non English text, the abstract must include the title. A summary in a second language may also be added; it should not exceed one manuscript page and is placed before the acknowledgements.

MAIN TEXT should be clearly divided into sections. Chapter headings should be given without designations. New paragraphs should be indicated by indentation. Footnotes should be avoided.

APPENDICES should be placed at the end of the main text.

ACKNOWLEDGEMENTS and dedications should be simply phrased and given before the references.

REFERENCES to articles and books should be limited to published works. Citation of references in the text: Dolcher & Pignatti (1971) or (Fukarek 1977; Clotet *et al.* 1988; Podani & Borhidi 1989). Citation of references:

Malyshev L.I., 1991. *Some quantitative approaches to problems of comparative floristics*. In: Nimis P.L. & Crovello T.J. (eds.), *Quantitative approaches to phytogeography*. Kluwer Academic Publishers, Dordrecht, Boston, London, 15-33.

Poelt J., 1969. *Bestimmungsschlüssel europäischer Flechten*. J. Cramer, Vaduz, 757 pp.

Trevisan R., Cappelletti E.M. & Caniato P., 1983. *Epicarp micro-characters in some european Angelica species*. Atti Accad. Naz. Lincei, Rend. Cl. Sci. fis., mat. e nat., 74: 77-82.

List in alphabetic sequence with: single authors, by date; two authors, alphabetically; then by date; three or more authors by date only.

NAME AND ADDRESS of the author(s) should be placed at the end of the article.

TABLES should be typed on separate pages, and not in the text itself. They should be submitted in camera-ready form to be reduced to page-size. Large tables should be prepared in the same way. Tables must be numbered. Author name(s) and table number should be written in pencil on each table and table copy. Title and legend to the tables should be typed on one or more separate pages.

ILLUSTRATIONS. Figures such as drawings and graphs should be done in India ink on smooth tracing paper. Names, numbers or other symbols should also be written in India ink or, preferably, by or a similar system. Reduction of these figures should be taken into account when preparing the lettering. All figures should be numbered. Author name(s) and figure number should be written in pencil on each figure and figure copy. Title and legend to the figures should be typed on one or more separate pages.

Legends to figures and tables are given both in the language of the main text and in English.

Plates should be avoided, but when used, they should be numbered and supplied with informative title and legend as the figures.

EDITORIAL POLICY. Manuscripts are considered for publication under the assumption that they have not been published elsewhere. Manuscripts are evaluated by reviewers whose decision is binding. Published articles are the property of STUDIA GEOBOTANICA.

CORRESPONDENCE on matters related to manuscripts or concerning offprints should be addressed to the Secretary of the Editorial Board.

Secretaries:

Dr. Michele CODOGNO, Dr. Mauro TRETACH

Dipartimento di Biologia - Università degli Studi di Trieste

Via L. Giorgieri, 10 I-34127 TRIESTE (ITALIA)

Phone and fax number: +39 / 40 / 568855

@mail: tretach@univ.trieste.it

STUDIA GEOBOTANICA

An international journal

Vol. 16 1998

EDITORS

G. Cristofolini - Bologna
Lj. Ilijanić - Zagreb
E. Mayer - Ljubljana
P.L. Nimis - Trieste
L. Orłóci - London, On.
F. Pedrotti - Camerino
S. Pignatti - Roma
A. Pirola - Pavia
J. Poelt - Graz
L. Poldini - Trieste
E. Wilkus Pignatti - Trieste

EDITOR IN CHIEF

† D. Lausi - Trieste

SECRETARIES

M. Codogno - Trieste
M. Tretiach - Trieste




Dipartimento di Biologia
Sezione di Geobotanica ed Ecologia vegetale

Università di Trieste

Il 16 marzo 1997 è venuto a mancare il Prof. Duilio Lausi, che è stato uno dei fondatori di questa rivista, e suo Direttore per lunghi anni. Chi lavora nell'ambito della Geobotanica conosce l'importante contributo da Lui dato in questa disciplina. In questa sede abbiamo voluto ricordare la Sua figura attraverso il ricordo di un amico e collega che per lunghi anni gli è stato vicino nella Sua ricca attività didattica e scientifica.

La Redazione

Un ricordo di Duilio Lausi (1923-1997)



Inutilmente cerco di ricordare quando incontrai Duilio Lausi per la prima volta. La sua capacità di comunicare in modo semplice e diretto con tutti lo rendeva immediatamente familiare e forse per questo non trovo un inizio della nostra amicizia, proprio come avviene con i parenti più stretti. Nonostante la irregolarità delle occasioni di trovarci, sempre per ragioni di lavoro e negli ultimi anni in particolare, per il dottorato di ricerca in geobotanica, il ritrovarsi significava riprendere un discorso già iniziato da tempo, sapendo che sarebbe stato presto interrotto, ma solo fino alla prossima riunione o telefonata. Una sequenza che purtroppo è giunta al termine.

Mi rendo conto che per commemorarlo in modo tradizionale su questa rivista di cui fu uno dei fondatori e animatore intelligente e attivo, dovrei svolgere un'analisi delle sue ricerche, dei suoi viaggi e soprattutto della sua opera di docente. In realtà non è un compito facile per la mole e la diversificazione della sua opera scientifica. Questa analisi dovrà essere fatta con maggiore distacco emotivo, per inquadrarla degnamente nel contesto generale della ricerca geobotanica contemporanea di cui, si può ben dire fino da ora senza temere di accondiscendere all'amicizia, fu tra i principali protagonisti.

Penso invece che sia necessario ora parlare della sua umanità, con l'esperienza diretta che con il passare del tempo e dei colleghi e amici, verrà a mancare lasciando al giudizio solo l'obiettività dei fatti. E per una volta

e per un amico, mi si conceda di essere soggettivo e di dire, con molta difficoltà, ciò che Duilio era per me, sperando di trovare qualche condivisione ma non sicuramente completa, perché Duilio Lausi non ebbe una personalità semplice o quantomeno riconducibile ad un paradigma, sebbene di pregio. A me è sempre apparso felicemente complesso: ansioso nelle tappe della ricerca di cui comunicava volentieri le quasi certezze, ma prima ancora i dubbi; aperto ad ogni idea nuova che fornisse un contributo originale, ma cauto di fronte a deviazioni ingiustificate dai metodi dimostratisi utili; disponibile con i giovani, ma severo nel valutare il loro impegno nelle ricerche.

Durante le discussioni scientifiche nelle aule dei convegni o in campagna, assumeva un ruolo inizialmente disposto ad ascoltare, cortese e attento, poi diveniva attivo e vivace interlocutore fino a condurre con arguzia e competenza la discussione. La sua presenza assicurava quella dose di fermento che manteneva elevato il grado di attenzione, dimostrando con molta semplicità che la ricerca, pur basandosi su dati concreti, è soprattutto un processo razionale che richiede e produce cultura.

Mai un viaggio in treno è sembrato troppo veloce e divertente come quando si faceva in sua compagnia. Era un grande estimatore delle ferrovie, non certo per il servizio di cui spesso lamentava la scomodità dei collegamenti con Trieste, ma come mezzo per dedicarsi al lavoro o alla conversazione.

I rapporti con gli allievi del dottorato meritano un cenno particolare: sapeva ascoltarli e non lesinava critiche, ma sempre in termini costruttivi, fornendo indicazioni utili sia per i metodi più adatti sia per le fonti. Ogni argomento posto in discussione era assunto come il più importante e veniva trattato con la massima serietà e approfondimento.

Tra i richiami più frequenti che rivolgeva agli allievi c'era la correttezza del rilevamento di campagna. Ricordo che durante una escursione in Valmalenco nel 1978 di un corso di fitosociologia alpina fece eseguire un rilievo con incremento di superficie su vegetazione di greto, a partire da un piccolo quadrato da 100 cm², dimostrando che la struttura della vegetazione da analizzare richiedeva appunto un approccio di quel tipo. Ne uscì un rilievo da manuale che descriveva in modo adeguato l'andamento della curva specie-superficie.

Intese l'insegnamento nel modo più sacrale immaginabile. Per Duilio la responsabilità del maestro era enorme e in molte discussioni sulle vicende che nei nostri anni attraversava l'università emergeva di frequente questo concetto come una sua preoccupazione continua sul dovere di passare le nozioni e soprattutto il comportamento corretto agli allievi. Più di una volta lo sentii esprimere questa idea in termini quasi dinastici: il metodo fitosociologico poteva essere insegnato sul terreno solo da coloro che lo avevano appreso direttamente da Braun-Blanquet o da loro allievi: una sorta di discendenza che garantisse l'aderenza più stretta possibile alla prassi del rilievo della vegetazione, per ridurre il più possibile la inevitabile componente soggettiva implicita nell'applicazione di campionamenti non distruttivi. Il requisito di trasmissione diretta del metodo era certamente una estremizzazione. Da buon biologo sapeva che non si può eliminare la variabilità anche negli operatori, ma la sua indicazione era molto utile per ricordare che corriamo continuamente il rischio di confrontare dati eterogenei.

Da queste posizioni agiva come un conservatore dell'approccio di Braun-Blanquet alla fitosociologia, ma era anche un sostenitore dell'applicazione delle analisi numeriche, raffinato e critico sulla scelta dei metodi e della interpretazione dei risultati, che, affermava, dovevano sempre essere verificati con la realtà.

Disponeva di un'ampia cultura naturalistica, soprattutto ecologica, dimostrando che le delimitazioni delle discipline sono una necessità umana, ma non devono costituire un limite per lo studio della natura.

La sua scomparsa ci ha lasciato un grosso vuoto, in particolare nel corso di dottorato di ricerca in Geobotanica

Augusto Pirola

THE SEGETAL VEGETATION OF VINEYARDS AND CROP FIELDS IN FRIULI-VENEZIA GIULIA (NE ITALY)

Livio POLDINI, Giuseppe ORIOLO and Gianluca MAZZOLINI

Dipartimento di Biologia, Università di Trieste, Via Giorgieri 10, I-34127 Trieste

Keywords: crops, Friuli-Venezia Giulia, Italy, segetal vegetation, syntaxonomy, vineyards.

Abstract: The segetal vegetation of vineyards and crop fields of Friuli-Venezia Giulia (NE Italy) was studied. 165 phytosociological relevés were analyzed by numerical classification and compared with those of Central European isoecic coenoses. Four associations were detected: the coenoses of vineyards and of weeded summer cultures of the plain and the hill belts, corresponding to those already described from Central Europe (*Geranio rotundifolii-Allietum vinealis* R. Tx. ex von Rochow 1966 and *Echinochloo-Setarietum pumilae* Felföldy 1942 corr. Mucina 1993, respectively), and the coenoses of weeded cultures of the mountain belt and of winter crop fields, described as new associations (*Galeopsido tetrahit-Galinsogietum* ass. nova and *Papaveretum apuli* ass. nova, respectively). A new subassociation of the *Echinochloo-Setarietum pumilae* is proposed, and the ecological concept of agroform is introduced. The results of this study indicate that the standardization of agronomic techniques has drastically reduced the biodiversity of these associations.

Introduction

The coenoses of vineyards and of crops (maize, soy-bean, potatoes, wheat and barley) are subjected to a very high anthropogenic pressure which changes continuously, due to the transition from artisan and semi-artisan techniques to an industrial agriculture. This has deeply modified the flora and vegetation of agroecosystems. The disappearance of many archaeophytes and the income and stabilisation of many neophytes represent very good examples of this phenomenon (Martini & Poldini 1995).

The standardization of agronomic techniques and the increased use of herbicides have drastically reduced the biodiversity of these ecosystems and the number of thermophilous species has strongly decreased. This is testified, for instance, by the disappearance of some syntaxa such as the *Polygono-Chenopodion polyspermi* alliance, described by Pignatti (1952) from the Venetian plain (Krippelová 1979). For this reason, it is necessary to reconsider the syntaxonomy of relevés from Friuli-Venezia Giulia. Nowadays it may be more appropriate to include them in the isoecic coenosis of Central Europe (*Geranio-Allietum* for vineyards of Rhine valley and *Echinochloo-Setarietum* for weeded crops of Central Europe). As a consequence, these coenoses have a very large distributional range that includes several biogeographical regions with different potential vegetations.

The weed communities of autumn-winter cereals (wheat and barley) and those of cultures of the mountain belt (potatoes and beans) are well differentiated; they are described for the first time in this paper.

When considering the strong floristic homogenization of agroecosystems, we adopted the syntaxonomical scheme proposed by Mucina (1993) that groups all the crop coenoses into a single class (*Stellarietea mediae*), rather than considering the coenosis of autumn-winter crops as belonging to the independent class of *Secalietea* (Oberdorfer 1993).

As already noted for other ecosystems (Chiappella Feoli & Poldini 1993; Poldini & Oriolo 1994), some of the relevés presented in this paper have a documentary value for vanishing plant communities.

Data and methods

165 phytosociological relevés regarding the main cultures of Friuli-Venezia Giulia have been considered. Karst relevés have been taken from Poldini (1989).

The relevés have been compared with those of vineyards of Slovenia (Vipava valley, Seljak 1989) and of Central Europe (Oberdorfer 1993; Wilmanns 1989), and with those of weeded crop fields from Italy [eastern Venetian plain (Lorenzoni 1963) and

Ferrara province (Gerdol & Piccoli 1979)], Slovenia (Vipava valley, Seljak 1989), Germany (Oberdorfer 1993), Croatia (Regula-Bevilacqua 1979) and Slovakia (Krippelová 1981).

Cover values have been converted according to van der Maarel (1979), and numerical data analysis has been carried out with Syn-Tax (Podani 1993) and Lagoli (Lagonegro & Feoli 1985) softwares.

The floristic nomenclature follows Ehrendorfer (1973), and the syntaxonomical one Mucina (1993), who refers to Ries (1992). Ecological indices have been desumed from Landolt (1977) and Poldini (1991). Data concerning pollination mechanisms have been drawn from Faegri & van der Pijl (1971), those concerning dispersion from Müller-Schneider (1986). Life forms and chorotypes have been taken from Poldini (1991).

Abbreviations in the tables are as follows: d = differential species of a part of an association; DAss = differential species of association; DAll = differential species of alliance; DO = differential species of order; O = order species; K = class species.

The concept of agroform

The evolution of cultivation techniques causes fast changes in the floristic composition and in the interspecific relationships within crop coenoses. These may change according to specific modifications of the cultivation techniques (for instance the passage from weeding to mulching) (Wilmanns 1989, 1991). In this respect we define as «agroform» of a crop coenosis the result of ecological modifications caused by agronomic techniques, provided that the connection with the original association is still evident.

In the crop associations of Friuli-Venezia Giulia three agroforms of vineyards (showing a transition from weeding to mulching or to mowing) and one of summer weeded crop fields (showing a very high use of herbicides) have been detected. Their description is given in the following chapters.

Syntaxonomical considerations

In order to define the syntaxonomical scheme of crop and vineyard associations of Friuli-Venezia Giulia, two of the most representative schemes were considered, that of Oberdorfer (1993), and that of Mucina (1993).

The scheme proposed by Oberdorfer (1993) is mainly based on Braun-Blanquet (1936). According

to this scheme, crops and ruderal communities with annual cycle are divided into two different classes, *Secalietea* Br.-Bl. 1952 and *Chenopodietea* Br.-Bl. in Br.-Bl. *et al.* 1952. The first class includes associations of autumn-winter cereals (for instance barley and wheat), while the second one includes the other crop coenoses (mainly summer weeded crops and vineyards) in the order *Polygono-Chenopodietalia* (Tx. *et Lohm.* in Tx. 1950) J. Tx. in Lohm. *et al.* 1962, and the ruderal vegetation with annual cycle in the order *Sisymbrietalia* J. Tx. in Lohm. *et al.* 1962. On the contrary, the scheme proposed by Mucina (1993) refers to Tüxen's modification (Tüxen 1950) of Braun-Blanquet's scheme. It considers a single class, *Stellarietea mediae* R. Tx., Lohm. *et Preis.* in R. Tx. 1950, subdivided in four orders: *Centaureetalia cyani* R. Tx., Lohm. *et Preis.* in R. Tx. 1950 for cultures of basic soils, *Chenopodietalia albi* R. Tx. (1937) 1950 for cultures of acid soils, *Sisymbrietalia* J. Tx. in Lohm. *et al.* 1962 for ruderal vegetation with annual cycle and, finally, *Eragrostietalia* J. Tx. *ex Poli* 1966 for cultures and ruderal associations with south-eastern European distribution.

The synthetic table (Tab. 1) highlights a substantial group of species, common to all the coenoses (*Chenopodium album*, *Veronica persica*, *Capsella bursa-pastoris*, *Cirsium arvense*, *Fallopia convolvulus*, *Stellaria media*, *Erigeron annuus*, *Sinapis arvensis*), that can be considered as characteristic of *Stellarietea*. This is in line with the fact that strong changes in agronomic techniques are levelling the floristic differentiation of the crop field associations (a valid example is the vanishing of many archaeophytes in wheat fields). Therefore, we think that Mucina's scheme fits better with the present situation in Friuli-Venezia Giulia, also because the only coenosis of *Centaureetalia* found in Friuli-Venezia Giulia (*Papaveretum apuli*) has a high presence of *Stellarietea* species.

We have introduced some significant changes to the original scheme of Mucina (1993):

a) the definition of *Centaureetalia* and *Chenopodietalia* orders seems to be more linked to the vegetation cycle than to soil characteristics. Therefore the spring-summer developing coenoses of vineyards and of summer weeded fields belong to *Chenopodietalia*, while the only association of autumn-winter cereals (*Papaveretum apuli*) belongs to *Centaureetalia*.

b) we placed the coenosis of Friuli-Venezia Giulia vineyards (*Geranio-Allietum*) into the *Chenopodietalia* order, not into the *Centaureetalia* order.

- Segetal vegetation in Friuli-Venezia Giulia -

Tab. 1 - Synthetic table of syntaxonomic units present in Friuli-Venezia Giulia (frequency values).

1 = *Papaveretum apuli*; 2 = *Geranio-Allietum*; 3 = *Echinochloo-Setarietum*, Friuli cool variant;
 4 = *Echinochloo-Setarietum*, Friuli warm variant; 5 = *Echinochloo-Setarietum xanthetosum*;
 6 = *Echinochloo-Setarietum*, Karst warm variant; 7 = *Echinochloo-Setarietum*, Karst cool variant;
 8 = *Galeopsido-Galinsogetum*.

		1	2	3	4	5	6	7	8
Char. & diff. sp. (DAss1) ass. & all. (<i>Papaveretum apuli</i> and <i>Caucalidion</i>)									
	Papaver apulum	76							
	Avena fatua	59							
	Valerianella rimosa	35							
	Fumaria vaillantii	12							
	Valerianella eriocarpa	12							
	Ajuga chamaepitys	12							
	Ranunculus arvensis	88	9						
	Legousia speculum-veneris	82	9						
	Consolida regalis	65	6						
DAss1	Chaenorrhinum minus	29		6					
	Valerianella locusta	24	14						
DAss1	Silene vulgaris	47	3	13				8	4
DAss1	Oxalis fontana	41		6					4
Diff. sp. order (<i>Centaureetalia</i>)									
	Centaurea cyanus	71							
	Raphanus raphanistrum	47							
	Buglossoides arvensis	41							
	Papaver argemone	24							
	Lolium temulentum	18							
	Rapistrum rugosum	12							
	Legousia hybrida	12							
DAss2	Anthemis arvensis	59	23						
DAss2	Veronica arvensis	53	31						
DAss2	Cerastium glomeratum	53	66						
	Aphanes arvensis	6	3						
	Arabidopsis thaliana	35	3						
	Papaver rhoeas	100	26					25	
	Fumaria officinalis	82	20					8	
DAss2	Alopecurus myosuroides	65	46					8	
	Myosotis arvensis	47	14						26
	Viola arvensis	47	6	6				8	
	Matricaria chamomilla	29			14				9
Char. and diff. (DAss2) ass. and all. (<i>Geranio-Allietum</i> and <i>Fumario-Euphorbion</i>)									
	Calepina irregularis	63							
DAss2	Cerastium tenoreanum	57							
	Muscari racemosum	34							
	Allium vineale	31							
	Ornithogalum umbellatum	23							
	Valerianella carinata	11							
	Geranium rotundifolium	3							
	Thlaspi arvense	3							
DAss2	Crepis taraxacifolia	24	83						
K, DAss2	Lactuca serriola	35	63						
DAss2	Geranium dissectum	6	60						
K, DAss2	Bromus sterilis	24	49						

Geranium molle	6	17				17	17	4
Brassica napus	12	20					8	9
Cardamine hirsuta	6	23	6					4
Geranium pusillum		3				17	17	4
Datura stramonium	12				40		8	
Oxalis corniculata	6		13					4
Geranium columbinum		23					8	13
Sherardia arvensis		3				17		4
Setaria viridis						67	42	26
Companions								
Agropyron repens	29	34	13	14	40	67	75	4
Calystegia sepium	94	51	88	57	100	33	42	65
Taraxacum officinale (aggr.)	12	94	63	57	60	67	67	26
Polygonum aviculare (aggr.)	76	3	25	14	60		58	9
Convolvulus arvensis	35	77	25	14		100	83	74
Artemisia vulgaris	24	11	6	43			8	4
Trifolium repens	24	14	31	14			8	17
Trifolium pratense	6	11	25			33	42	17
Daucus carota	29	14	13		20	17		13
Sorghum halepense	59		6	57	40	17	8	
Equisetum arvense	24		13	57	60		17	22
Plantago lanceolata	6	20	6			33	17	
Rumex crispus	18	54	19			17	17	
Plantago major	12	9	13		40		8	
Verbena officinalis	6		6	14	40		8	
Diplotaxis tenuifolia	6		6	14		33	17	
Rorippa sylvestris	29	6	6				8	
Potentilla reptans	18	11		14		33		
Galium mollugo	6	3				17		4
Lolium multiflorum	53	51					8	4
Rumex obtusifolius	18	40					17	4
Medicago lupulina	35		19			17	25	
Leucanthemum vulgare (aggr.)	6	3				17		9
Mentha spicata				14	20	17	8	
Arenaria serpyllifolia	59	20	6					
Arrhenatherum elatius	6	11						4
Glechoma hederacea	6		13					9
Medicago sativa	12	3					8	
Achillea millefolium	6					17		9
Mentha longifolia	71	29		14				
Silene alba	59	6						9
Cynodon dactylon		11	6			17		
Poa annua		46	13					17
Aristolochia clematitis		3				17	17	
Agrostis stolonifera		3	13			17		
Trifolium incarnatum /molinieri		20				17	25	
Leontodon hispidus		3				17	17	
Poa sylvicola	82	57						
Artemisia verlotorum	29	3						
Acalypha virginica			6	14				
Rubus caesius	18			14				
Prunella vulgaris						17		4
Conyza albida	24							
Ranunculus bulbosus		29						
Abutilon theophrasti					20			

Although this attribution is questionable, we wanted to put in evidence that this association lacks archaeophytic species, whereas species typical of weeded crops (*Lamium purpureum*, *Sonchus asper*, etc.) are frequent.

In addition to floristic criteria, we considered also structural (presence of bulbous geophytes), functional (presence of C4 plants) and chorological (presence of adventitious species) parameters for the definition of the coenoses.

Table 2 shows that geophytes reach their maximum value in *Geranio-Allietum*, while the highest values of neophytes and C4 species are found in *Echinochloo-Setarietum*; these two categories change together because most of successful neophytes are C4 plants. From Tab. 2 it is

also evident that the vineyard vegetation has a low level of disturbance, as indicated by the scarce participation of neophytes. The dendrogram obtained from the cluster analysis of all relevés (Fig. 1) shows four main clusters, as follows:

Table 2 - Parameters used in the definition of coenoses (A = *Papaveretum apuli*; B = *Geranio-Allietum*; C = *Echinochloo-Setarietum*; D = *Galeopsido-Galinsogetum*). Values are weighed by frequency.

	A	B	C	D
Bulbous geophytes	0.001	0.038	0	0
Neophytes	0.082	0.005	0.180	0.085
C4 species	0.026	0.010	0.215	0.058




Fig. 1 - Simplified dendrogram of 165 phytosociological relevés (similarity ratio, complete linkage). 1 = relevés of autumn-winter crops (*Papaveretum apuli*); 2 = relevés of summer weeded crops of hills, valley bottoms, and plains; 2a = *Echinochloo-Setarietum*, Friulian cool variant; 2b = *Echinochloo-Setarietum*, Friulian warm variant; 2c = *Echinochloo-Setarietum xanthetosum*; 2d = *Echinochloo-Setarietum*, *Sorghum halepense* agroform; 3a = *Galeopsido-Galinsogetum*; 3b = *Echinochloo-Setarietum*, Karstic cool variant; 3c = *Echinochloo-Setarietum*, Karstic warm variant; 4 = relevés of vineyards (*Geranium-Allietum*); 4a = *Geranio-Allietum*, typical form; 4b = *Geranio-Allietum*, *Taraxacum officinale* agroform; 4c = *Geranio-Allietum*, *Arrhenatherum elatius* agroform; 4d = *Geranio-Allietum*, *Agrostis stolonifera* agroform.

- cluster 1 = relevés of autumn-winter cereals (*Papaveretum apuli*);
 cluster 2 = relevés of weeded summer cultures of hills, valley bottoms and plain;
 cluster 3 = relevés of weeded summer cultures of Karst and mountains;
 cluster 4 = relevés of vineyards (*Geranio-Allietum*).

By following ecological and biogeographical considerations, relevés of cluster 2 can be ascribed to a Friuli sub-race of *Echinochloo-Setarietum*; these relevés can be further subdivided into a cool variant (cluster 2a), a warm variant (cluster 2b), a subassociation (cluster 2c, *xanthietosum*) and an agroform (cluster 2d). Similarly, relevés of cluster 3 can be ascribed to a Karstic sub-race of *Echinochloo-Setarietum*, with a warm variant (cluster 3c) and a cool one (cluster 3b), and to *Galeopsido-Galinsoghetum* (cluster 3a). Cluster 4 groups all relevés of vineyards.

The synthetic table of all relevés belonging to clusters 2a,b,c,d and 3a,b,c is reported in Tab. 3. This matrix has undergone a hierarchical classification based on the complete link method, in order to clarify the syntaxonomy of weeded crop fields. The dendrogram is reported in Fig. 2.




Fig. 2 – Dendrogram of weeded crop coenoses of Friuli – Venezia Giulia (similarity ratio, complete linkage). 1 = *Echinochloo-Setarietum*, Friulian cool variant; 2 = *Echinochloo-Setarietum*, Friulian warm variant; 3 = *Echinochloo-Setarietum xanthietosum*; 4 = *Echinochloo-Setarietum, Sorghum halepense* agroform; 5 = *Galeopsido-Galinsoghetum*; 6 = *Echinochloo-Setarietum*, Karstic cool variant; 7 = *Echinochloo-Setarietum*, Karstic warm variant.

Synoptic table of studied vegetation types

STELLARIETEA MEDIAE R. Tx., Lohm. et
 Preis. in R. Tx. 1950

CHENOPODIETALIA ALBI R. Tx. (1937) 1950

Veronico-Euphorbion Sissingh ex Passarge
 1964 (syntax. syn.: **Fumario-Euphor-**
bion T. Müller in Görs 1966)

- 1) *Geranio rotundifolii-Allietum vineale*
 R. Tx. ex von Rochow 1951 (syntax. syn.:
Cerastio-Geranium dissecti Poldini
 1980)

Panico-Setarion Sissingh in Westhoff *et al.*
 1946

- 2) *Echinochloo-Setarietum pumilae*
 Felföldy 1942 corr. Mucina 1993
 (syntax. syn.: *Panico-Polygonetum*
persicariae Pignatti 1953; *Hibisco-*
Digitarietum sanguinalis Poldini
 1980; *Hibisco-Sorghetum halepensis*
 Poldini 1989 non Horvatic et Hodak
 1960; *Oxalido - Chenopodietum*
 Sissingh 1950)
 subass. *xanthietosum italicum* subass.
 nova hoc loco

- 3) *Galeopsido tetrahit-Galinsoghetum*
parviflorae ass. nova hoc loco

CENTAUREETALIA CYANI R. Tx., Lohm. et
 Preis. in R. Tx. 1950

Caucalidion lappulae (R. Tx. 1950) von
 Rochow 1951

- 4) *Papaveretum apuli* ass. nova hoc
 loco

Associations

Ass.: *Geranio rotundifolii-Allietum vineale* R. Tx.
 ex von Rochow 1951 geogr. var. with
Cerastium tenoreanum, geogr. var. nova hoc
 loco (Tab. 4).

Syntax. syn.: *Cerastio-Geranium dissecti* Poldini
 1980.

“The vegetation of vineyards”

Tab. 3 - Simplified comparison table of weeded coenosis of Friuli-Venezia Giulia (frequency values). 1 = *Echinochloo-Setarium*, Friulian cool variant; 2 = *Echinochloo-Setarium*, Friulian warm variant; 3 = *Echinochloo-Setarium xanthetosum*; 4 = *Echinochloo-Setarium*, *Sorghum halepense* agroform; 5 = *Echinochloo-Setarium*, Karstic warm variant; 6 = *Echinochloo-Setarium*, Karstic cool variant; 7 = *Galeopsido-Galinsogetum*.

	1	2	3	4	5	6	7
Charac. and diff. (DAss1) species of ass. (<i>Echinochloo-Setarium</i>)							
K,DAss1 <i>Amaranthus retroflexus</i>	88	100	100	91	83	100	4
DAss1 <i>Solanum nigrum</i>	31	86	60	78	83	67	
<i>Portulaca oleracea</i>	19	71		5	33	8	
d2 <i>Hibiscum trionum</i>		14	40	5	100	17	
d2 <i>Bidens tripartita</i>		14	100	5			39
d2 <i>Amaranthus graecizans</i>		14			67	50	
d2 <i>Bidens frondosa</i>	6		60				
d2 <i>Polygonum lapathifolium</i>		43	100				
d2 <i>Xanthium italicum</i>		14	80				
d2 <i>Lycopus europaeus</i>			40				
Charac. and diff. (DAss2) species of ass. (<i>Galeopsido-Galinsogetum</i>)							
DAss2 <i>Galinsoga ciliata</i>	13			5			26
DAss2 <i>Galium aparine</i>						17	70
K, DAss2 <i>Mentha arvensis</i>						8	39
K, DAss2 <i>Stachys palustris</i>						25	39
DAss2 <i>Galeopsis speciosa</i>							48
DAss2 <i>Galeopsis tetrahit</i>							65
DAss2 <i>Aethusa cynapium</i>							30
Charac. and diff. (DAll) of all. (<i>Panico-Setarion</i>)							
K, DAll <i>Digitaria sanguinalis</i>	75	86	40	82	100	33	9
DO, DAll <i>Echinochloa crus-galli</i>	88	100	100	22	33	42	13
DAll <i>Setaria pumila</i>	100	43	100	48	83	50	48
<i>Chenopodium polyspermum</i>	63	14	40	8		50	30
DAll, DAss2 <i>Galinsoga parviflora</i>	100	29		73		42	74
<i>Panicum miliaceum</i>					17	8	4
Charac. and diff. (DO) species of ord. (<i>Chenopodietalia</i>)							
DO <i>Euphorbia helioscopia</i>	75	43	60	49	50	83	65
DO <i>Sonchus oleraceus</i>	63	43	60	22	67	58	74
DO <i>Sonchus asper</i>	19	14	60	13		50	
DO <i>Lamium purpureum</i>	44			9		33	57
<i>Diploaxis muralis</i>	6				83	42	
<i>Sonchus arvensis</i>	6					50	13
<i>Erodium cicutarium</i>					33	17	

Class species (*Stellarietea*)

Chenopodium album	100	100	100	73	100	92	100
Veronica persica	44	29	20	26	83	67	26
Cirsium arvense	13	29		13	67	58	43
Capsella bursa-pastoris	81	14		59	17	83	74
Fallopia convolvulus	38	14		5	33	83	83
Polygonum persicaria	81	100		91	33	83	96
Erigeron annuus	31	57	60	30			17
Conyza canadensis	6		20	8	17	8	
Stellaria media	56			21	33	67	96
Senecio vulgaris		29	20	23	33	42	
Vicia sativa (aggr.)			20	5	17	17	4
Mercurialis annua	6			40		42	4
Geranium molle				5	17	17	4
Setaria viridis				14	67	42	26
Sinapis arvensis				5	33	33	9
Datura stramonium			40	18		8	
Papaver rhoeas						25	
Myosotis arvensis							26

Companions

Agropyron repens	13	14	40	26	67	75	4
Calystegia sepium	88	57	100	87	33	42	65
Taraxacum officinale (aggr.)	63	57	60	22	67	67	26
Sorghum halepense	6	57	40	57	17	8	
Equisetum arvense	13	57	60	17		17	22
Polygonum aviculare	25	14	60	22		58	9
Convolvulus arvensis	25	14		52	100	83	74
Ranunculus repens	38		20	38	17	17	22
Diplotaxis tenuifolia	6	14		5	33	17	
Artemisia vulgaris	6	43		14		8	4
Trifolium repens	31	14		8		8	17
Daucus carota	13		20	9	17		13
Plantago major	13		40	5		8	
Mentha spicata		14	20	9	17	8	
Verbena officinalis	6	14	40			8	
Medicago lupulina	19			9	17	25	
Rumex crispus	19			5	17	17	
Trifolium pratense	25				33	42	17
Cynodon dactylon	6			27	17		
Plantago lanceolata	6				33	17	
Lythrum salicaria		14	20	5			
Potentilla reptans		14		9	33		
Trifolium incarnatum/molinieri					17	25	
Abutilon theophrasti			20				

Tab. 4 - *Geranio rotundifolii*-*Allietum vineale* R. Tx. ex von Rochow 1951 geogr. var. with *Cerastium tenoreanum* var. nova hoc loco

Relevé number		1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	
Altitude (x 10 m)		14	3	30	3	19	5	52	19	23	19	22	23	18	31	12	17	19	2	2	2	2	2	2	2	3	3	2	6	3	5	3	
Exposure								SO		NW			S															S		E			
Slope (°)								10		20			10															20		20			
Number of species (sporadic included)		28	21	25	27	34	17	25	35	25	28	28	23	32	17	31	48	42	41	31	33	30	27	26	23	19	26	32	24	26	25	29	
Charac. and diff. (DAss) spec. of ass. and all. (<i>Fumario-Euphorbion</i>)		typical form																															
DAss	Geranium dissectum										1	+	+	+	1		+	1	+	+	+	+	1	+	r		2	1	3	+			
DAss K	Lactuca serriola		+					1	+							+	1	+		+	+	+	+	1	+	+	+	+	+	+	+	+	
	Allium vineale								+	+		+					+	+	+				+	1	+	+	+	+	+	+	+	+	
DAss K	Bromus sterilis					2		1	2	2	3	1	3	3					2				1	+		+	2			+	+		
	Muscari racemosum				+				+		+						+		+												+		
DAss K	Veronica hederifolia	2	r	1				1	1	+		1	+	+					+		+					r			r				
DAss K	Veronica arvensis	+				+			+	+							+	+	r										1				
	Ornithogalum umbellatum										+				2		+	+		+		+						r					
DAss.	Fumaria officinalis	1					2										+		r	+													
DAss K	Anthemis arvensis															+	1		1					+				1	+			+	
DAss	Valerianella locusta																+		1						1			1	1				
DAss	Muscari comosum	+															+	+															
	Valerianella carinata								+	2												+	+										
	Thlaspi arvense				1																												
	Geranium rotundifolium (Gagea villosa)																																
Diff. spec. SE-alpine race																																	
DAss	Crepis taraxacifolia	+	+	1				+	+	+	1	1	1	1	1	+	+	1	+	r	+	+	+	+	+	+	+	+	+	+	+	+	
	Poa sylvicola	+			+				+							2	+	1	+	1	+	1	1	+	+	1	+	1	+	1	+	+	
	Rumex obtusifolius	+														1	+	+						+	r	+	+	+	+	+	+	1	
	Vicia sativa (aggr.)	+			+	+		1	1	1	+	+	+			+	1	+	1	+	+	+	+	+	+	1	1	1		+	r	+	
DAss	Cerastium glomeratum	+	+						+							+	+	+	+	+	+	1	+	+	+	1	1	1		+	1	1	
	Lolium multiflorum		1	1	2	1						1	1	1					1	1					+	+	+	+	+	+	+	+	
	Rumex crispus				1	+		1	1	1	1	1	2	1		1	1	+	1	1	1	1	+	1	+								
d	Calepina irregularis				2				3	1	3	3	3	3	2							2	3	2	3	1	2	3	2	2	3	1	2
DAss	Cerastium tenoreanum	+				1			+	+	+			+					+	2	1	+		+	+	+	1	+	+	1	+	+	
DAss	Alopecurus myosuroides					1											2	1		1		2	+	3	3	2	+	1		+	+	1	
Charac. and diff. (DO) spec. of ord. (<i>Chenopodietalia</i>)																																	
DO	Euphorbia helioscopia	1	+	+		+	+	1		+	+	+	+				1	1	1	1	1	+	2	2	1	1	+	1	1	1	+	2	
DO	Lamium purpureum		1					+	+	+		+	+				+		+				+				+						
DO	Sonchus oleraceus				+	+		+									+		r				1		+	+	+	+	+	+	+	+	
DO	Sonchus asper		r																+	+	+	+				+	+	+	1	+	+	+	
	Erodium cicutarium	+	+	+	1			+				+										+	+	+			1	+	1		1	+	1
	Diplotaxis muralis	+	+		+									+								+	+	+							+		
	Lamium amplexicaule						+	2			+	+												+									
	Echinochloa crus-galli											1					1	1															
	Digitaria sanguinalis																1					1											
	Amaranthus retroflexus																										+	+					+
	Sonchus arvensis																	+															
Class spec. (<i>Stellarietea</i>)																																	
	Veronica persica	3	3	1	3	3	2	1	3	3	3	3	3	2	2	2	2	1	1	1	1	2	+	2	1	1	2	3	2	2	1	2	1
	Stellaria media	3	2	+	4	2	3	4	3	3	2	3	3	2	3	2	2	1	+	1	2	2	1	1	+		+	1	2	1	+	+	+
	Senecio vulgaris	+	r	+	+	1	1	+	+	+	+	+	+	+	+	+	+	1	+	r	1	+	2	+	r	r		+	+	+	+	+	+
	Cirsium arvense			2	2	1	2	1	1				1	1	1	2	1	2	1	2	1	2	+	1	1		1		1	1	+	+	+
	Capsella bursa-pastoris	+	1	+	+	1	1	1	+		+		+	+	1	1	+	+	r	+	2	+						+	+	+	+	+	+
	Erigeron annuus		+			+	+										+	+		+	+	+	+				+	1	+	+	+	+	+
	Cardamine hirsuta			1							+	+	+								+	+	+		r								
	Papaver rhoeas	+															+	+	r			+	+									r	
	Geranium molle			r																1							r		+				
	Geranium columbinum					+	+		+	+	+						+																
	Brassica napus	+	+																			+			+		+	r				+	+
	Mercurialis annua					+											+															+	+
	Bromus hordeaceus								+					+									+										+
	Myosotis arvensis			+	+	+					+															+							
	Fallopia convolvulus	+				+	+	+				+																					
	Anagallis arvensis	+											+				+	+				+											
	Conyza canadensis					+	+										+																

- Segetal vegetation in Friuli-Venezia Giulia -

Tab. 4 – Continued.

		26		30		32		33		34		35		36		37		38		39		40		41		42		43		44		45		46		47		48		49		50		51		52		53		54		55		56		57		58		59		60		61		62			
		5		15		33		15		13		2		2		24		2		2		2		24		2		2		24		3		12		13		13		11		12		13		5		2		2		1		2		1		26		25		24		2		2		2	
		26		30		32		33		34		35		36		37		38		39		40		41		42		43		44		45		46		47		48		49		50		51		52		53		54		55		56		57		58		59		60		61		62			
		26		30		32		33		34		35		36		37		38		39		40		41		42		43		44		45		46		47		48		49		50		51		52		53		54		55		56		57		58		59		60		61		62			
		26		30		32		33		34		35		36		37		38		39		40		41		42		43		44		45		46		47		48		49		50		51		52		53		54		55		56		57		58		59		60		61		62			
		26		30		32		33		34		35		36		37		38		39		40		41		42		43		44		45		46		47		48		49		50		51		52		53		54		55		56		57		58		59		60		61		62			
		26		30		32		33		34		35		36		37		38		39		40		41		42		43		44		45		46		47		48		49		50		51		52		53		54		55		56		57		58		59		60		61		62			
		26		30		32		33		34		35		36		37		38		39		40		41		42		43		44		45		46		47		48		49		50		51		52		53		54		55		56		57		58		59		60		61		62			
		26		30		32		33		34		35		36		37		38		39		40		41		42		43		44		45		46		47		48		49		50		51		52		53		54		55		56		57		58		59		60		61		62			
		26		30		32		33		34		35		36		37		38		39		40		41		42		43		44		45		46		47		48		49		50		51		52		53		54		55		56		57		58		59		60		61		62			
		26		30		32		33		34		35		36		37		38		39		40		41		42		43		44		45		46		47		48		49		50		51		52		53		54		55		56		57		58		59		60		61		62			
		26		30		32		33		34		35		36		37		38		39		40		41		42		43		44		45		46		47		48		49		50		51		52		53		54		55		56		57		58		59		60		61		62			
		26		30		32		33		34		35		36		37		38		39		40		41		42		43		44		45		46		47		48		49		50		51		52		53		54		55		56		57		58		59		60		61		62			
		26		30		32		33		34		35		36		37		38		39		40		41		42		43		44		45		46		47		48		49		50		51		52		53		54		55		56		57		58		59		60		61		62			
		26		30		32		33		34		35		36		37		38		39		40		41		42		43		44		45		46		47		48		49		50		51		52		53		54		55		56		57		58		59		60		61		62			
		26		30		32		33		34		35		36		37		38		39		40		41		42		43		44		45		46		47		48		49		50		51		52		53		54		55		56		57		58		59		60		61		62			
		26		30		32		33		34		35		36		37		38		39		40		41		42		43		44		45		46		47		48		49		50		51		52		53		54		55		56		57		58		59		60		61		62			
		26		30		32		33		34		35		36		37		38		39		40		41		42		43		44		45		46		47		48		49		50		51		52		53		54		55		56		57		58		59		60		61		62			
		26		30		32		33		34		35		36		37		38		39		40		41		42		43		44		45		46		47		48		49		50		51		52		53		54		55		56		57		58		59		60		61		62			
		26		30		32		33		34		35		36		37		38		39		40		41		42		43		44		45		46		47		48		49		50		51		52		53		54		55		56		57		58		59		60		61		62			
		26		30		32		33		34		35		36		37		38		39		40		41		42		43		44		45		46		47		48		49		50		51		52		53		54		55		56		57		58		59		60		61		62			
		26		30		32		33		34		35		36		37		38		39		40		41		42		43		44		45		46		47		48		49		50		51		52		53		54		55		56		57		58		59		60		61		62			
		26		30		32		33		34		35		36		37		38		39		40		41		42		43		44		45		46		47		48		49		50		51		52		53		54		55		56		57		58		59		60		61		62			
		26		30		32		33		34		35		36		37		38		39		40		41		42		43		44		45		46		47		48		49		50		51		52		53		54		55		56		57		58		59		60		61		62			
		26		30		32		33		34		35		36		37		38		39		40		41		42		43		44		45		46		47		48		49		50		51		52		53		54		55		56		57		58		59		60		61		62			
		26		30		32		33		34		35		36		37		38		39		40		41		42		43		44		45		46		47		48		49		50		51		52		53		54		55		56		57		58		59		60		61		62			
		26		30		32		33		34		35		36		37		38		39		40		41		42		43		44		45		46		47		48		49		50		51		52		53		54		55		56		57		58		59		60		61		62			
		26		30		32		33		34		35		36		37		38		39		40		41		42		43		44		45		46		47		48		49		50		51		52		53		54		55		56		57		58		59		60		61		62			
		26		30		32		33		34		35		36		37		38		39		40		41		42		43		44		45		46		47		48		49		50		51		52		53		54		55		56		57		58		59		60		61		62			
		26		30		32		33		34		35		36		37		38		39		40		41		42		43		44		45		46		47		48		49		50		51		52		53		54		55		56		57		58		59		60		61		62			
		26		30		32		33		34		35		36		37		38		39		40		41		42		43		44		45		46		47		48		49		50		51		52		53		54		55		56		57		58		59		60		61		62			
		26		30		32		33		34		35		36		37		38		39		40		41		42		43		44		45		46		47		48		49		50		51		52		53		54		55		56		57		58		59		60		61		62			
		26		30		32		33		34		35		36		37		38		39		40		41		42		43		44		45		46		47		48		49		50		51		52		53		54		55		56		57		58		59		60		61		62			
		26		30		32		33		34		35		36		37		38		39		40		41		42		43		44		45		46		47		48		49		50		51		52		53		54		55		56		57		58		59		60		61		62			
		26		30		32		33		34		35		36		37		38		39		40		41		42		43		44		45		46		47		48		49		50		51		52		53		54		55		56		57		58		59		60		61		62			
		26		30		32		33		34		35		36		37		38		39		40		41		42		43		44		45		46		47		48		49		50		51		52		53		54		55		56		57		58		59		60		61		62			
		26		30		32		33		34		35		36		37		38		39																																																	

[illegible]

Floristic composition. This is the agrocoenosis with the highest number of bulbous geophytes (*Allium*

Syntaxonomy. The vegetation of vineyards of Karst and Collio was described by Poldini (1980, 1989) with the name of *Cerastio-Geranium dissecti*. At that time the author had already put in evidence the strong differentiation from the isoecic coenosis of

Tab. 4 – Continued.

[illegible]

Central Europe (*Geranio-Allietum*). In this respect, we faced the problem of differentiating between geographical race and association. Data analysis is not useful to detect the shifting point between these two *syntaxa*, and it is therefore necessary to evaluate both the biogeography and the potential vegetation of the survey area (Poldini *et al.* 1996). The potential vegetation (as synonym of the synphytosociological term «head association») of Friuli-Venezia Giulia is represented by mature oaks woods: *Seslerio-Quercetum petraeae* (Poldini 1964 n.n.) Poldini 1982, *Carici umbrosae-Quercetum petraeae* Poldini ex Marincek 1994, *Asparago tenuifolii-Quercetum roboris* (Lausi 1966) Marincek 1994 (basonym: *Quercus-Carpinetum boreo-italicum* Pignatti ex Lausi 1966). These woods are particularly rich in

species with submediterranean - SE European distribution. Such a strong characterization could suggest that the coenosis of Friuli-Venezia Giulia vineyards belongs to an independent association. Nevertheless, we observed that, when anthropogenic pressure is very high, the same association («tail association») develops from different head associations (Géhu 1996).

Table 5 shows the main differences between the vineyard vegetation of Friuli-Venezia Giulia and that of Central Europe. The community occurring in Friuli-Venezia Giulia is characterized by several thermophilous elements, such as *Alopecurus myosuroides*, *Calepina irregularis*, *Calystegia sepium*, *Cerastium tenoreum*, *Crepis taraxacifolia*, *Geranium dissectum*, *Poa sylvicola*, *Rumex*

Tab. 5 – Simplified, synthetic table of Centro-European *Geranio-Allietum* and of the SE-Alpine race (frequency values). A = data from Oberdorfer (1993); B = data from Wilmanns (1989); C = data from Tab. 4.

	A	B	C		A	B	C
Species of ass. and all. (<i>Veronico-Euphorbion</i>)							
<i>Allium vineale</i>	70	75	40	<i>Capsella bursa-pastoris</i>	25	13	75
<i>Bromus sterilis</i>	20	50	53	<i>Chenopodium album</i>	35	19	10
<i>Fumaria officinalis</i>	54	25	18	<i>Cirsium arvense</i>	56	38	68
<i>Geranium rotundifolium</i>	21	44	8	<i>Conyza canadensis</i>	13	19	8
<i>Lactuca serriola</i>	22	75	60	<i>Geranium pusillum</i>	11	13	3
<i>Muscari racemosum</i>	51	56	35	<i>Mercurialis annua</i>	52	81	23
<i>Ornithogalum umbellatum</i>	23	31	28	<i>Papaver rhoeas</i>	9	6	28
<i>Valerianella carinata</i>	2	13	10	<i>Senecio vulgaris</i>	84	56	85
<i>Veronica arvensis</i>	13	25	28	<i>Stellaria media</i>	80	88	98
<i>Cerastium glomeratum</i>	4		60	<i>Veronica persica</i>	64	94	100
<i>Gagea villosa</i>	16	13		<i>Anagallis arvensis</i>	16		18
<i>Geranium dissectum</i>	17		63	<i>Fallopia convolvulus</i>	1		18
<i>Thlaspi arvense</i>	6		5	<i>Geranium columbinum</i>	15		20
<i>Veronica hederifolia</i>		88	33	<i>Malva neglecta</i>	17		5
<i>Allium oleraceum</i>	14			<i>Setaria viridis</i>	21		8
<i>Allium scordoprasum</i>		13		<i>Sinapis arvensis</i>	20		5
<i>Alopecurus myosuroides</i>			40	<i>Cardamine hirsuta</i>		31	20
<i>Anthemis arvensis</i>			20	<i>Vicia sativa</i> (aggr.)		19	75
<i>Calepina irregularis</i>			55	<i>Brassica napus</i>			18
<i>Cerastium tenoreum</i>			58	<i>Diploaxis muralis</i>			25
<i>Crepis taraxacifolia</i>			78	<i>Erigeron annuus</i>			33
<i>Lolium multiflorum</i>			45				
<i>Muscari comosum</i>			10	Companions			
<i>Poa sylvicola</i>			50	<i>Agropyron repens</i>	41	25	38
<i>Valerianella locusta</i>			13	<i>Convolvulus arvensis</i>	41	69	80
				<i>Galium aparine</i>	10	25	8
Ord. (<i>Chenopodietalia</i>) species				<i>Lolium perenne</i>	4	25	5
<i>Euphorbia helioscopia</i>	78	19	80	<i>Poa annua</i>	22	69	40
<i>Lamium amplexicaule</i>	30	6	18	<i>Polygonum aviculare</i> (aggr.)	15	19	3
<i>Lamium purpureum</i>	89	100	33	<i>Taraxacum officinale</i>	91	94	88
<i>Solanum nigrum</i>	61	13	10	<i>Diploaxis tenuifolia</i>	1	25	
<i>Sonchus arvensis</i>	15	6	5	<i>Lapsana communis</i>	14		3
<i>Sonchus asper</i>	81	19	38	<i>Ranunculus repens</i>	26		13
<i>Sonchus oleraceus</i>	91	19	48	<i>Veronica polita</i>	58	6	
<i>Echinochloa crus-galli</i>	27		10	<i>Calystegia sepium</i>			48
<i>Erodium cicutarium</i>	29		40	<i>Mentha longifolia</i>			25
<i>Euphorbia peplus</i>	33			<i>Ranunculus bulbosus</i>			25
				<i>Rumex crispus</i>			48
Class (<i>Stellarietea</i>) species				<i>Rumex obtusifolius</i>			35
<i>Amaranthus retroflexus</i>	19	19	15	<i>Trifolium incarnatum/molineri</i>			23

crispus, and *Vicia sativa* / *angustifolia*. With the exception of *Calepina irregularis*, these species are not exclusive of vineyards, but penetrate from surrounding habitats. They can be considered as the geographical differentials of a Southeastern Alpine race of *Geranio-Allietum*. The ecological-structural meaning of the high frequency of spring bulbous geophytes, which distinguishes this agrocoenosis from the others, has been considered particularly important and has led us to include the relevés of Friuli-Venezia Giulia into *Geranio-Allietum*.

At higher syntaxonomical levels, the attribution of this coenosis is evident; *Geranium dissectum* and *Fumaria officinalis* show adherence to the alliance *Veronico-Euphorbion*. Since *Geranio-Allietum* is the only known association of this alliance in Friuli-Venezia Giulia, alliance and association species coincide. *Sonchus oleraceus*, *S. asper*, *Euphorbia helioscopia*, *Lamium purpureum*, *L. amplexicaule*, *Diplotaxis muralis* and *Erodium cicutarium* show adherence to *Chenopodietalia*.

Centaureetalia species are particularly frequent in these relevés. This depends on the growth time of *Geranio-Allietum* (spring) when also the autumn-winter cultures have their maximum blooming. *Stellarietea* species are very numerous. Among them, *Veronica persica*, *Stellaria media*, *Senecio vulgaris*, *Cirsium arvense*, *Capsella bursa-pastoris* and *Erigeron annuus* are the most frequent.

Variability. It is low, although the relevés come from areas with strong pedological differences (karstic «terre rosse», flysch and gravels). A certain differentiation (floristic rather than ecological) exists between relevés of the Collio hills near Gorizia (*Alopecurus myosuroides*) and those from the Karst plateau near Trieste (*Ranunculus bulbosus* and *Trifolium incarnatum/molinieri*, maybe once used as fodder plants).

Agroforms. The variety of cultivation techniques in viticulture has a strong influence on vineyard coenoses. It is therefore possible to identify three agroforms that are often more frequent than the typical association.

— *Taraxacum officinale* agroform (Tab. 4, rels. 36-44): it occurs in vineyards characterized by light tilling once or twice a year, combined with mulching between the rows and the use of desiccatives beneath the rows. Sometimes *Taraxacum officinale* dominates these vineyards, because its roots, rich of reserve material and buds, make it resistant to mulching and tolerant to the use of desiccatives and to occasional tap-root fragmentation caused by

harrowing, which may even foster its propagation (Wilmanns 1991). In this agroform all species of *Geranio-Allietum* are well represented, especially *Geranium dissectum*.

— *Agrostis stolonifera* agroform: it occurs in vineyards characterized by the absence of tilling (Tab. 4, rels. 52-62); it spreads on heavy clay soils with high water table, such as those of the springs area («risorgive»). The weeds control is made by mulching, which helps the spreading of creeping perennial herbs (*Potentilla reptans*, *Agrostis stolonifera* and *Ranunculus repens*, etc.) or of plants with tap-roots (*Taraxacum officinale*). Among the differential species of *Geranio-Allietum*, *Allium vineale* reaches high cover values, because, in absence of tilling, it forms large tufts.

— *Arrhenatherum elatius* agroform (Tab. 4, rels. 45-51): it occurs in vineyards where soil tilling is replaced by mowing and natural manuring. Besides *Arrhenatherum elatius*, other species of *Arrhenatherion* dominate this agroform, such as *Galium mollugo*, *Silene alba* and *Plantago lanceolata*. Therefore, the characteristic species of vineyards almost disappear, being replaced by perennial herbs.

The two first agroforms correspond to those described by Wilmanns (1989, 1991) from the Rhine valley.

Life forms. This association, in its typical aspect, shows a high percentage of geophytes (11.3%), although therophytes dominate (67.0%) and the percentage of hemicryptophytes is also high (21.4%). The increase in hemicryptophytes in the agroforms is caused by the reduction of soil tilling, replaced by mowing and the use of desiccatives.

Chorotypes. The low percentage of adventitious species is noteworthy; this fact confirms that vineyards are scarcely disturbed cultures. The high percentage of euromediterranean species (26.6%) and the presence of stenomediterranean ones (0.6%) well arrange with the Mediterranean gravitation of the SE-Alpine race of *Geranio-Allietum*; these chorotypes decrease in agroforms where species with European distribution (more resistant to disturbance) prevail.

Pollination and dispersal. There are some differences between the typical association and its agroforms. Autogamy and entomophily are progressively substituted by anemophily, proceeding from the association to its agroforms. In *Geranio-Allietum*, due to ground tilling, autogamy is very important; in fact this strategy guarantees quick self-fecundation of species (Faegri & van der Pijl 1971)

and fructification before the next tilling. In the agroforms with *Agrostis stolonifera* and *Arrhenatherum elatius* the mowing favours the spreading of anemophilous *Gramineae*. The agroform with *Taraxacum officinale* is a transition coenosis, because of the co-presence of tilling and mulching.

Sinecology. The coenosis develops at the beginning of spring in vineyards where artisan cultivation techniques are used. The absence of tilling until late spring is essential for the survival of the coenosis, which would be destroyed by early tilling, because it is formed mainly by spring flowering plants. The light ploughing at the beginning of June determines the closure of the seasonal cycle of the association. During summer harrowing is carried out to keep under control the summer species of *Stellarietea* (especially *Setaria viridis*, *Erigeron annuus*, *Amaranthus retroflexus* and *Mercurialis annua*).

This association develops on «terre rosse», silts, clays, and gravels; these soils are not heavily modified by viticulture, since it needs only light manuring.

The altitude ranges from sea level to 350-400 m (Collio hills and Karst plateau).

Synchorology. The presence of *Geranio-Allietum* has been reported in Germany, Austria, North-Eastern Italy and Slovenia.

Ass.: *Echinochloo-Setarietum pumilae* Felföldy 1942 corr. Mucina 1993 geogr. var. with *Calystegia sepium*, geogr. var. nova hoc loco (Tab. 6).

Syntax. syn.: *Panico-Polygonetum persicariae* Pignatti 1953, *Hibisco-Digitarietum sanguinalis* Poldini 1980, *Hibisco-Sorghetum halepensis* Poldini 1989 non Horvatic et Hodak 1960; *Oxalido-Chenopodietum* Sissingh 1950

“The vegetation of weeded summer cultures of the plain and of the hill belts”

Charact. and diff. (D) species: *Amaranthus retroflexus* (D), *Solanum nigrum* (D), *Portulaca oleracea* and *Calystegia sepium*.

Floristic composition. This association is dominated by *Setaria pumila*, *Digitaria sanguinalis*, *Echinochloa crus-galli*, *Amaranthus retroflexus*, *Polygonum persicaria*, *Euphorbia helioscopia*, *Chenopodium album*, *C. polyspermum*, *Convolvulus arvensis* and *Sonchus oleraceus*. *Portulaca oleracea*, *Solanum nigrum*, *Capsella bursa-pastoris* and *Taraxacum officinale* are often present, but with low cover values.

Syntaxonomy. In the attempt to attribute the relevés

of Friuli-Venezia Giulia, we first considered two associations already described for weeded crops: *Panico-Polygonetum* Pign. 1953 (Pignatti 1952), surveyed in the eastern Venetian plain, and *Echinochloo-Setarietum pumilae* Felföldy 1942 corr. Mucina 1993, present in Central Europe (Mucina 1993), and we compared synthetic tables of weeded cultures of northern Italy (Gerdol & Piccoli 1979; Lorenzoni 1963; Poldini 1989) with those of Central Europe (Krippelová 1981; Regula-Bevilacqua 1979; Seljak 1989; Oberdorfer 1993) by means of numerical analysis. The classification (Fig. 3) shows that the relevés from Ferrara separate at a high level (0.63 of dissimilarity), while the cluster of relevés from the Venetian plain join the group of associations of Central Europe at a lower level of similarity (0.54). Hence all the examined relevés have been gathered into a single coenosis. Due to nomenclatural priority, the correct name is *Echinochloo-Setarietum*, whereas the binomial *Panico-Polygonetum* and *Oxalido-Chenopodietum* are syntaxonomic synonyms.




Fig. 3 – Dendrogram of weeded crops of Europe, obtained by classification of synthetic tables (similarity ratio, mean linkage). 1 = *Panico-Polygonetum* (Italy, Ferrara, Gerdol & Piccoli 1979); 2 = *Setario-Galinsotum* (Croatia, Regula-Bevilacqua 1979); 3 = *Setario-Galinsotum* (Slovakia, Krippelová 1981); 4 = *Oxalido-Chenopodietum* (Slovakia, Krippelová 1981); 5 = *Panico-Polygonetum* (Italy, Eastern Venetian plain, Lorenzoni 1963); 6 = *Oxalido-Chenopodietum* (Italy, Eastern Venetian Plain, Lorenzoni 1963); 7 = *Setario-Galinsotum* (Slovenia, Seljak 1989); 8 = *Oxalido-Chenopodietum* (Slovenia, Seljak 1989); 9 = *Setario-Galinsotum* (Germany, Oberdorfer 1993); 10 = relevés from Friuli – Venezia Giulia.

Although the eastern Venetian plain and Central Europe belong to different biogeographical regions with different «head associations» (see Syntaxonomy of *Geranio-Allietum*), it is also true that coenoses of weeded cultures are subject to a levelling anthropogenic pressure.

The attribution of this association to the *Panico-Setarion* alliance is due to the high participation of species of the tribe *Panicoideae* (*Setaria pumila*, *Digitaria sanguinalis* and *Echinochloa crus-galli*), of *Galinsoga parviflora* and *Chenopodium polyspermum*. The framing in *Chenopodietalia* is justified by the presence of *Sonchus oleraceus*, *S. asper*, *Lamium purpureum* and *Euphorbia helioscopia*. *Chenopodium album*, *Capsella bursa-pastoris*, *Erigeron annuus*, *Stellaria media*, *Fallopia convolvulus*, *Veronica persica* and *Setaria viridis* show adherence to the *Stellarietea* class.

Variability. Because of its very large geographical distribution, *Echinochloo-Setarietum* exhibits a certain variability. In Friuli-Venezia Giulia a Southeastern-Alpine race is present, which is characterized by some thermophilous species such as *Calystegia sepium*, *Euphorbia helioscopia*, *Polygonum persicaria* and *Chenopodium polyspermum*. This SE-Alpine race is not homogenous; on the basis of biogeographical considerations, it is possible to segregate two sub-races (Fig. 4):

a) Karstic sub-race (Tab. 6, rels. 46-63), distinguished by: *Veronica persica*, *Agropyron repens*, *Cirsium arvense*, *Amaranthus graecizans*, *Diplotaxis muralis*, *Erodium cicutarium*, *Trifolium incarnatum/molineri* and *Stachys palustris*;

b) Friuli sub-race (Tab. 6, rels. 1-23), distinguished by the absence, or the negligible presence, of Karstic differential species.

Within the two sub-races, parallel ecological responses to changes in temperature and in soil moisture can be observed. We can therefore identify a cool variant and a warm one of each subrace. The cool variants are differentiated by *Galinsoga parviflora*, *Stellaria media*, *Fallopia convolvulus*, *Lamium purpureum* and *Trifolium pratense*, while the warm ones by *Sorghum halepense* and *Hibiscus trionum*. In the Karst the cool variant is found at the bottom of dolines, while the warm one is found in the south-exposed terraced cultures; the warm variant has already been described as a race of *Hibisco-Sorghetum halepensis* Horvatic et Hodak 1960 by Poldini (1989), who framed it in the Mediterranean alliance *Diplotaxidion* Br.-Bl. 1936. According to the new results, this alliance is excluded from Friuli-Venezia Giulia. In Friuli, the cool variant is found at the bottom of pre-Alpine valleys and on the Friulian hills, while the warm one is found in the plain.

Subassociation. One new subassociation has been detected:

xanthietosum italicum subass. nova, holotypus Tab. 6 rel. 24 (Tab. 6, rels. 24-28). This subassociation has been recorded in the crop fields of reclaimed areas of Friuli, where the soil texture limits the water flow, increasing the soil moisture. It is differentiated by *Polygonum lapathifolium* (which completely replaces *P. persicaria*), *Bidens tripartita*, *B. frondosa*, *Xanthium italicum* and *Lycopus europaeus*. These




Fig. 4 – Variability of *Echinochloo-Setarietum*.

Tab. 6 – *Echinochloa-Setarietum pumilae* Felföldy 1942 corr. Mucina 1993 geogr. var. with *Calystegia sepium* var. nova hoc loco.

Relevés n°	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	
Altitude (x10m)	40	37	35	40	30	30	39	43	35	2	5	6	34	34	28	26	3	1	1	1	1	11	2	1	25	1	30		
Total n. of species (spor. incl.)	18	21	23	18	20	18	19	15	22	9	13	27	13	19	14	21	19	11	19	10	17	19	16	19	21	16	22	17	
Char. & diff. (DAss) sp. of ass.	Friuli sub-race																xanthietosum												
	cool variant								warm variant																				
DAss	Amaranthus retroflexus	1	+	1	+	1	2	1	1	1	1	2	2	2	2	2	2	2	1	1	1	1	1	1	1	+	+	+	
DAss	Calystegia sepium	2	2	+	2	1	+	+	1	+	+	+	+	1	1	+	2	+	1	1	1	+	+	1	+	1	1	1	
DAss	Solanum nigrum				+			r	+		+						+	1	1	1	1	+	+	1	1	1	1	1	
	Portulaca oleracea								3	1	+						+	1	1		+	r							
Char. & diff. (DAll) sp. alliance																													
DAll	Digitaria sanguinalis	+	+		2	1	3	+	r	1	1	1	1	1	1	+	2	1	1		1	1	1		+	+			
DAll	Setaria pumila	2	2	1	2	1	+	2	2	2	+	1	1	1	1	2	1					1	+	1	1	1	1	1	
DAll	Echinochloa crus-galli	+	1			+	+	2	2	4	+	+	2	2	+	1	1	2	2	2	1	2	1	1	1	1	+	2	2
DAll	Panicum miliaceum																												
Diff. sp. SE-alpine race																													
K	Polygonum persicaria	+	1	+	+	1	1	2	1	1		1	1	+		+	r	1	2	2	2	2	1						
DO	Euphorbia helioscopia	1	1	+	1	+		1	+	r				+	+	+	+		+						+	+		+	
DAll	Chenopodium polyspermum	2	2	1	+	+	1	2	2	+	+						+								+			1	
Diff. sp. Karst subrace																													
K	Veronica persica	+	1	+	+				+				+			+	+	+	+						r		+		
	Agropyron repens															+	+										+	+	
K	Cirsium arvense				1								r									+	+						
K	Setaria viridis																												
	Amaranthus graecizans																	+											
O	Diploaxis muralis			+																									
O	Erodium cicutarium																												
	Trifolium incarnatum\ molinieri																												
	Stachys palustris																												
Diff. sp. warm variant																													
	Sorghum halepense																1			+	+	+	+				+	+	
	Hibiscus trionum																	+							+	+			
Diff. sp. cool variant																													
DAll	Galinsoga parviflora	3	3	4	2	2	4	3	3	3	4	2	2	1	3	1	1	+	+										
K	Stellaria media	2	1	2	1	2	2	3	1								1												
K	Fallopia convolvulus	+	1	2			+	1	+									r											
DO	Lamium purpureum	+	2	+	1	+	1							r															
	Trifolium pratense			+				+	+			r																	
Diff. sp. subass. (xanthietosum)																													
K	Polygonum lapathifolium																	r	+				r		1	1	+	1	2
	Bidens tripartita																							1	1	2	2	2	2
K	Xanthium italicum																							1	+	+	+	1	
	Bidens frondosa			+																						+	+	+	1
	Lycopus europaeus																									+	+		
Char. & diff. (DO) sp. Chenopodietaia																													
DO	Sonchus oleraceus	2	2	1	+		r	1	2	1				+		+	+	+	+							+	+	+	+
DO	Sonchus asper						r			+						r			r							+	+		r
	Sonchus arvensis															r													
	Galinsoga ciliata	2															r												
	Euphorbia peplus												r																
	Lamium amplexicaule																												
Class species																													
	Chenopodium album	1	2	2	2	1	1	3	2	1	1	1	1	+	2	1	1	3	2	1	1	1	+	+	+	1	1	+	1
	Capsella bursa-pastoris	1	1	+			+	+	1	+		+	+	+	+	+	1												
	Erigeron annuus			+		+												+											
	Senecio vulgaris																	r		r									r
	Mercurialis annua												+																
	Sinapis arvensis																												
	Datura stramonium																										+	+	
	Conyza canadensis																												+
	Vicia sativa (aggr.)																												+
	Anagallis arvensis												+																
	Geranium molle																												
	Oxalis fontana												+																+
	Geranium pusillum																												
	Cardamine hirsuta						r																						
	Oxalis corniculata			+				+																					
	Viola tricolor			r					+																				
	Brassica napus																												
	Acalypha virginica												1											+					
Companions																													
	Taraxacum officinale	+	+	+		+	+	1	+			+	+	+	+	+	+	+	+			r	r	+	+	+	+	+	
	Convolvulus arvensis				1		+		1	+																			
	Polygonum aviculare							1					r	+	+										+	+		1	1
	Ranunculus repens				+	1	+		1				+	+														1	1
	Equisetum arvense					r											+									1	1	1	1
	Cynodon dactylon												r																

Tab. 6 - Continued.

[illegible]

species, except *Polygonum lapathifolium* (typical of cultures), come from neighbouring humid, nitrophilous habitats. This subassociation is similar to the *Polygono-Xanthietum italicum* Pirola & Rossetti 1974 which has been described from the eutrophic, disturbed gravel-bed of the Reno river, near Bologna (Pirola & Rossetti 1974), and to *Polygono-Bidentetum* (W. Koch 1926) Lohm. 1950, a ruderal association of *Bidentetea tripartiti* Tx., Lohm. et Prsg. 1950 (Regula-Bevilacqua 1979).

Agroform. One agroform has been defined:

— *Sorghum halepense* agroform (Tab. 6, rels. 29-45): it is typical of intensive cultivation with use of heavy ploughing and high input of herbicides and of mineral manure. The relevés are characterized by very low cover and strong quantitative modification of the interspecific relationships (Ferrari *et al.* 1979). The permanence of association differential species assures the link with the typical coenosis.

Life forms and chorotypes. Therophytes prevail, and, in the Friuli subrace, they reach the highest value among the studied coenoses (73.1%). Geophytes are scarce; only in the *Sorghum halepense* agroform (15.1%) and in the Karst subrace (14.8%) their value is significant, due to the presence of *Sorghum halepense* (Friuli) and of *Agropyron repens* (Karst).

The most frequent chorotypes are Cosmopolitan, adventitious and palaeotemperate.

Synecology. *Echinochloa-Setarietum* is present in weeded summer cultivation, such as maize, soy-bean, and, to a lesser degree, potatoes. After sowing, these cultures undergo weeding and hoeing, but nowadays weeds are mainly controlled by the use of herbicides.

Nowadays the maize cultures can produce a biomass of 450-500 q ha⁻¹ in 100-200 days. Maize exploits at best the modern techniques of manuring, irrigation, etc., but it is very sensitive to weed competition. This is due to its inability to recover from damages caused by the development of weeds, because of its short growth season. Moreover, it is the only cereal sowed in rows, and for this reason it is not able to express the typical spatial competitiveness of *Gramineae*. In Friuli-Venezia Giulia, this susceptibility is enhanced by low spring temperatures, which delay the early growth stages. Nevertheless, maize cultivation has become an easy task, thanks to the use of triazinic herbicides (Cassani 1986). Their intensive and extensive use have selected some resistant weeds. As we can observe from the phytosociological relevés, in this

particular ecosystem some species are now very frequent and widespread, as they have become resistant (*Amaranthus retroflexus*, *Chenopodium album* and *Solanum nigrum*) or tolerant (monocots such as *Echinochloa crus-galli*, *Digitaria sanguinalis* and *Sorghum halepense*) to triazinic compounds (Pollastro 1989). Infestation by these species is very dangerous to maize, because most of them belong to the same family (*Gramineae*) and are therefore very competitive. Moreover, except *Chenopodium album* and *Solanum nigrum*, they are C4 plants (as maize). Under favourable environmental conditions, C4 species are known to be very competitive against C3 species (Baker 1972, Collins & Jones 1985). Moreover, since triazinic substances operate on photosynthesis, C4 weeds take further advantage of their alternative CO₂ assimilation pathway, and can survive to these herbicides (Gutierrez *et al.* 1974).

Indeed, an overgrowth of weeds soon after sowing would seriously hamper the development of any crop. From an agronomic point of view, the damage is greater when the weed coenosis develops at the same time as the crop plant. Conversely, the growth of weeds at the end of July, when maize and soy-bean dominate, is usually tolerated and the crop coenosis consequently reaches its maximum development only in late summer (second half of August). It is then destroyed by the harvest in late August-early September (potatoes) or at the end of October-beginning of November (maize and soy-bean). This tilling calendar allows most of the weeds to fructify, thus increasing the soil seed-bank responsible for the spring renovation of the coenosis. Fragmented roots and underground organs, and weed-seeds contained in stable manure (artisan fields) also contribute considerably to such process.

The species which dominate *Echinochloa-Setarietum* (*Amaranthus retroflexus*, *Digitaria sanguinalis*, *Echinochloa crus-galli*, *Setaria pumila*, *S. viridis*, *Sorghum halepense*) are almost all C4, and thus need high temperature for their peculiar metabolism.

The association mainly occurs on basic soils, from gravels to silt and clay but also on the «terre rosse» of the Karst plateau.

The altitude ranges from sea level to 350-400 m. In Friuli the warm form reaches 100-150 meters, being replaced by the cool form at higher altitudes.

Synchorology. This association occurs in Central Europe but it is also present south of the Alps and in the Italian and Slovenian Karst; the southern limit of

its distribution is yet unknown.

Ass.: *Galeopsido tetrahit-Galinsogetum parviflorae* ass. nova hoc loco (Tab. 7).

"The vegetation of weeded summer cultivation of the mountains"

Holotypus: Tab. 7 rel. 10.

Diff. species: *Galium aparine*, *Galeopsis tetrahit*, *G. speciosa*, *Mentha arvensis*, *Stachys palustris*, *Galinsoga ciliata* and *Aethusa cynapium*.

Floristic composition. The coenosis is dominated by *Galinsoga parviflora*, although it is sometimes entirely replaced by the congeneric *G. ciliata*. *Sonchus oleraceus*, *Chenopodium album*, *Polygonum persicaria*, *Stellaria media* and *Fallopia convolvulus* are very frequent and show high cover values. There are also interesting penetrations of *Galeopsis speciosa* and *G. tetrahit* from the edge of the surrounding woods.

Syntaxonomy. The syntaxonomical framing of the relevés has been rather problematic. Although the floristic link with *Echinochloo-Setarietum* is evident, most of the *Panicoideae* species (differentials of that association) are missing. Due to their C4 metabolism, they are not able to grow (or are not competitive) in mountain climate. Friuli relevés have been compared with *Galinsogeto-Portulacetum* Br.-Bl. 1949 (Braun-Blanquet 1949; Pedrotti 1959a, 1959b), described for the Central Alps, where the cultivation of autumn-winter cereals is possible thanks to the continental climate, with spring and summer less rainy and warmer than in Carnia. The floristic compositions of the two coenoses are significantly different, and they cannot be referred to the same association. The floristic differences are due to the coexistence of *Centaureetalia* and *Chenopodetalia* species in the Central Alps association. Therefore we prefer to consider the Carnic relevés as a new, independent association with the name of *Galeopsido tetrahit-Galinsogetum parviflorae*.

This association has to be referred to the *Panico-Setarion* alliance for the presence of *Galinsoga parviflora*, *Setaria pumila* and *Chenopodium album*. Among *Chenopodietalia* species *Sonchus oleraceus*, *Euphorbia helioscopia* and *Lamium purpureum* are the most frequent; among the *Stellarietea* ones *Chenopodium album*, *Polygonum persicaria*, *Stellaria media*, *Capsella bursa-pastoris* and *Fallopia convolvulus*.

Life forms and chorotypes. Therophytes are dominant (71.4%), followed by hemicryptophytes

(20.6%) and geophytes (8.1%).

Because of the particular climatic conditions in which this coenosis occurs, large distribution chorotypes dominate; among these, Cosmopolitan (32.6%), eurasiatic (20.7%) and circumboreal (12.5%) are the best represented. For the same reason, the frequency values of eurymediterranean (1.9%) and steno-mediterranean species (0.3%) are very low, whereas those of adventitious (14.2%) and palaeotemperate (12.6%) are similar to the other coenoses studied in this paper.

Synecology. The association has been found in small fields near villages in cool mountain valleys, especially in Carnia. These fields are mainly cultivated with potatoes and beans (sometimes in the same plot of land) and, due to the excessive altitude, maize is only seldom present. This is a rare example of artisan agriculture, where the few tilling works consist in a single hoeing operation before sowing and in a light weeding during the growth period. In the potato fields, harvest is at the end of September, while bean fields do not undergo further tilling. The soil is left to rest till the next sowing. These fields hardly exceed 300 m².

Galeopsido-Galinsogetum occurs in peculiar ecological conditions, in areas with high rainfalls (1600-2500 mm per year), from 400 to 1100 m on sea level, on rendzinas and brown soils, with a high input of organic matter (manuring).

Synchorology. So far, this association has been recorded only in weeded fields of the mountains of Friuli.

Ass.: *Papaveretum apuli* ass. nova hoc loco (Tab. 8)
"The vegetation of autumn-winter cereals of the plains of Veneto and Friuli"

Holotypus: Tab. 8, rel. 3.

Charact. and diff. species (D): *Papaver apulum*, *Oxalis fontana* (D).

Floristic composition. The coenosis is dominated by *Centaureetalia* (*Papaver rhoeas*, *Fumaria officinalis*, *Centaurea cyanus*, *Alopecurus myosuroides* and *Anthemis arvensis*) and by *Caucalidion* (*Ranunculus arvensis*, *Legousia speculum-veneris*, *Avena fatua* and *Consolida regalis*) species. *Chenopodium album*, *Polygonum persicaria*, *Calystegia sepium* and *Poa sylvicola* are almost always present.

Syntaxonomy. The framing of this association was problematic. Pignatti (1952) found in the Venetian plain an association which was identified as *Alchemillo arvensis-Matricarietum* R. Tx. 1937. The

same author, following Tuxen's scheme (Tuxen 1950) for Central Europe, placed it in the alliance *Scleranthion annui* (Kreusman et Vlieger 1939) Sissingh in Westhoff *et al.* 1946 (which includes acid soil coenoses), although he noted the absence of alliance species, replaced by differential species of *Caucalidion* (which includes basiphilous associations). In this respect Pignatti admitted the existence of one or more basiphilous associations of *Caucalidion*, but did not formally distinguish them (Ferro 1990). In this paper a new association is therefore described with the name of *Papaveretum apuli*.

Ranunculus arvensis, *Avena fatua*, *Legousia speculum-veneris* and *Consolida regalis* show the adherence to *Caucalidion*, *Papaver rhoeas*, *Fumaria officinalis*, *Centaurea cyanus*, *Alopecurus myosuroides* and *Veronica arvensis* to *Centaureetalia*, while the presence of *Chenopodium album*, *Polygonum persicaria*, *Lactuca serriola* and *Anagallis arvensis* supports the framing in *Stellarietea*.

Life forms and chorotypes. Therophytes are the most frequent life form (67.3%), but the percentage of hemicryptophytes is also high (25.5%).

Among the studied coenoses, *Papaveretum apuli* shows the highest value of stenomediterranean (6.5%) and eurymediterranean (22.9%) species. The remarkable percentage of neophytes (12.2%) underscores the level of anthropogenic disturbance of this association.

Synecology. This coenosis develops in cultures of autumn-winter cereals and is therefore characterized by the presence of spring therophytes. These species must fructify before the harvest in June.

The soil texture is heterogeneous: it varies from gravel to silt and clay; the soils mainly derive from calcareous substrata. This coenosis occurs below 100-150 m on sea level.

Synchorology. So far, this association has been detected only in Friuli-Venezia Giulia, but it is probably present in the whole of the Venetian plain.

Conclusions

The continuous adjustment of the segetal flora to the changing cultural techniques causes high instability in the floristic composition of the coenoses. Therefore, in the characterization of these plant communities, we privileged some ecological-structural characters (life form / growth form, and species with C4 metabolism) which remain constant.

The crop field and vineyard vegetation of Friuli-

Venezia Giulia can be ascribed to four associations. This situation, due to the great changes in cultivation techniques, does not coincide with that described about 40 years ago by Pignatti (1952) and Lorenzoni (1963). Among the associations described at that time, *Panico-Polygonetum* is still valid as a well defined geographical race of *Echinochloo-Setarietum*. In Friuli-Venezia Giulia the variability of this coenosis is very high, and it is possible to identify two geographical sub-races (Karst and Friuli), two ecological-altitudinal forms, one subassociation on drained soils (*xanthetosum italicum*) and one agroform in fields with a strong use of herbicides (with *Sorghum halepense*). In the same way *Cerastio-Geranietum dissecti*, described by Poldini (1980), is considered here as a geographical race of *Geranio-Allietum*, containing three different agroforms (with *Taraxacum officinale*, with *Agrostis stolonifera* and with *Arrhenatherum elatius*) induced by different cultivation techniques. Other two coenoses of autumn-winter cereals fields and of small weeded fields of the mountain belt are here described for the first time. They are, respectively, *Papaveretum apuli* and *Galeopsido tetrahit-Galinsogietum parviflorae*.

Appendix

Tab. 4 - *Geranio rotundifolii-Allietum vineale*

Localities: 1: Spessa di Cividale 9946/1 (UD); 2: Villanova di Farra 0047/3 (GO); 3: Sgonico 0248/3 (TS); 4: Samatorza 0248/1 (TS); 5: Pliskovica 0349/1 (SLO); 6: Bonetti 0147/3 (GO); 7: Sales 0248/3 (TS); 8: Dutovlje 0248/2 (SLO); 9: Godnje 0349/1 (SLO); 10: ibidem; 11: ibidem; 12: ibidem; 13: Tomaj 0348/3 (SLO); 14: Kreplje 0348/3 (SLO); 15: near Rocca Bernarda 9946/1 (UD); 16: Moimacco (near Villa Puppi) 9946/1 (UD); 17: Felettis (Palmanova) 0045/1 (UD); 18: Virco 0144/1 (UD); 19: Moraro 0047/3 (GO); 20: between Mossa and Capriva 0047/3 (GO); 21: Corona (Mariano del Friuli) 0046/2 (GO); 22: Villanova di Farra 0047/3 (GO); 23: Moraro 0047/3 (GO); 24: Capriva del Friuli 0047/1 (GO); 25: Villanova di Farra 0047/3 (GO); 26: ibidem; 27: Moraro 0047/3 (GO); 28: Capriva del Friuli 0047/1 (GO); 29: Savogna d'Isonzo 0147/1 (GO); 30: between Lucinico and Giasbana 0047/2 (GO); 31: between Villanova and S. Lorenzo 0047/2 (GO); 32: Lucinico 0047/3 (GO); 33: Corno di Rosazzo 0046/2 (UD); 34: Gagliano di Cividale 9946/2 (UD); 35: Cividale 9946/2 (UD); 37: Campeggio (Faedis) 9846/2 (UD); 38: Orzano (Remanzacco) 9946/3 (UD); 39: Varmo 0143/2 (UD); 40: Vendoglio 9845/1 (UD); 41: Buttrio 9744/3 (UD); 42: Sterpo 0044/3 (UD); 43: between Ariis and Torsa 0144/2 (UD); 44: Faedis 9846/1 (UD); 45: Tauriano 9843/3 (PN); 46: ibidem; 47: ibidem; 48: ibidem; 49: ibidem; 50: ibidem; 51: ibidem; 52: Capriva del Friuli 0047/1 (GO); 53: Torsa 0144/2 (UD); 54: Iutizzo (Codroipo) 0043/4 (UD); 55: S. Pietro di Varmo 0043/4 (UD); 56: Sevegliano 0045/3 (UD); 57: Muzzana del Turgnano

Tab.8 - *Papaveretum apuli* ass. nova hoc loco

Relevé number		1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17			
Altitude (x 10 m)		6	6	4	23	9	11	2	25	10	10	8	11	2	1	1	9	24			
Number of species (spor. incl.)		31	30	37	37	36	41	35	32	31	39	40	29	19	30	36	37	46			
Char. & diff. (DAss) sp. ass.																			Pr.	Fr.	Cl.
K, DAss	Papaver apulum	+	+	1	+	+			1	+	2	1	+			+	2	2	13	76	IV
	Oxalis fontana		+	+	1	+							+			+	+		7	41	III
Char. & diff. (DAll) sp. all.																					
(Caucadon)																					
DAll	Ranunculus arvensis	+	+	1	+	1	2	+	2	+	+	1	+		+		+	2	15	88	V
	Legousia speculum-veneris			1	+	+	+	+	2	+	3	1	+		2	+	1	1	14	82	V
DAll	Consolida regalis	1	+	+	+	+	+	+	+		3	1					+	+	11	65	IV
	Avena fatua	+	+	1			1	1	+	2	+				+	+			10	59	III
DAll	Silene vulgaris		+		+	+	+				+	1	1		+				8	47	III
	Valerianella rimosa									+	+	+	+				+	+	6	35	II
DAll	Chaenorrhinum minus				+			+			+						+	+	5	29	II
DAll	Valerianella locusta			2					+			+					+		4	24	II
	Valerianella eriocarpa	+														+			2	12	I
	Fumaria vaillantii			+									+						2	12	I
	Ajuga chamaepitys											1	+						2	12	I
Char. & diff. (DO) sp. ord.																					
(Centaureetana)																					
DO	Papaver rhoeas	1	2	2	+	1	+	+	1	1	4	3	1	2	2	2	3	2	17	100	V
DO	Fumaria officinalis	+	+	1	1	1	+	1	1	+	1	+				+	+	1	14	82	V
	Centaurea cyanus	2	1	4	1	+	+	2	4	1	2		3				1		12	71	IV
DO	Alopecurus myosuroides	+	+	1				3	2	1	2	2	1		+	3			11	65	IV
DO	Anthemis arvensis	+	2	1		+		+	1	1	1	2	1						10	59	III
DO	Veronica arvensis				+	+	+				+	+	+		+		+	+	9	53	III
	Cerastium glomeratum				+	+	+				1	1	1	+			1	+	9	53	III
DO	Viola tricolor	3		+	+	+			1	+	1	1							8	47	III
DO	Myosotis arvensis	+		+			+	+	+	+	+						+		8	47	III
	Raphanus raphanistrum				1	1		+	+	2	+						+	1	8	47	III
	Buglossoides arvensis	+	1			1	2			2		+	+						7	41	III
	Arabis thaliana					+					2	+					+	1	6	35	II
DO	Matricaria chamomilla			+			+									4	+	2	5	29	II
	Papaver argemone				+		+	+							+				4	24	II
	Lolium multiflorum						1	1	2										3	18	I
	Legousia hybrida	+	+																2	12	I
	Rapistrum rugosum													+	+				2	12	I
	Aphanes arvensis			1															1	6	I
Class species (Stellarietea)																					
	Chenopodium album	+	4	+	+	1	+	+	+	+	+	+		2	+	+	2	2	16	94	V
	Polygonum persicaria		+		1	+	+	+	1	+	+	+		2	1	3	+	+	14	82	V
	Capsella bursa-pastoris	+	+	1	1	+	+	+	1				+	+		+	+	+	13	76	IV
	Anagallis arvensis	+	+	+	1	+	+	+			+			+	+	+	1	+	13	76	IV
	Veronica persica	+	+	1	1	+					+	1		+	+	+	1	+	13	76	IV
	Stellaria media	+	+	+	1	+	+	+	+		1			+	+	1	+		13	76	IV
	Vicia sativa		+	+	+	1		1	1	+	+	+			+	+			12	71	IV
	Fallopia convolvulus	2		+		1	1	+	+	+	1	1		1			1		11	65	IV
	Erigeron annuus					2	+		+		+	1	2				1	1	9	53	III
	Sinapis arvensis			+	+	1	+		1	2					+				7	41	III
	Lactuca serriola			+							1	+				2	2	3	6	35	II
	Senecio vulgaris	+										+				1	+	1	5	29	II
	Cirsium arvense		+	+						1						1			5	29	II
	Bromus sterilis					+	1	+									+		4	24	II
	Mercurialis annua	+	+														+		3	18	I
	Sonchus oleraceus	+													+	1			3	18	I
	Amaranthus retroflexus			+													+	1	3	18	I
	Euphorbia helioscopia				+							+					+		3	18	I
	Sonchus asper				+												+	+	3	18	I
	Veronica hederifolia					+		+		+									3	18	I
	Conyza canadensis					+			+						+				3	18	I

Tab.8 - Continued.

<i>Brassica napus</i>	+	1																2	12	I
<i>Xanthium italicum</i>					+		1											2	12	I
<i>Datura stramonium</i>						+								+				2	12	I
<i>Lamium purpureum</i>					+									+				2	12	I
<i>Geranium dissectum</i>			+															1	6	I
<i>Geranium molle</i>			+															1	6	I
<i>Mentha arvensis</i>				1														1	6	I
<i>Muscari comosum</i>					+													1	6	I
<i>Euphorbia falcata</i>										1								1	6	I
<i>Setaria pumila</i>											+							1	6	I
<i>Oxalis corniculata</i>													+					1	6	I
<i>Amaranthus graecizans</i>																1		1	6	I
<i>Euphorbia peplus</i>															+			1	6	I
<i>Cardamine hirsuta</i>															+			1	6	I
<i>Digitaria sanguinalis</i>																1		1	6	I
<i>Galinsoga parviflora</i>																+		1	6	I
Companions																				
<i>Calystegia sepium</i>	2	1		1	1	+	1	1	1	+	1	1	2	+	1	1	1	16	94	V
<i>Poa sylvicola</i>			+	1	2	2	2		1	2	2	1	1	2	1	1	1	14	82	V
<i>Polygonum aviculare</i>	1		+	1	1	+	+	1	1	1			1	1	+	+		13	76	IV
<i>Mentha longifolia</i>	+	+	+	1	1	1		+	+	+			+	+	+	+		12	71	IV
<i>Sorghum halepense</i>	+	1	+	+						+	1	+			1	1	2	10	59	III
<i>Silene alba</i>	+	+				+	+	+	+	1	1	1			1			10	59	III
<i>Arenaria serpyllifolia</i>		1				+	+			1	+	1		+	1	+	+	10	59	III
<i>Lolium multiflorum</i>	1	+		1					2	1		1			+	1	1	9	53	III
<i>Medicago lupulina</i>			+								+	+			+	+		6	35	II
<i>Convolvulus arvensis</i>				+		1				+				1	1		1	6	35	II
<i>Artemisia verlotorum</i>	1					+	+	1							+			5	29	II
<i>Daucus carota</i>				+	+	+				2				1				5	29	II
<i>Agropyron repens</i>				1	1			+					2	+				5	29	II
<i>Ranunculus arvensis</i>						+	+				+					1		5	29	II
<i>Rorippa sylvestris</i>						+		+				+			1		+	5	29	II
<i>Galium aparine</i>						+		+						+	+	+		5	29	II
<i>Trifolium repens</i>		+		1									+	1				4	24	II
<i>Artemisia vulgaris</i>				+						2	1					2		4	24	II
<i>Conyza albida</i>						+				+						1	1	4	24	II
<i>Crepis taraxacifolia</i>						+				+	2					2		4	24	II
<i>Equisetum arvense</i>													1	2	+	+		4	24	II

0144/4 (UD); 58: Colloredo di Montalbano 9844/2 (UD); 59: Buttrio 9744/3 (UD); 60: Nimis 9745/4 (UD); 61: S. Croce 0248/3 (TS); 62: ibidem.

Sporadic species: *Daucus carota*: + (1), + (15), + (17), + (19), r (35), + (37), + (48), + (56), + (59); *Urtica dioica*: + (28), + (37), + (43), 1 (45), 3 (50), 1 (51), 4 (54), + (56), r (60); *Trifolium incarnatum/molineri*: 1 (4), 1 (7), + (8), + (10), + (11), + (12), + (13); *Arenaria serpyllifolia*: + (13), + (17), + (20), r (24), r (26), + (31), r (35); *Cynodon dactylon*: + (13), 1 (17), r (26), + (34), 1 (43); *Raphanus landra*: + (16), + (17), + (19), + (22), 1 (34), 2 (35); *Lamium maculatum*: + (10), + (21), + (22), + (36), + (62); *Clematis vitalba*: r (15), + (16), + (49), + (50), + (62); *Artemisia vulgaris*: + (18), + (21), 1 (26), 1 (32), + (49); *Ajuga reptans*: + (38), + (55), + (58), + (59), + (60); *Crepis foetida*: + (3), + (4), + (20), + (21); *Galium aparine*: + (5), + (12), r (23), r (58); *Ranunculus acris*: + (15), r (44), + (55), + (60); *Polygonum aviculare*: + (1), + (48), + (50); *Brassica rapa*: 1 (4), + (10), + (21); *Cerastium pumilum*: + (6), 2 (14), r (61); *Galium lucidum*: + (15), 1 (37), + (38); *Medicago sativa*: r (35), + (52), + (56); *Pastinaca sativa*: + (39), 2 (51), + (60); *Glechoma hederacea*: + (44), + (54), 1 (57); *Anthoxanthum odoratum*: + (58), + (59), + (60); *Picris hieracioides*: + (39), 1 (47); *Cardaria draba*: 1 (4), +

(13); *Malva sylvestris*: + (5), + (44); *Poa pratensis*: + (86), + (62); *Sisymbrium officinale*: + (10), + (34); *Rorippa sylvestris*: + (13), 1 (16); *Allium ampeloprasum*: 1 (14), r (44); *Aristolochia clematidis*: + (16), + (62); *Sambucus ebulus*: + (16), + (54); *Leucanthemum vulgare*: + (17), + (56); *Artemisia verlotorum*: 1 (19), + (61); *Ranunculus sardous*: 1 (21), + (31); *Capsella rubella*: + (29), 1 (36); *Silene vulgaris*: + (34), + (60); *Holcus lanatus*: + (37), + (58); *Verbena officinalis*: + (45), + (46); *Crepis capillaris*: + (48), + (51); *Veronica chamaedrys*: 1 (54), 1 (60); *Brachypodium sylvaticum*: + (61), + (62); *Carex pairae*: 1 (61), + (62); *Rubus caesius*: r (61), + (62); *Linaria vulgaris*: + (3); *Bromus madritensis*: + (4); *Erophila verna*: + (5); *Vicia hirsuta*: + (8); *Plantago media*: + (13); *Medicago minima*: + (13); *Trifolium campestre*: + (13); *Inula salicina*: + (15); *Tussilago farfara*: + (15); *Ononis spinosa*: + (16); *Solanum dulcamara*: + (16); *Leontodon hispidus*: + (17); *Sanguisorba minor*: r (18); *Arabidopsis thaliana*: + (19); *Bromus molliformis*: + (19); *Ranunculus parviflorus*: + (23); *Cirsium vulgare*: + (28); *Fragaria vesca*: + (28); *Lapsana communis*: + (34); *Senecio inaequidens*: + (34); *Tragopogon orientalis*: + (34); *Lotus corniculatus*: + (37); *Saponaria officinalis*: + (37); *Hypochaeris radicata*: + (38); *Cerastium brachypetalum*: + (40); *Carex hirta*: + (42); *Euphorbia*

lathyris: r (42); *Aegopodium podagraria*: + (44); *Buphthalmum salicifolium*: + (44); *Hypochoeris glabra*: + (52); *Valeriana officinalis*: + (52); *Pimpinella major*: + (57); *Cruciata laevipes*: + (59); *Allium sphaerocephalon*: 1 (61); *Avena barbata*: 1 (61); *Medicago lupulina*: r (62); *Eupatorium cannabinum*: 1 (62).

Tab. 6 - Echinochloo-Setarietum pumilae

Localities: 1: Enemonzo 9543/3 (UD); 2: Villa Santina 9543/4 (UD); 3: Esemont di Sotto 9543/3 (UD); 4: Formeaso 9544/3 (UD); 5: Socchieve 9643/1 (UD); 6: Enemonzo 9543/3 (UD); 7: Esemont di Sopra 9543/3 (UD); 8: Esemont di Sopra 9543/3 (UD); 9: Terzo 9544/3 (UD); 10: Subida (Cormons) 0046/2 (GO); 11: Moraro 0047/3 (GO); 12: Capriva del Friuli 0047/3 (GO); 13: Terzo 9544/3 (UD); 14: Esemont di Sotto 9543/3 (UD); 15: Cavazzo 9644/1 (UD); 16: Amaro 9644/2 (UD); 17: Manzano 0046/1 (UD); 18: Gorgo (Latisana) 0244/1 (UD); 19: Gorgo (Latisana) 0244/1 (UD); 20: Gorgo (Latisana) 0244/1 (UD); 21: Latisana 0144/3 (UD); 22: Perteole 9945/1 (UD); 23: Flambruzzo 0144/1 (UD); 24: Marano 0245/1 (UD); 25: Aquileia 0246/1 (UD); 26: S. Giorgio di Nogaro 0145/3 (UD); 27: Preval (Cormons) 0047/1 (GO); 28: ibidem; 29: Fagagna 9844/4 (UD); 30: Tolmezzo 9544/3 (UD); 31: Terzo 9544/3 (UD); 32: Gemona 9744/2 (UD); 33: ibidem; 34: Artega 9744/4 (UD); 35: Flambruzzo (Rivignano) 0144/1 (UD); 36: Romans d'Isonzo 0146/2 (UD); 37: Terzo 9544/3 (UD); 38: Fagagna 9844/4 (UD); 39: Maiano 9844/1 (UD); 40: Terzo 9544/3 (UD); 41: Cavazzo 9644/1 (UD); 42: ibidem; 43: Gemona 9744/2 (UD); 44: Cavazzo 9644/1 (UD); 45: Codroipo 0043/1 (UD); 46: Pliskovica 0349/1 (SLO); 47: Ferletti 0147/3 (GO); 48: Medeazza 0247/2 (TS); 49: Sales 0248/3 (TS); 50: S. Michele 0147/1 (GO); 51: Pliskovica 0349/1 (SLO); 52: Medeazza 0247/2 (TS); 53: Dutovlje 0349/1 (SLO); 54: Lokev 0349/1 (SLO); 55: Villanova di Farra 0047/3 (GO); 56: Grozzana 0349/3 (TS); 57: ibidem; 58: Basovizza 0349/1 (TS); 59: Percedol 0248/3 (TS); 60: S. Canziano 0349/2 (SLO); 61: Rupinpiccolo 0248/4 (TS); 62: ibidem; 63: Sgonico 0248/3 (TS).

Sporadic species: *Medicago lupulina*: r (5), + (9), + (14), + (29), r (37), + (46), r (52), + (54), + (55); *Trifolium repens*: + (2), + (5), + (13), + (14), + (15), r (23), + (40), + (66); *Artemisia vulgaris*: r (15), r (18), + (21), + (22), r (32), + (33), + (37), r (55); *Diplotaxis tenuifolia*: r (6), 1 (20), + (29), + (48), + (50), + (25); *Rumex crispus*: + (12), r (15), 1 (16), + (32), + (50), r (52), + (54); *Plantago major*: r (3), + (6), + (25), + (27), + (32), + (57); *Daucus carota*: r (3), + (14), + (27), + (29), r (38), + (46); *Mentha spicata*: + (22), + (25), + (34), 1 (38), + (47), r (55); *Plantago lanceolata*: r (12), + (47), + (51), + (57), + (66); *Verbena officinalis*: + (12), r (21), r (24), r (25), + (55); *Potentilla reptans*: r (21), + (29), + (36), 1 (47), 2 (52); *Vicia sepium*: + (7), + (9), r (30), r (43); *Rumex obtusifolius*: r (29), + (34), 2 (54), 1 (58); *Silene vulgaris*: r (1), r (14), r (55); *Agrostis stolonifera*: + (7), r (15), 1 (51); *Rubus caesius*: r (22), 1 (38), 1 (39); *Lythrum salicaria*: r (21), + (28), r (36); *Aristolochia clematidis*: + (51), + (53), 1 (66); *Poa annua*: + (3), + (12); *Glechoma hederacea*: r (5), 1 (12); *Rorippa sylvestris*: + (11), 1 (52); *Arenaria serpyllifolia*: + (14), r (33); *Cirsium vulgare*: r (16), r (39); *Heliotropium europaeum*: + (29), r (52); *Armoracia rusticana*: 1 (5); *Chaenarrhinum minus*: r (14); *Mentha longifolia*: + (21); *Cichorium intybus*: r (23); *Abutilon theophrasti*: + (26); *Polygonum mite*: + (27); *Picris hieracioides*: + (29); *Helianthus tuberosus*: + (30); *Aegopodium podagraria*: + (31); *Hypericum perforatum*: r (33); *Lysimachia nummularia*: r (36); *Sambucus ebulus*: + (37); *Vicia cracca*: r (37); *Brassica nigra*: r (38); *Picris echioides*: + (38); *Crepis rheoadifolia*: + (46); *Nicandra*

physalodes: + (47); *Chondrilla juncea*: r (50); *Achillea millefolium*: + (51); *Centaurea weldeniana*: r (51); *Galium mollugo*: + (51); *Leucanthemum vulgare*: r (51); *Prunella vulgaris*: + (51); *Misopates orontium*: r (52); *Stachys annua*: + (53); *Lolium multiflorum*: + (57); *Medicago sativa*: 1 (57); *Euphorbia falcata*: + (63).

Tab. 7 - Galeopsido tetrahit-Galinsogetum parviflorae

Localities: 1: Socchieve 9543/3 (UD); 2: S. Pietro di Zuglio 9544/1 (UD); 3: Cercivento 9443/3 (UD); 4: Ampezzo 9542/4 (UD); 5: Avaglio (Lauro) 9543/3 (UD); 6: Forni Avoltri 9442/2 (UD); 7: Ravascletto 9443/3 (UD); 8: between Liariis and Clavais (Ovaro) 9543/1 (UD); 9: Magnanins (Ravascletto) 9443/3 (UD); 10: Lenzone (Ovaro) 9543/1 (UD); 11: Cella (Ovaro) 9543/1 (UD); 12: Lenzone (Ovaro) 9543/1 (UD); 13: Liariis (Ovaro) 9543/1 (UD); 14: Oltris (Ampezzo) 9542/4 (UD); 15: between Trava and Avaglio (Lauro) 9543/3 (UD); 16: Osais (Val Pesarina) 9442/2 (UD); 17: Pesariis 9442/4 (UD); 18: Cercivento 9443/3 (UD); 19: Sauris di sotto 9542/1 (UD); 20: Villa Santina 9543/2 (UD); 21: Paularo 9444/3 (UD); 22: Mersino sup. (M. Matajur) 9847/3 (UD); 23: Malga Vidoni (M. Chiampon) 9745/1 (UD).

Sporadic species: *Poa annua*: + (7), + (10), + (17), 1 (20); *Trifolium pratense*: + (13), + (21), + (22), + (23); *Trifolium repens*: + (4), + (13), + (22), + (23); *Daucus carota*: + (10), + (12), + (23); *Aegopodium podagraria*: + (16), + (19), + (23); *Silene alba*: r (2); + (19); *Polygonum aviculare*: + (4), + (8); *Urtica dioica*: + (4), 1 (23); *Nepeta cataria*: + (5), + (15); *Lamium album*: + (6), + (16); *Glechoma hederacea*: + (11), + (20); *Myosoton aquaticum*: 1 (11), 1 (12); *Leucanthemum vulgare*: + (19), + (22); *Phalaris canariensis*: + (1); *Artemisia vulgaris*: r (2); *Lamium maculatum*: + (2); *Rumex obtusifolius*: + (6); *Myosotis sylvatica*: r (8); *Atriplex tatarica*: r (13); *Galium lucidum*: + (14); *Geranium pyrenaicum*: + (15); *Agropyron repens*: 1 (19); *Lathyrus pratensis*: + (19); *Ranunculus acris*: + (19); *Silene vulgaris*: + (19); *Vicia hirsuta*: r (19); *Vicia sepium*: + (19); *Dactylis glomerata*: + (20); *Arrhenatherum elatius*: + (21); *Borago officinalis*: + (21); *Euphorbia lathyris*: + (21); *Lotus corniculatus*: + (22); *Centaurea nigrescens*: + (23); *Crepis capillaris*: + (23); *Galium mollugo*: + (23); *Holcus lanatus*: + (23); *Lolium multiflorum*: 1 (23); *Malva neglecta*: + (23); *Prunella vulgaris*: + (23); *Veronica chamaedrys*: + (23).

Tab. 8 - Papaveretum apuli

Localities: 1: Lucinico 0047/3 (GO); 2: Farra d'Isonzo 0047/3 (GO); 3: Gradisca 0147/1 (GO); 4: S. Vito di Fagagna 9944/4 (UD); 5: Mereto di Tomba 9944/3 (UD); 6: Orzano (Remanzacco) 9946/2 (UD); 7: Palmanova 0045/4 (UD); 8: Fagagna 9844/4 (UD); 9: Cernegons 9954/2 (UD); 10: Udine 9945/2; 11: Udine Sud 9945/2 (UD); 12: Premariacco 9946/2 (UD); 13: Cervignano 0146/4 (UD); 14: Carlino (Marano) 0145/3-0245/1 (UD); 15: Torviscosa 0145/4 (UD); 16: Pradamano 9945/2 (UD); 17: Cicconico di Fagagna 9844/3 (UD);

Sporadic species: *Potentilla reptans*: + (1), 1 (13), 1 (14); *Rumex crispus*: + (1), + (6), 1 (15); *Reseda lutea*: + (4), + (10), + (12); *Rumex obtusifolius*: 1 (8), + (9), 1 (11); *Rubus caesius*: + (13); + (14), + (15); *Medicago sativa*: 2 (4), 3 (6); *Ajuga genevensis*: + (5), 1 (11); *Plantago major*: + (6), 1 (9); *Helianthus tuberosus*: 1 (7), 1 (14); *Cerastium holosteoides*: + (10), + (17); *Taraxacum officinale*: 1 (10), + (17); *Saxifraga tridactylites*: + (11), + (16); *Acinos arvensis*: + (11), + (12); *Fagopyrum tataricum*: 1 (1); *Achillea millefolium*: 1 (2); *Melilotus officinalis*: + (2); *Trifolium pratense*: + (2); *Vicia cracca*: + (2); *Aphanes arvensis*: 1 (3);

Bifora radians: + (3); *Lithospermum officinale*: 4 (3); *Diplotaxis tenuifolia*: + (4); *Fallopia dumetorum*: 1 (4); *Galium mollugo*: + (4); *Arrhenatherum elatius*: + (5); *Glechoma hederacea*: + (5); *Myosoton aquaticum*: + (5); *Rumex acetosa*: + (5); *Bidens frondosa*: 1 (7); *Geum urbanum*: + (7); *Brassica oleracea*: + (8); *Humulus lupulus*: + (9); *Oxalis acetosella*: + (9); *Hypochoeris radicata*: + (10); *Senecio inaequidens*: 1 (10); *Galium lucidum*: 1 (11); *Plantago lanceolata*: 1 (11); *Saponaria officinalis*: 1 (11); *Verbena officinalis*: + (11); *Aegopodium podagraria*: 2 (13); *Equisetum ramosissimum*: + (14); *Festuca arundinacea*: + (14); *Plantago media*: 1 (14); *Lysimachia vulgaris*: + (15); *Dactylis glomerata*: + (16); *Erophila verna*: + (16); *Leucanthemum vulgare*: + (16); *Trifolium campestre*: + (16); *Crepis foetida*: 1 (17); *Rumex conglomeratus*: + (17).

Acknowledgments

The work was carried out with the financial support of M.U.R.S.T. «Analisi causale della vegetazione dell'Italia nord-orientale» (responsibile Prof. Dr L. Poldini).

References

- Baker H.G., 1972. *The evolution of weed*. Ann. Rev. Ecol. Syst., 5:1-24.
- Braun-Blanquet J., 1936. *Classe des Rudereto-Secalinetales. Groupements messicoles, culturaux et nitrophiles-ruderaux du cercle de végétation méditerranéen*. Prodrome des groupements végétaux, fasc. 3, Montpellier.
- Braun-Blanquet J., 1949. *Übersicht der Pflanzengesellschaften Rätien*. (II). Vegetatio, 1:129-146.
- Cassani S., 1986. *Un alleato prezioso*. Lottantiparassitaria, XXXVIII, 3: 3.
- Chiappella Feoli L. & Poldini L., 1993. *Prati e pascoli del Friuli (NE Italia) su substrati basici*. Studia Geobot., 13: 3-140.
- Collins R.P. & Jones M.B., 1985. *The influence of climatic factors on the distribution of C4 species in Europe*. Vegetatio, 64: 121-129.
- Ehrendorfer F., 1973. *Liste der Gefüßpflanzen Mitteleuropas*. 2nd ed., Fischer Verlag, Jena, Stuttgart, New York, 318 pp.
- Faegri K. & Pijl van der L., 1971. *The principles of pollination ecology*. 2nd ed., Pergamon Press, Oxford, 291 pp.
- Ferrari C., Catizone P., Speranza M. & De Polzer S., 1979. *Studio degli effetti selettivi degli erbicidi ammidici e triazinici sulla vegetazione infestante le colture di mais nella pianura padano-veneta*. Notiz. Fitosoc., 15: 63-74.
- Ferro G., 1990. *Revisione della vegetazione segetale mediterranea ed europea dell'ordine Secalietalia*. Braun-Blanquetia 6:1-59.
- Géhu J., 1996. *Epistémologie de la territorialité en phytosociologie*. Giorn. Bot. Ital., 130: 189-199.
- Gerdol R. & Piccoli F., 1979. *Considerazioni sulla sinsistemica della vegetazione infestante delle colture sarchiate del Ferrarese*. Arch. Bot., 55: 114-123.
- Gutierrez M., Gracen V.E. & Edward G.E., 1974. *Biochemical and cytological relationships in C4 plants*. Planta Berl., 119: 547-579.
- Krippelová T., 1979. *Sur la problématique des communautés des alliances Polygono-Chenopodion Koch 1926 em. Sissingh 1946 et Panico-Setarion Sissingh 1946 dans le bassin de Kosice (Slovaquie Sud-Est)*. Notiz. Fitosoc., 15: 21-25.
- Krippelová T., 1981. *Synanthrope Vegetation des Beckens Košická kotlina*. Vegetácia CSSR B4, pp.214.
- Lagonegro M. & Feoli E., 1985. *Analisi multivariata di dati. Manuale d'uso di programmi BASIC per personal computer*. Libreria Goliardica, Trieste, 182 pp.
- Landolt E., 1977. *Ökologische Zeiger zur Schweizer Flora*. Ber. Geobot. Inst. ETH, 68: 64-207.
- Lorenzoni G. G., 1963. *La vegetazione infestante del mais nel Friuli, nel Veneto e in Lombardia*. Maydica, VIII, quaderno 2, 42 pp.
- Maarel van der E., 1979. *Transformation of cover-abundance values in phytosociology and its effects on community similarity*. Vegetatio, 39: 97-114.
- Martini F. & Poldini L., 1995. *The hemerophytic flora of Friuli-Venezia Giulia (NE Italy)*. Flora Medit., 5: 229-246.
- Mucina L., 1993. *Stellarietea mediae*. In: Mucina L., Grabherr G. & Ellmauer T (eds.), *Die Pflanzengesellschaften Österreichs. Teil I: Anthropogene Vegetation*. Fischer Verlag, Jena, Stuttgart, New York, 110-168.
- Müller-Schneider P., 1986. *Verbreitungsbiologie der Blütenpflanzen Graubündens*. Veröff. Geobot. Inst. ETH, Stiftung Rübel, 85: 5-263.
- Oberdorfer E., 1993. *Süddeutsche Pflanzengesellschaften. Teil III: Wirtschaftswiesen und Unkrautgesellschaften*. Fischer Verlag, Jena, Stuttgart, New York, 48-114.
- Pedrotti F., 1959a. *La vegetazione delle colture sarchiate di patata in Val di Sole*. Studi Trentini di Scienze Naturali, 36: 73-91.
- Pedrotti F., 1959b. *Sulla vegetazione infestante le colture sarchiate in Val di Sole*. Giorn. Bot. Ital., 66: 344-345.
- Pignatti S., 1952. *Introduzione allo studio fitosociologico della pianura veneta orientale con particolare riguardo alla vegetazione litoranea*. Arch. Bot., 28: 265-329.
- Pirola A. & Rossetti A., 1974. *Polygono-Xanthietum italicum ass. nova, vegetazione di greto del corso medio del Reno (Bologna)*. Notiz. Fitosoc., 8: 15-27.
- Podani J., 1993. *Syn-Tax-PC: computer programs for multivariate data analysis in ecology and systematics. Version 5.0, user's guide*. Scientia Publishing, Budapest, 104 pp.
- Poldini L., 1980. *Übersicht über die Vegetation des Karstes von Triest und Görz (NO-Italien)*. Studia Geobot., 1:79-130.
- Poldini L., 1989. *La vegetazione del Carso isontino e triestino*. Lint, Trieste, 315 pp.
- Poldini L., 1991. *Atlante corologico delle piante vascolari nel Friuli-Venezia Giulia. Inventario floristico regionale*. Regione Autonoma Friuli-Venezia Giulia, Direzione Regionale delle Foreste e dei Parchi & Università degli Studi di Trieste, Dipartimento di Biologia, Udine, 899 pp.
- Poldini L. & Oriolo G., 1994. *La vegetazione dei prati da sfalcio e dei pascoli intensivi (Arrhenatheretalia e Poo-Trisetetalia) in Friuli (NE Italia)*. Studia Geobot., 14, suppl. 1: 3-48.
- Poldini L., Mazzolini G. & Oriolo G., 1996. *La vegetazione spontanea dei vigneti nei territori nord-est adriatici: Geranio rotundifolii-Allietum R. Tx. ex von Rochow 1951 o Cerastio-Geranium dissecti Poldini 1980?* Giorn. Bot. Ital., 130: 151-152.
- Pollastro L., 1989. *Il diserbo del mais*. Lottantiparassitaria, XLI: 10-11.
- Regula-Bevilacqua L., 1979. *Ruderalna i korovna vegetacija na području Strahinsce*. Acta Bot. Croat., 38: 105-122.
- Ries C., 1992. *Überblick über die Ackerunkrautvegetation Österreichs und ihre Entwicklung in neuerer Zeit*. Dissert. Botanicae, vol. 187.




Fig. 1 - Distribuzione geografica di *Quercus trojana* Webb.
Geographic distribution of *Quercus trojana* Webb.

Area di studio

Sotto il profilo bioclimatico l'area pugliese interessata dai fragneti (Fig. 2) rientra, secondo la classificazione proposta da Rivas-Martinez *et al.* (1991), nel tipo mesomediterraneo subumido inferiore. Infatti sulla base dei diagrammi ombrotermici (Fig. 3), relativi alle stazioni termopluviometriche ricadenti all'interno dell'areale del fragno, si rileva che le temperature medie annue sono comprese fra 14°C e 16°C, mentre le precipitazioni medie annue vanno da circa 600 mm a 700 mm. *Quercus trojana* nel territorio pugliese si insedia normalmente su terre rosse mediterranee originatesi da calcari miocenici, dove forma boschi puri o talora misti a *Quercus ilex* L. e *Q. virgiliana* (Ten.) Ten., o più raramente a *Q. cerris* L. e *Q. calliprinos* Webb.




Fig. 2 - Aree di *Quercus trojana* Webb in Puglia (da Maselli 1940, modif.).

Distribution of *Quercus trojana* in Puglia (from Maselli 1940, modified).

Risultati e Discussione

Sulla base della loro composizione floristica, esigenze ecologiche e correlazioni dinamiche, i boschi a *Quercus trojana* presenti nelle Murge pugliesi mostrano un chiaro carattere termofilo. Infatti in queste formazioni sono presenti numerose sclerofille arboree, arbustive e lianose come *Quercus ilex*, *Phillyrea latifolia*, *Rubia longifolia*, *Rosa sempervirens*, *Lonicera implexa*, *Pistacia lentiscus*, *Viburnum tinus*, *Ruscus aculeatus*, ecc. che normalmente si accompagnano ad altri elementi termofili decidui come *Pyrus amygdaliformis*, *Calicotome infesta*, *Lonicera etrusca*, *Fraxinus ornus*, *Quercus virgiliana*, e *Pistacia terebinthus*.

Nel loro complesso questi fragneti, rientranti nella fascia climacica del *Quercion ilicis*, si insediano su substrati calcarei a quote comprese tra i 100 e i 500 m, dove costituiscono un tipo di vegetazione climacica molto peculiare ed esclusiva in Italia di quest'area pugliese. Abbastanza ricco e ben rappresentato è il contingente di specie appartenenti alla suddetta alleanza e alla relativa classe *Quercetea ilicis*. Oltre alle specie legnose sopracitate, si rinvenivano pure numerosi altri elementi di questi *syntaxa*, come *Cyclamen repandum*, *Carex distachya*, *Asparagus acutifolius*, *Viola dehnhardtii*, *Luzula forsteri*, *Poa sylvicola*, *Tamus communis*, *Carex hallerana*. Per la caratterizzazione floristica dei suddetti *syntaxa* ci si è attenuti soprattutto al lavoro di revisione di *Quercetea ilicis* effettuato da Rivas Martinez (1975) e Brullo & Marcenò (1985).




Fig. 3 - Diagrammi ombrotermici di alcune stazioni delle Murge Pugliesi. I dati di Castellaneta e Gioia del Colle sono stati tratti da Bianco *et al.* (1988); quelli di Turi e Locorotondo da Bianco (1961a).

Climatic diagrams of selected localities in the Murge (Puglia). Data from Bianco et al. (1988) (Castellaneta and Gioia del Colle), and from Bianco (1961a) (Turi and Locorotondo).

L'appartenenza di questa formazione all'alleanza *Quercion ilicis* era già stata evidenziata da Chiesura Lorenzoni *et al.* (1971) e successivamente confermata da Lorenzoni e Chiesura Lorenzoni (1987), che in particolare attribuiscono le cenosi in oggetto ad un *Quercetum trojanae* n.n., in seno al quale differenziano varie subassociazioni.

In effetti, i fragneti delle Murge pugliesi si differenziano abbastanza bene floristicamente dalle altre formazioni boschive del *Quercion ilicis*, oltre che per la dominanza di *Quercus trojana*, anche per la presenza di alcuni elementi abbastanza rari e significativi nel territorio italiano, esclusivi o quasi di queste formazioni. Si tratta in particolare di *Euphorbia apios* e *Potentilla detommasii*, specie entrambe del Mediterraneo nord-orientale, qui abbastanza diffuse e frequenti, come pure di *Arum apulum*, un raro endemismo pugliese. Queste specie nel loro complesso permettono di differenziare una nuova associazione, proposta come *Euphorbio apii-Quercetum trojanae* (holotypus: ril. 11, Tab. 1).

In situazioni microclimatiche e pedologiche più

mesiche i boschi a *Q. trojana* della Puglia si arricchiscono in specie dei *Quercetalia pubescentis-petraeae* e dei *Quercus-Fagetea*, fra cui in particolare *Buglossoides purpureo-coerulea*, *Silene italica*, *Euonymus europaeus*, *Poa sylvicola*, *Brachypodium sylvaticum*, *Quercus cerris*, *Fraxinus ornus*. Queste specie permettono di differenziare un aspetto più mesofilo proposto come subass. *poetosum sylvicolae* (holotypus ril. 10, Tab. 1), che si distingue abbastanza bene dal tipo anche sotto il profilo ecologico.

Sulla base di osservazioni personali inedite e di dati di letteratura (Agostini 1967, Macchia 1987, Bartolo *et al.* 1985, De Marco & Caneva 1985, Brullo *et al.* 1987), nel territorio pugliese l'associazione in oggetto viene sostituita in situazioni più xeriche dall'*Oleo - Quercetum virgilianae* Brullo 1984, che in genere è legato a un clima di tipo termomediterraneo subumido inferiore, o dal *Thymo-Pinetum halepensis* De Marco & Caneva 1985 quando si passa al termomediterraneo secco superiore. In situazioni microclimatiche più

Tab. 1 - *Euphorbia apii* - *Quercetum trojanae*

Numero del rilievo	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15
Quota (m)	320	320	320	310	310	310	420	450	440	440	450	450	430	380	360
Superficie (m ²)	100	100	100	100	100	100	100	100	100	100	100	100	100	100	100
Copertura arborea (%)	100	100	100	100	100	100	100	100	100	100	100	100	100	100	100
Copertura arbustiva (%)	40	40	30	50	50	40	60	60	70	50	50	50	50	60	60
Copertura erbacea (%)	40	30	30	50	30	50	60	60	50	60	60	70	70	70	60
Specie caratteristiche di associazione															
<i>Quercus trojana</i> Webb	4	5	5	4	5	5	5	5	5	5	5	5	5	3	4
<i>Euphorbia apios</i> L.	1	1	2	1	1	+	2	1	1	2	1	1	1	2	1
<i>Potentilla detommasii</i> Ten.	*	+	*	+	1	*	*	+	2	1	1	1	+	1	2
<i>Arum apulum</i> (Carano) Bedalov	*	*	*	*	*	*	*	*	*	*	*	*	*	+	*
Specie differenziali di subassociazione															
<i>Poa sylvicola</i> Guss.	*	*	*	*	*	*	2	1	1	2	2	2	2	3	2
<i>Euonymus europaeus</i> L.	*	*	*	*	*	*	+	+	1	2	2	1	1	1	2
<i>Buglossoides purpureoacerulea</i> (L.) Johnst.	1	+	*	*	+	1	+	1	2	2	2	2	1	2	1
<i>Brachypodium sylvaticum</i> (Hudson) Beauv.	*	*	*	*	*	*	2	1	2	2	2	2	3	*	*
<i>Geum urbanum</i> L.	*	*	*	*	*	*	+	1	1	1	1	1	+	*	+
<i>Galium odoratum</i> (L.) Scop.	*	*	*	*	*	*	+	1	+	1	+	*	*	*	*
<i>Fraxinus ornus</i> L.	*	*	*	*	*	*	1	1	1	1	+	*	*	*	*
<i>Quercus cerris</i> L.	*	*	*	*	*	*	*	*	*	*	*	*	*	3	2
Specie caratteristiche di <i>Quercion ilicis</i> e di <i>Quercetea ilicis</i>															
<i>Rubia longifolia</i> Poiret	1	2	2	2	1	1	*	2	2	2	2	1	1	1	1
<i>Rosa sempervirens</i> L.	1	1	+	+	*	1	*	1	1	1	2	2	2	2	2
<i>Carex distachya</i> Desf.	1	1	+	1	+	*	*	1	+	1	1	1	1	3	2
<i>Pyrus amygdaliformis</i> Vill.	1	1	+	1	+	*	*	+	2	1	2	1	1	*	1
<i>Asparagus acutifolius</i> L.	1	2	+	2	2	2	*	*	*	*	2	1	1	1	1
<i>Quercus virgiliana</i> (Ten.) Ten.	1	1	+	+	1	1	1	1	1	1	1	1	1	2	2
<i>Phillyrea latifolia</i> L.	1	+	1	1	+	1	2	1	1	+	1	*	*	*	*
<i>Lonicera etrusca</i> Santi	*	*	+	1	+	*	*	1	1	1	1	1	+	1	1
<i>Quercus ilex</i> L.	2	1	1	2	2	1	1	+	1	*	*	*	*	*	*
<i>Cyclamen repandum</i> Sibth. et Sm.	1	+	1	+	+	*	*	*	*	2	1	1	1	+	*
<i>Viola dehnhardtii</i> Ten.	+	+	+	+	+	*	*	*	*	1	1	1	+	+	+
<i>Calicotome infesta</i> (C.Presl) Guss.	1	*	+	2	1	1	*	2	1	1	1	1	2	1	+
<i>Ruscus aculeatus</i> L.	*	*	*	*	*	*	*	*	1	1	1	2	1	1	*
<i>Tamus communis</i> L.	*	*	*	*	*	*	2	1	2	1	1	1	*	*	*
<i>Lonicera implexa</i> Aiton	*	*	*	*	*	*	1	2	1	*	*	*	1	1	*
<i>Luzula forsteri</i> (Sm.) DC.	+	*	+	1	+	*	*	*	*	*	*	*	*	+	*
<i>Pistacia lentiscus</i> L.	*	*	*	*	*	*	3	2	2	*	*	*	*	*	*
<i>Pistacia terebinthus</i> L.	*	*	*	*	*	*	*	*	*	1	2	1	*	*	*
<i>Viburnum tinus</i> L.	*	*	*	*	*	*	1	1	1	*	*	*	*	*	*
<i>Carex hallerana</i> Asso	*	+	*	+	+	*	*	*	*	*	*	*	*	*	*
<i>Pulicaria odora</i> (L.) Reichenb.	+	*	+	*	+	*	*	*	*	*	*	*	*	*	*
Altre specie															
<i>Ranunculus neapolitanus</i> Ten.	2	1	1	1	*	1	1	1	1	2	1	1	2	1	+
<i>Crataegus monogyna</i> Jacq.	1	+	1	1	*	+	1	1	2	1	1	1	1	1	2
<i>Dactylis glomerata</i> L.	1	*	+	1	*	2	2	1	1	2	2	2	2	2	2
<i>Geranium sanguineum</i> L.	1	*	+	1	1	2	1	+	+	*	1	1	+	1	+
<i>Silene italica</i> (L.) Pers.	+	1	+	1	1	+	1	*	*	*	1	2	1	2	2
<i>Allium subhirsutum</i> L.	+	+	+	1	1	+	*	*	1	+	1	+	+	*	*
<i>Hedera helix</i> L.	1	+	*	*	1	*	2	1	1	2	2	2	2	2	1
<i>Stachys</i> cfr. <i>salviifolia</i> Ten.	1	+	*	+	1	+	1	*	*	*	+	+	+	+	+
<i>Carex serrulata</i> Biv.	+	*	+	1	+	+	*	*	+	*	+	+	+	1	+
<i>Crepis leontodontoides</i> All.	1	+	*	+	+	*	2	*	*	*	+	1	*	1	1
<i>Rubus ulmifolius</i> Schott	1	+	+	1	2	1	*	3	*	*	*	*	*	1	2
<i>Euonymus europaeus</i> L.	*	*	*	*	*	*	*	+	1	2	2	1	1	1	2
<i>Crepis vesicaria</i> L.	*	*	*	*	*	*	2	1	1	+	1	1	+	*	1
<i>Anthoxanthum odoratum</i> L.	1	+	*	+	*	*	*	*	+	+	1	1	1	1	+
<i>Arabis hirsuta</i> (L.) Scop.	*	*	*	*	*	*	+	+	+	+	1	1	1	*	+
<i>Teucrium chamaedrys</i> L.	+	+	*	+	*	+	*	*	*	*	+	+	+	1	+
<i>Prunus spinosa</i> L.	*	*	*	*	*	*	*	1	1	1	1	+	+	2	2
<i>Anemone hortensis</i> L.	*	*	*	*	*	*	*	+	1	*	1	+	1	+	+
<i>Festuca rubra</i> L.	*	*	+	2	+	*	1	*	*	*	*	*	*	2	1
<i>Asphodelus microcarpus</i> Salzm. et Viv.	*	*	*	*	*	*	2	*	*	*	+	1	1	2	+
<i>Cistus monspeliensis</i> L.	1	1	+	1	+	*	2	*	*	*	*	*	*	*	*
<i>Sanguisorba muricata</i> Gremler	*	*	*	+	*	*	*	*	1	+	1	1	+	*	*
<i>Silene latifolia</i> Poiret	*	*	*	*	*	*	*	*	+	*	+	1	1	+	+
<i>Cistus eriocephalus</i> Viv.	1	+	*	*	*	+	*	1	+	*	*	*	*	*	*
<i>Dorycnium hirsutum</i> (L.) Ser.	1	+	+	*	*	*	1	1	*	*	*	*	*	*	*
<i>Silene vulgaris</i> (Moench) Garcke	*	*	*	*	*	*	1	1	+	*	+	*	*	*	*
<i>Galium verum</i> L.	+	+	*	*	*	*	*	*	*	*	*	*	*	+	1
<i>Leontodon tuberosus</i> L.	*	*	*	*	*	*	+	*	*	*	*	*	+	1	+

fresche e umide, come il fondo di valloni o versanti bene ombreggiati, in genere si insediano leccete mesofile del *Quercion ilicis* riferibili in massima parte al *Pistacio-Quercetum ilicis* Brullo & Marcenò 1985. Nell'estrema punta salentina, limitatamente alle aree più interne, interessate da un bioclina termomediterraneo subumido superiore, la formazione climatofila è rappresentata dall' *Arbutio-Quercetum calliprini* Brullo *et al.* 1987. Abbastanza diffuse lungo le stazioni costiere della Puglia sono inoltre le formazioni dell' *Oleo-Ceratonion*, appartenenti per lo più al *Myrto-Lentiscetum* (Molinier 1954 em. O. Bolos 1962) Rivas Martinez 1975.

Da osservazioni personali inedite, nei processi di degradazione, legati soprattutto al taglio, incendio e pascolo, l'*Euphorbio apii-Quercetum trojanae* tende ad essere sostituito, in seguito alla distruzione dello strato arboreo ed arbustivo, da garighe del *Cisto-Ericion* Horvatić 1958, che a loro volta, con l'accentuarsi dei fenomeni di erosione del suolo, lasciano il posto a praticelli effimeri del *Thero-Brachypodion* Br.-Bl. 1925.

Sulla base dei dati di letteratura, boschi a *Quercus trojana* sono stati segnalati nella Penisola Balcanica da Horvat *et al.* (1974), che riportano per la Macedonia un *Quercetum trojanae* Em 1958 em. Horvat 1959 attribuito all' *Ostryo-Carpinion orientalis* Horvat (1954) 1958, alleanza dei *Quercetalia pubescentis* Br.-Bl. 1932. Questa associazione, per la presenza e in genere abbondanza di *Carpinus orientalis*, *Ostrya carpinifolia*, *Sorbus aria*, *Acer obtusatum*, *Quercus pubescens*, *Fraxinus ornus*, *Coronilla emeroides*, *Buxus sempervirens*, *Cornus mas*, *Festuca heterophylla*, *Campanula trachelium*, *Polygonatum odoratum*, ecc., come pure per le sue esigenze ecologiche, mostra un marcato carattere mesofilo e pertanto si differenzia nettamente dall'associazione pugliese. Altre associazioni a *Quercus trojana*, ascritte sempre all' *Ostryo-Carpinion orientalis*, sono state segnalate per varie località dell'ex Jugoslavia. Si tratta in particolare del *Quercetum pubescenti-macedonicae* Horvat 1936 n.n., *Quercetum trojanae hercegovinicum* Stefanović 1962 n.n., *Quercetum trojanae montenegrinum* Blečić & Lakusić 1975 n.n., *Quercetum macedonicae* Em in Horvat 1950 n.n. (cfr. Horvat 1936, 1950, Fukarek & Fabijanić 1968, Stefanović & Fabijanić 1969, Em 1969). Queste associazioni non possono essere comparate dal punto di vista floristico con quella pugliese in quanto i vari autori citano semplicemente il nome dell'associazione

senza fornire rilievi o elenchi floristici. E' comunque da ritenere probabile che boschi termofili a *Quercus trojana* affini a quelli pugliesi, rientranti anch'essi nel *Quercion ilicis* siano presenti anche nei territori balcanici e anatolici, soprattutto nelle stazioni collinari più meridionali.

Appendice 1

Ril. 1-6: Tra Putignano e Alberobello (Bari), 8.5.84; ril. 7 Bosco delle Pianelle (Taranto), 7.5.1992; ril. 8-10: Martinafranca (Taranto), 7.5.1992; ril. 11-13: Alberobello (Bari), 7.5.1992; ril. 14-15 Gioia del Colle (Bari), 7.5.1992.

Appendice 2

Specie sporadiche di Tab. 1 - Ril. 1: *Asperula aristata* L. Fil., *Briza maxima* L., *Hippocrepis varia*, *Oryzopsis miliacea* (L.) Asch. et Schweinf., *Pimpinella anisoides* Brignati, *Stachys arenaria* Vahl, *Thesium italicum* DC. Ril. 2: *Briza maxima* L., *Hippocrepis varia*, *Stachys arenaria* Vahl, *Thesium italicum* DC. Ril. 3: *Asperula aristata* L. Fil., *Pimpinella anisoides* Brignati, *Stachys arenaria* Vahl, *Saxifraga bulbifera* L. Ril. 4: *Geranium purpureum* Vill., *Lolium perenne* L., *Prunella vulgaris* L., *Saxifraga bulbifera* L. Ril. 6: *Briza maxima* L., *Geranium purpureum* Vill., *Lolium perenne* L., *Saxifraga bulbifera* L. ril. 7: *Cistus salvifolius* L., *Orchis morio* L., *Orchis papilionacea* L. Ril. 8: *Cistus salvifolius* L., *Festuca heterophylla* Lam., *Pteridium aquilinum* (L.) Kuhn Ril. 9: *Festuca heterophylla* Lam., *Pteridium aquilinum* (L.) Kuhn, *Platanthera chlorantha* (Custer) Rehb. Ril. 11: *Clinopodium arundanum* Boiss. Ril. 12: *Bromus erectus* Hudson, *Clinopodium arundanum* Boiss., *Hordeum bulbosum* L., *Ornithogalum comosum* L. Ril. 13: *Clinopodium arundanum* Boiss., *Festuca heterophylla* Lam., *Ornithogalum comosum* L.; *Pimpinella peregrina* L. Ril. 14: *Elaeoselinum asclepium* (L.) Bertol., *Galium aparine* L., *Ornithogalum comosum* L., *Pimpinella peregrina* L., *Quercus calliprinos* Webb, *Silene otites* (L.) Wibel, *Eryngium campestre* L., *Origanum heracleoticum* L. Ril. 15: *Pimpinella peregrina* L., *Quercus calliprinos* Webb, *Silene otites* (L.) Wibe, *Elaeoselinum asclepium* (L.) Bertol., *Galium aparine* L., *Thapsia garganica* L. *Genista* sp., *Rhamnus oleoides* L.

Riassunto

In questo lavoro vengono esaminati i boschi calcicoli caratterizzati dalla dominanza di *Quercus trojana* presenti in Puglia. Dal punto di vista fitosociologico questa formazione rientra nel *Quercion ilicis*, alleanza dei *Quercetalia ilicis*, localizzandosi nel territorio delle Murge a 100-500 m di quota. Si tratta di un'associazione climatofila legata a un bioclina di tipo mesomediterraneo subumido inferiore, la quale viene proposta come *Euphorbio apii-Quercetum trojanae*. Ne vengono esaminate la composizione floristica, l'ecologia, la distribuzione e le correlazioni sindinamiche.

Bibliografia

- Agostini R., 1967. *Aspetti fitosociologici delle pinete a Pinus halepensis Mill. del territorio di Taranto*. Arch. Bot. Biogeogr. Ital., 4, 43: 373-401.
- Bartolo G., Brullo S., Minissale P. & Spampinato G., 1985. *Osservazioni fitosociologiche sulle pinete a Pinus halepensis Miller del bacino del Fiume Tellaro (Sicilia sud-orientale)*. Boll. Acc. Gioenia Sci. Nat., 18 (325): 255-270.
- Bianco P., 1958. *Querceti a Quercus trojana Webb nel territorio di San Michele di Bari*. Giorn. Bot. Ital., 65: 43-100.
- Bianco P., 1961a. *Ricerche sul ciclo riproduttivo di specie del genere Quercus della flora italiana. VI - contributo alla biologia di Quercus trojana Webb in Puglia*. Ann. Acc. Ital. Sc. Forest., 10: 59-96.
- Bianco P., 1961b. *Studio biometrico su Quercus trojana Webb della Puglia*. Ann. Acc. Ital. Sc. Forest., 10: 115-133.
- Bianco P., Brullo S., Pignatti E. & Pignatti S. 1988. *La vegetazione delle rupi calcaree della Puglia*. Braun-Blanquetia, 2: 133-151.
- Brullo S. & Marcenò C., 1985. *Contributo alla conoscenza della classe Quercetea ilicis in Sicilia*. Not. Fitosoc., 19(1): 183-229.
- Brullo S., Minissale P., Signorello P. & Spampinato G., 1987. *Studio fitosociologico delle garighe ad Erica manipuliflora del Salento (Puglia meridionale)*. Arch. Bot. Biogeogr. Ital., 62: 201-214.
- Chiesura Lorenzoni F., Curti L. & Lorenzoni G. G., 1971. *Considerazioni sulle cenosi a Quercus calliprinos Webb e a Q. macedonica DC. in Puglia*. Atti I Simposio Nazionale sulla conservazione della Natura. Bari, 255-262.
- Corti R. & Magini E., 1955. *Fragno*. In Pavari A. (ed.), *Le Cupulifere*. Monti & Boschi, 6: 553-556.
- Crivellari D., 1950. *Inchiesta sulla distribuzione del genere Quercus in Puglia*. Giorn. Bot. Ital., 57: 335-350.
- De Marco G. & Caneva G., 1985. *Analisi sintassonomica fitogeografica comparata di alcune significative cenosi a Pinus halepensis Mill. in Italia*. Not. Fitosoc. 19: 155-176.
- D'Amato F., 1949. *Sull'areale pugliese della Quercus trojana Webb*. Atti Soc. Tosc. Sc. Nat., 66B: 89-114.
- Em H., 1969. *Der Buchsbaum (Buxus sempervirens) in der Vegetation Macedoniens*. Mitt. Ostalp.-din. Pflanzensoz. Arbeitsgem. 9: 311-322.
- Francini Corti E., 1966. *Aspetti della vegetazione pugliese e contingente paleoegeico meridionale nella Puglia*. Ann. Acc. Ital. Sc. Forest., 15: 137-193.
- Fukarek P. & Fabijanić B., 1968. *Versuch einer pflanzensoziologischen Gliederung der Wald- und Šibliak-Gesellschaften Bosniens und der Hercegovina*. In: Tüxen R. (ed.), *Pflanzensoziologische Systematik*. Den Haag, 112-123.
- Hedge I.C. & Yaltirik P. H., 1982. *Quercus*. In: , Davis P.H., Edmondson J.R., Miel R.R. & Kit Tan (eds.), *Flora of Turkey and the East Aegean Islands*, vol. 7: 659-683.
- Horvat I., 1936. *Istraživanje vegetacije planina Vardarske banovine*, 2. Ljet. Jugosl. Akad. Zn. Unj. Zagreb, 48: 211-227.
- Horvat I., 1950. *Les associations forestières en Yugoslavie*. Inst. šum. istr. Zagreb.
- Horvat I., Glavac V. & Ellenberg H., 1974. *Vegetation Südsteuropas*. Geobotanica selecta. 4. Fisher, Stuttgart.
- Jalas J. & Suominen J., 1976. *Atlas Florae Europaeae*. 3: 69. Helsinki.
- Linzalone M., 1955. *Boschi misti a Quercus trojana Webb e Quercus pubescens W. a sud di Gioia del Colle*. Giorn. Bot. Ital. 62: 468-477.
- Lorenzoni G. G. & Chiesura Lorenzoni F., 1987. *First phytosociological interpretation of Quercus trojana Webb vegetation in the Murge region (Bari - Taranto - South Italy)*. Acta Bot. Croat., 46: 95-103.
- Macchia F., 1987. *Il fitoclima del Salento*. Not. Fitosoc. 19: 29-60.
- Maselli V.G., 1940. *Contributo alla conoscenza delle querce d'Italia. Il fragno*. Riv. Forest. Ital., 2: 20-35.
- Rivas-Martinez S., 1975. *La vegetation de la clase Quercetea ilicis en España y Portugal*. Anal. Inst. Bot. Cavanilles 31: 205-259.
- Rivas-Martinez S., Bascónes J.C., Díaz T.E., Gonzáles F.F. & Loidi J., 1991. *Vegetación del Pirineo occidental y Navarra*. Itinera Geobot. 5: 5-456.
- Solazzo E., 1955. *Boschi misti a Quercus trojana Webb e Quercus pubescens W. nel territorio di Acquaviva delle fonti*. Giorn. Bot. Ital. 62: 487-498.
- Stefanović V. & Fabijanić B., 1969. *Zerreiche und Waldgesellschaften mit Zerreiche des Dinarischen und Apenninischen Gebietes*. Mitt. Ostalp.-din. Pflanzensoz. Arbeitsgem. 9: 287-299.

Received: December 11, 1996

Accepted: October 17, 1997

fresche e umide, come il fondo di valloni o versanti bene ombreggiati, in genere si insediano leccete mesofile del *Quercion ilicis* riferibili in massima parte al *Pistacio-Quercetum ilicis* Brullo & Marcenò 1985. Nell'estrema punta salentina, limitatamente alle aree più interne, interessate da un bioclina termomediterraneo subumido superiore, la formazione climatofila è rappresentata dall' *Arbuto-Quercetum calliprini* Brullo *et al.* 1987. Abbastanza diffuse lungo le stazioni costiere della Puglia sono inoltre le formazioni dell' *Oleo-Ceratonion*, appartenenti per lo più al *Myrto-Lentiscetum* (Molinier 1954 em. O. Bolos 1962) Rivas Martinez 1975.

Da osservazioni personali inedite, nei processi di degradazione, legati soprattutto al taglio, incendio e pascolo, l'*Euphorbio apii-Quercetum trojanae* tende ad essere sostituito, in seguito alla distruzione dello strato arboreo ed arbustivo, da garighe del *Cisto-Ericion* Horvatić 1958, che a loro volta, con l'accentuarsi dei fenomeni di erosione del suolo, lasciano il posto a praticelli effimeri del *Thero-Brachypodion* Br.-Bl. 1925.

Sulla base dei dati di letteratura, boschi a *Quercus trojana* sono stati segnalati nella Penisola Balcanica da Horvat *et al.* (1974), che riportano per la Macedonia un *Quercetum trojanae* Em 1958 em. Horvat 1959 attribuito all' *Ostryo-Carpinion orientalis* Horvat (1954) 1958, alleanza dei *Quercetalia pubescentis* Br.-Bl. 1932. Questa associazione, per la presenza e in genere abbondanza di *Carpinus orientalis*, *Ostrya carpinifolia*, *Sorbus aria*, *Acer obtusatum*, *Quercus pubescens*, *Fraxinus ornus*, *Coronilla emeroides*, *Buxus sempervirens*, *Cornus mas*, *Festuca heterophylla*, *Campanula trachelium*, *Polygonatum odoratum*, ecc., come pure per le sue esigenze ecologiche, mostra un marcato carattere mesofilo e pertanto si differenzia nettamente dall'associazione pugliese. Altre associazioni a *Quercus trojana*, ascritte sempre all' *Ostryo-Carpinion orientalis*, sono state segnalate per varie località dell'ex Jugoslavia. Si tratta in particolare del *Quercetum pubescenti-macedonicae* Horvat 1936 n.n., *Quercetum trojanae hercegovanicum* Stefanović 1962 n.n., *Quercetum trojanae montenegrinum* Blečić & Lakusić 1975 n.n., *Quercetum macedonicae* Em in Horvat 1950 n.n. (cfr. Horvat 1936, 1950, Fukarek & Fabijanić 1968, Stefanović & Fabijanić 1969, Em 1969). Queste associazioni non possono essere comparate dal punto di vista floristico con quella pugliese in quanto i vari autori citano semplicemente il nome dell'associazione

senza fornire rilievi o elenchi floristici. E' comunque da ritenere probabile che boschi termofili a *Quercus trojana* affini a quelli pugliesi, rientranti anch'essi nel *Quercion ilicis* siano presenti anche nei territori balcanici e anatolici, soprattutto nelle stazioni collinari più meridionali.

Appendice 1

Rill. 1-6: Tra Putignano e Alberobello (Bari), 8.5.84; ril. 7 Bosco delle Pianelle (Taranto), 7.5.1992; rill. 8-10: Martinafranca (Taranto), 7.5.1992; rill. 11-13: Alberobello (Bari), 7.5.1992; rill. 14-15 Gioia del Colle (Bari), 7.5.1992.

Appendice 2

Specie sporadiche di Tab. 1 - Ril. 1: *Asperula aristata* L. Fil., *Briza maxima* L., *Hippocrepis varia*, *Oryzopsis miliacea* (L.) Asch. et Schweinf., *Pimpinella anisoides* Brignati, *Stachys arenaria* Vahl, *Thesium italicum* DC. Ril. 2: *Briza maxima* L., *Hippocrepis varia*, *Stachys arenaria* Vahl, *Thesium italicum* DC. Ril. 3: *Asperula aristata* L. Fil., *Pimpinella anisoides* Brignati, *Stachys arenaria* Vahl, *Saxifraga bulbifera* L. Ril. 4: *Geranium purpureum* Vill., *Lolium perenne* L., *Prunella vulgaris* L., *Saxifraga bulbifera* L. Ril. 6: *Briza maxima* L., *Geranium purpureum* Vill., *Lolium perenne* L., *Saxifraga bulbifera* L. ril. 7: *Cistus salvifolius* L., *Orchis morio* L., *Orchis papilionacea* L. Ril. 8: *Cistus salvifolius* L., *Festuca heterophylla* Lam., *Pteridium aquilinum* (L.) Kuhn Ril. 9: *Festuca heterophylla* Lam., *Pteridium aquilinum* (L.) Kuhn, *Platanthera chlorantha* (Custer) Rchb. Ril. 11: *Clinopodium arundanum* Boiss. Ril. 12: *Bromus erectus* Hudson, *Clinopodium arundanum* Boiss., *Hordeum bulbosum* L., *Ornithogalum comosum* L. Ril. 13: *Clinopodium arundanum* Boiss., *Festuca heterophylla* Lam., *Ornithogalum comosum* L.; *Pimpinella peregrina* L. Ril. 14: *Elaeoselinum asclepium* (L.) Bertol., *Galium aparine* L., *Ornithogalum comosum* L., *Pimpinella peregrina* L., *Quercus calliprinos* Webb, *Silene otites* (L.) Wibel, *Eryngium campestre* L., *Origanum heracleoticum* L. Ril. 15: *Pimpinella peregrina* L., *Quercus calliprinos* Webb, *Silene otites* (L.) Wibe, *Elaeoselinum asclepium* (L.) Bertol., *Galium aparine* L., *Thapsia garganica* L. *Genista* sp., *Rhamnus oleoides* L.

Riassunto

In questo lavoro vengono esaminati i boschi calcicoli caratterizzati dalla dominanza di *Quercus trojana* presenti in Puglia. Dal punto di vista fitosociologico questa formazione rientra nel *Quercion ilicis*, alleanza dei *Quercetalia ilicis*, localizzandosi nel territorio delle Murge a 100-500 m di quota. Si tratta di un'associazione climatofila legata a un bioclina di tipo mesomediterraneo subumido inferiore, la quale viene proposta come *Euphorbio apii-Quercetum trojanae*. Ne vengono esaminate la composizione floristica, l'ecologia, la distribuzione e le correlazioni sindinamiche.

Bibliografia

- Agostini R., 1967. *Aspetti fitosociologici delle pinete a Pinus halepensis Mill. del territorio di Taranto*. Arch. Bot. Biogeogr. Ital., 4, 43: 373-401.
- Bartolo G., Brullo S., Minissale P. & Spampinato G., 1985. *Osservazioni fitosociologiche sulle pinete a Pinus halepensis Miller del bacino del Fiume Tellaro (Sicilia sud-orientale)*. Boll. Acc. Gioenia Sci. Nat., 18 (325): 255-270.
- Bianco P., 1958. *Querceti a Quercus trojana Webb nel territorio di San Michele di Bari*. Giorn. Bot. Ital., 65: 43-100.
- Bianco P., 1961a. *Ricerche sul ciclo riproduttivo di specie del genere Quercus della flora italiana. VI - contributo alla biologia di Quercus trojana Webb in Puglia*. Ann. Acc. Ital. Sc. Forest., 10: 59-96.
- Bianco P., 1961b. *Studio biometrico su Quercus trojana Webb della Puglia*. Ann. Acc. Ital. Sc. Forest., 10: 115-133.
- Bianco P., Brullo S., Pignatti E. & Pignatti S. 1988. *La vegetazione delle rupi calcaree della Puglia*. Braun-Blanquetia, 2: 133-151.
- Brullo S. & Marcenò C., 1985. *Contributo alla conoscenza della classe Quercetea ilicis in Sicilia*. Not. Fitosoc., 19(1): 183-229.
- Brullo S., Minissale P., Signorello P. & Spampinato G., 1987. *Studio fitosociologico delle garighe ad Erica manipuliflora del Salento (Puglia meridionale)*. Arch. Bot. Biogeogr. Ital., 62: 201-214.
- Chiesura Lorenzoni F., Curti L. & Lorenzoni G. G., 1971. *Considerazioni sulle cenosi a Quercus calliprinos Webb e a Q. macedonica DC. in Puglia*. Atti I Simposio Nazionale sulla conservazione della Natura. Bari, 255-262.
- Corti R. & Magini E., 1955. *Fragno*. In Pavari A. (ed.), *Le Cupulifere*. Monti & Boschi, 6: 553-556.
- Crivellari D., 1950. *Inchiesta sulla distribuzione del genere Quercus in Puglia*. Giorn. Bot. Ital., 57: 335-350.
- De Marco G. & Caneva G., 1985. *Analisi sintassonomica fitogeografica comparata di alcune significative cenosi a Pinus halepensis Mill. in Italia*. Not. Fitosoc. 19: 155-176.
- D'Amato F., 1949. *Sull'areale pugliese della Quercus trojana Webb*. Atti Soc. Tosc. Sc. Nat., 66B: 89-114.
- Em H., 1969. *Der Buchsbaum (Buxus sempervirens) in der Vegetation Macedoniens*. Mitt. Ostalp.-din. Pflanzensoz. Arbeitsgem. 9: 311-322.
- Francini Corti E., 1966. *Aspetti della vegetazione pugliese e contingente paleoegeico meridionale nella Puglia*. Ann. Acc. Ital. Sc. Forest., 15: 137-193.
- Fukarek P. & Fabijanić B., 1968. *Versuch einer pflanzensoziologischen Gliederung der Wald- und Šibliak-Gesellschaften Bosniens und der Hercegovina*. In: Tüxen R. (ed.), *Pflanzensoziologische Systematik*. Den Haag, 112-123.
- Hedge I.C. & Yaltirik P. H., 1982. *Quercus*. In: , Davis P.H., Edmondson J.R., Miel R.R. & Kit Tan (eds.), *Flora of Turkey and the East Aegean Islands*, vol. 7: 659-683.
- Horvat I., 1936. *Istraživanje vegetacije planina Vardarske banovine*, 2. Ljet. Jugosl. Akad. Zn. Unj. Zagreb, 48: 211-227.
- Horvat I., 1950. *Les associations forestières en Yugoslavie*. Inst. šum. istr. Zagreb.
- Horvat I., Glavac V. & Ellenberg H., 1974. *Vegetation Südsteuropas*. Geobotanica selecta. 4. Fisher, Stuttgart.
- Jalas J. & Suominen J., 1976. *Atlas Florae Europaeae*. 3: 69. Helsinki.
- Linzalone M., 1955. *Boschi misti a Quercus trojana Webb e Quercus pubescens W. a sud di Gioia del Colle*. Giorn. Bot. Ital. 62: 468-477.
- Lorenzoni G. G. & Chiesura Lorenzoni F., 1987. *First phytosociological interpretation of Quercus trojana Webb vegetation in the Murge region (Bari - Taranto - South Italy)*. Acta Bot. Croat., 46: 95-103.
- Macchia F., 1987. *Il fitoclima del Salento*. Not. Fitosoc. 19: 29-60.
- Maselli V.G., 1940. *Contributo alla conoscenza delle querce d'Italia. Il fragno*. Riv. Forest. Ital., 2: 20-35.
- Rivas-Martinez S., 1975. *La vegetation de la clase Quercetea ilicis en España y Portugal*. Anal. Inst. Bot. Cavanilles 31: 205-259.
- Rivas-Martinez S., Bascónes J.C., Díaz T.E., Gonzáles F.F. & Loidi J., 1991. *Vegetación del Pirineo occidental y Navarra*. Itinera Geobot. 5: 5-456.
- Solazzo E., 1955. *Boschi misti a Quercus trojana Webb e Quercus pubescens W. nel territorio di Acquaviva delle fonti*. Giorn. Bot. Ital. 62: 487-498.
- Stefanović V. & Fabijanić B., 1969. *Zerreiche und Waldgesellschaften mit Zerreiche des Dinarischen und Apenninischen Gebietes*. Mitt. Ostalp.-din. Pflanzensoz. Arbeitsgem. 9: 287-299.

Received: December 11, 1996

Accepted: October 17, 1997

ON THE PRESENCE OF PHYTOSTEROLS IN SOME SPECIES OF *EUPHORBIA* L.

Laura COASSINI LOKAR and Fabrizio MARTINI

Dipartimento di Biologia, Università di Trieste, Via L. Giorgieri 10, I-34127 Trieste

Keywords: Chemotaxonomy, *Euphorbia*, Friuli- Venezia Giulia, Phytosterols.

Abstract: In five species of *Euphorbia* of the Friuli-Venezia Giulia region (NE Italy) (*E. amygdaloides*, *E. cyparissias*, *E. helioscopia*, *E. fragifera* and *E. flavicoma* subsp. *verrucosa*) ten phytosterols were isolated and identified, by means of HPLC and spectroscopic techniques, as β -sitosterol, stigmasterol, campesterol, colestano-3, 5, 6-triol, cycloartenol, squalene, stigmastanol, stigmasterol acetate, cholesterol and lanosterol. The ratio β -sitosterol/stigmasterol can be used to distinguish some of the species.

Introduction

The interest in plant sterols (phytosterols) has been growing over the last few years because of their biological importance. Numerous studies show that phytosterols are widespread, from the simplest organisms such as bacteria, algae and fungi (Chardon-Loriaux *et al.* 1976), to vascular plants, where they were isolated both in the vegetative and reproductive systems (Ingram *et al.* 1968). Recent studies have also identified them in the genus *Euphorbia* (Atallah & Nicholas 1972, Nielsen *et al.* 1979, Sekula & Nes 1980), together with various other secondary metabolites such as triterpenes, flavonoids, resins and essential oils. The highest concentrations were found in chloroplasts, microsomes and mitochondria, and in the plasma-lemma (Nes & Heftmann 1981), where they probably are structural elements of the membranes, regulating their stabilisation and the permeability (Grunwald 1968 and 1971). Studies were also carried out on the biosynthesis and the functions of phytosterols (Sharma 1970, Heftmann 1971) and of some derivatives, in particular sterol esters (Waller *et al.* 1981) and sterol glycosides (Smith 1969). Their concentration varies during the life cycle, as proved by the studies of Davis (1972) and Grunwald (1975) on tobacco, and of Ingram *et al.* (1968) on some *Cruciferae*. Germination (Bush & Grunwald 1972), growth (Geuns 1975), anthesis (Biswas *et al.* 1967) and senescence (Grunwald 1975) are significant stages in the biosynthesis of phytosterols, whose concentration is also influenced by environmental factors such as water availability,

temperature (Davis & Finkner 1972) and light (Bush *et al.* 1971). Moreover, the concentration of sterols particularly increases during the germination of the seeds, which seems to be linked to the biogenesis of the membranes (Duperon 1971; Bush & Grunwald 1972).

According to some authors (Atallah *et al.* 1975), all plant tissues can potentially synthesize sterols, but the degree of biosynthesis depends on the type and on the age of the tissue (Geuns 1975). As regards the pharmacological properties of phytosterols, Gupta *et al.* (1980) demonstrated their anti-inflammatory and antipyretic properties. Experiments on their contraceptive effects, due to stigmast-5ene-3-7 diol and to stigmast-5ene-3, 7 diol isolated in leaves of *Ananas comosus*, were conducted on Guinea pigs by Pakrashi *et al.* (1975). Phytosterols also have antitumoral and cytotoxic (Miles *et al.* 1974), antihyper-cholesteremic (Chandler *et al.* 1979), anticonvulsant (Farnsworth & Cordell 1976), antibacterial, antitubercular and expectorant effects (Benoit *et al.* 1976).

Examined species

The genus *Euphorbia* comprises about 700 species and has a world-wide distribution (except Antarctica), especially in hot and dry areas. They are herbaceous annuals and perennials, suffrutescent plants and trees, mainly with simple leaves. Thermophytic elements prevail also among the almost thirty species of the flora of the Friuli-

Venezia Giulia region (Pignatti 1982, Poldini 1980, 1991), the highest concentration of the species being in the Karst area around Trieste and Gorizia, with a submediterranean climate (Poldini *et al.* 1991).

Our study is based on five among the most widespread species of the subgenus *Esula* Pers.: *E. amygdaloides* L. and *E. cyparissias* L., of the sect. *Esula*; *E. helioscopia* L., *E. fragifera* Jan and *E. flavicoma* DC. subspecies *verrucosa* (Fiori) Pign. of the sect. *Helioscopia* Dumort.

E. amygdaloides is found growing all over Europe (Meusel *et al.* 1978). In the Friuli-Venezia Giulia region it mainly occurs in the mountains, but it also occurs sporadically in the lowlands (Poldini 1991).

The distribution of *E. cyparissias*, originally limited to central-southern and south-eastern Europe has become wider because of anthropization, and it now extends to the United States and New Zealand (Hult n & Fries 1986). In north-eastern Italy, it is very common in open, stony grasslands.

E. helioscopia, the only annual species considered, occurs in Europe and western Asia (Meusel *et al.* 1978). It is mainly found growing in synanthropic stands on loose, well-manured soil from sea level to about 1300 m.

The Illyrian endemic *E. fragifera* occurs from Montenegro to the Trieste Karst, where it is an important element of the sage garrigue. Two chemical varieties were distinguished in the Trieste Karst on the basis of essential oils composition. The different terpenoidic composition is probably correlated to different pollinators (Coassini Lokar *et al.* 1986).

Between the two intraspecies varieties of *E. flavicoma*, only subsp. *verrucosa*, with a Mediterranean-Pontic-Illyrian distribution, occurs in the Friuli Venezia-Giulia region (Pignatti 1982, Poldini 1991), usually in dry grasslands on limestone.

Data and Methods

Plant material - The samples (30 plants of each species at full anthesis) were randomly collected in the following areas of the Karst of Trieste (NE Italy): *E. amygdaloides*: Zolla di Monrupino and Mt. Orsario, 330 m; *E. cyparissias*: Opicina, 330 m; *E. fragifera*: Prosecco, 250 m; *E. flavicoma* subsp. *verrucosa*: between Gabrovizza and Sgonico, 260 m; *E. helioscopia*: Trieste, Scorcola, 100 m.

Voucher specimens were deposited in the Herbarium of the Department of Biology, University of Trieste (TSB). Nomenclature follows Pignatti (1982).

Preparation of crude extracts - The samples (epigeal and aerial organs) were dried at 50°C until constant weight and pulverized under liquid nitrogen in a stainless steel ball-mill prior to extraction. 5 g of each pulverized sample accurately weighted and exhaustively extracted according to the method proposed by Ghosh *et al.* (1985), were shaken again with a mixture of chloroform: methanol: water (30:60:30) in the ratio 5:120 p/V at room temperature (exhaustion control by preparative TLC). After centrifugation (20,000 rp/m for 10 min) the supernatant was separated from the methanolic phase with chloroform, evaporated *in vacuo* at 30°C; the residue was dissolved in 10 ml of n-hexane for HPLC in defined volumes (10 ml) and used directly for analysis. The purity and identity of the peaks in the crude extracts were controlled by UV spectra detection and compared with the standards (SIGMA Chem. Co., St. Louis, USA). The standards and the crude extracts were filtered on Millipore before the injection (7µ 1/10 ml of crude extract).

HPLC equipment - We used a Spectra Physic M-8750, Spectra Physic Programmer M 8700, septumless injector Model 8750, with a 10 µl sample loop, Bio Rad BIO-SIL-HP10 column (4×250 mm i.d. with BIORAD HP-10 pre-column, Spectra Physic SP Mod. 8840 wavelength detector, multiple Mega Series (C. Erba, Milano, Italy) Computing integrator.

Operating conditions - Flow rate 1,5 ml/min, room temperature, detection UV 220 nm (detector sensitivity 0.01 a.u.f.s.), PSI 150, chart speed 10 mm/min; all the determination were performed at room temperature (22°C) and at isocratic condition of elution. Standard deviation for retention times values 0.4-1.0 %; peak area reproducibility (for 5 analyses of standards) 1.9-3.1 %; mobile phase: n-hexane/ isopropanol/methanol (96:3.5:0.5 v/v).

Results

Ten of the phytosterols contained in aerial and hypogeal parts of the various samples of the five *Euphorbia* species considered were identified and quantified, as mean percentages (Tab.1): β-sitosterol, stigmasterol, campesterol, colestano-3β, 5α,

Tab.1 - Composition in phytosterols (% dry weight) in epigeal and hypogeal parts of the populations of the five species of *Euphorbia*.

Phytosterols	<i>E. amygdaloides</i>		<i>E. fragifera</i>		<i>E. helioscopia</i>		<i>E. cyparissias</i>		<i>E. verrucosa</i>	
	epig. %	hypog. %	epig. %	hypog. %	epig. %	hypog. %	epig. %	hypog. %	epig. %	hypog. %
β -sitosterol	88.95	84.61	93.97	79.53	79.90	71.92	64.28	50.30	83.50	37.95
stigmasterol	6.35	6.62	2.77	2.20	12.34	16.93	27.13	18.28	12.27	15.53
campesterol	2.20	2.53	1.94	3.09	3.09	4.21	5.01	6.78	1.78	6.49
coestan - 3 β , 5 α , 6 β - triol	1.17	2.14	-	3.65	2.41	1.75	-	20.32	0.28	9.63
cycloartenol	0.11	0.31	0.05	0.32	0.76	0.90	0.71	1.00	1.02	1.51
squalene	-	1.64	-	0.32	-	1.75	0.13	0.03	-	0.82
stigmastanol	-	-	0.05	0.58	0.83	1.27	2.62	2.91	1.07	1.33
stigmasterol acetate	1.22	2.15	1.22	-	-	-	-	-	-	16.07
cholesterol	-	-	-	0.19	-	-	-	-	-	-
lanosterol	-	-	-	-	0.04	-	-	-	-	-
Σ others unidentified	-	-	-	7.65	-	0.80	0.12	0.38	0.08	2.81

6 β -triol, cycloartenol, squalene, stigmastanol, stigmasterol acetate, cholesterol and lanosterol. β -sitosterol, stigmasterol and campesterol are present both in epigeal and hypogeal organs, with β -sitosterol prevailing in the aerial parts. Stigmasterol is present in lower percentages, and generally prevails in the hypogeal parts with the exception of *E. fragifera* and *E. cyparissias*, which present the greatest percentages. Campesterol and cycloartenol are always most abundant in the hypogeal parts. Squalene, in very low quantities, was only found in the hypogeal parts of all samples. Stigmastanol is absent only in *E. amygdaloides* and tends to prevail in the hypogeal parts. Stigmasterol- acetate is scarcely represented, with the exception of the hypogeal parts of *E. verrucosa*. The hypogeal parts of *E. cyparissias* and *E. verrucosa* contained the highest percentages of coestan- 3 β - 5 α - 6 β - triol.

Small quantities of lanosterol (in *E. helioscopia*) and of cholesterol (*E. fragifera*) were isolated. These triterpenic alcohols were also isolated by Falsone (1972) and Falsone & Schneider (1985) in *E. biglandulosa* Desf., together with their acetylic derivatives.

The relative position of the five species may be defined by the chemical variables, taken three by three. The analysis proved that they are discriminated only by the quantities of β -sitosterol, stigmasterol, Σ others in the hypogeal parts (Fig. 1A) and by the quantities of β -sitosterol, stigmasterol, campesterol in their epigeal organs (Fig. 1B).

If we examine the sterolic composition of the various plant organs, there is evidence that, as already established by Grunwald (1975), the ratio β -sitosterol/ stigmasterol in the hypogeal and aerial




Fig. 1 - Triangular diagram showing the phytosterol discriminant combination of epigeal (A) and of hypogeal (B) organs of the five examined species (A = *E. amygdaloides*; C = *E. cyparissias*; F = *E. fragifera*; H = *E. helioscopia*; V = *E. verrucosa*).

Tab. 2 - Ratio β -sitosterol/stigmasterol in epigeal and hypogeal parts in the populations of the examined species of *Euphorbia*.

Species	β - sitosterol/stigmasterol	
	epigeal	hypogeal
<i>E. fragifera</i>	33.92	36.15
<i>E. amygdaloides</i>	14.00	12.76
<i>E. helioscopia</i>	6.47	4.25
<i>E. verrucosa</i>	6.27	2.44
<i>E. cyparissias</i>	2.37	2.75

parts (Tab. 2) is effective to distinguish the examined species. If both hypogeal and aerial organs are considered, *E. amygdaloides*, *E. fragifera* and *E. cyparissias* are clearly distinguished in their epigeal parts, while *E. amygdaloides* and *E. fragifera* are distinguished by the ratio value in the hypogeal parts. *E. helioscopia* and *E. verrucosa* are distinguishable only on the basis of the ratio value in the hypogeal parts.

Conclusions

This study was focused on the qualitative and quantitative presence of sterols in some populations of five *Euphorbia* species occurring in the Friuli-Venezia Giulia region, analyzed in their anthesis phase. It highlighted the high productivity of these biologically active compounds, and in particular of β -sitosterol, which tends to prevail in the green parts of the plants, especially in the leaves, where, on the other side, the biosynthesis takes place. Most of the identified sterols may play an active role in the florigenic activity, particularly β -sitosterol and lanosterol, whose concentration of 100 p.p.m. is sufficient to lead to flowering in 50% of the plants of *Chrysanthemum* (Biswas *et al.* 1967). The presence of single sterol derivatives was proved to be linked to developmental stages, and a particular value was attributed to the ratio β -sitosterol /stigmasterol (Grunwald 1975). In our study, the value of the ratio remains sufficiently constant in the populations, both in epi- and hypogeal parts, and may be utilized for the discrimination at population level. Lanosterol was only present in *E. helioscopia*, and therefore is a significant discriminant. Lanosterol and its ester derivatives, which are not usually present in plant extracts, are also typical of other *Euphorbiae* with a more southern distribution

(Falsone 1972, Falsone & Schneider 1985); lanosterol and cycloartenol may result from the primary cyclization of squalene and are often found as acetylic derivatives, especially in latex. The 4-demethylsterols (β -sitosterol, campesterol and stigmasterol) prevail in the examined species, while saturated compounds (= stanols) are present in smaller amounts. Among sterol-esters, only stigmasterol-acetate was found.

It would be interesting to continue this analysis on other species of the genus, to define the composition of steroidal derivatives in plants and in latex alone, where ester derivatives and triterpenic alcohols seem to prevail, probably because of their highest solubility in water, which makes them less linked to the membrane processes. This could contribute to understand their role in various parts of the plant, and to establish the chemosystematic value of some of them inside the genus, or among other critical entities, such as *E. characias* and *E. wulfenii*.

Acknowledgments

Financial support was provided by the Italian M.U.R.S.T. (Ministry of Scientific Research and Technology) 40% to Prof. L. Coassini Lokar.

References

- Atallah A.M. & Nicholas H.J., 1972. Triterpenoids and steroids of *Euphorbia pilulifera*. *Phytochemistry*, 11: 1860.
- Atallah A.M., Aixel R.T., Ramsey R.B. & Nicholas H.J., 1975. Biosynthesis of sterols and triterpenes in *Pelargonium hortorum*. *Phytochemistry*, 14: 1529-1533.
- Benoit P.S., Fong H.H.S., Svoboda G.H. & Farnsworth N.R., 1976. Biological and phytochemical evaluation of plants. XIV. Anti-inflammatory evaluation of 163 species of plants. *Lloydia*, 39: 160-171.
- Biswas P.K., Paul K.B. & Henderson J.H.M., 1967. Effects of steroids on *Chrysanthemum* in relation to growth and flowering. *Nature*, 213: 917-918.
- Bush P.B. & Grunwald C., 1972. Sterol changes during germination of *Nicotiana tabacum* seeds. *Pl. Physiol.*, 50: 69-72.
- Bush P.B., Grunwald C. & Davis D.L., 1971. Changes in sterol composition during greening of etiolated barley shoots. *Pl. Physiol.*, 47: 745-749.
- Chandler R.F., Hooper S.N. & Ismail H.A., 1979. Antihypercholesteremic studies with sterols: dihydrocholesterol and 7-dehydrocholesterol. *Can. J. Pharm. Sci.*, 14: 61-62.
- Chardon-Loriaux I., Morisaki M. & Ikekawa N., 1976. Sterol profiles of red algae. *Phytochemistry*, 15: 723-725.
- Coassini Lokar L., Maurich V. & Poldini L., 1986. Chemical aspect of floral biology in *Euphorbia fragifera*. *Folia Geobot.*

- Phytotax., 21: 277-285.
- Davis D.L. & Finkner V.C., 1972. *Influence of temperature on sterol biosynthesis in Triticum aestivum*. Pl. Physiol., 52: 324-326.
- Davis D.L., 1972. *Sterol distribution within green and air cured tobacco*. Phytochemistry, 11: 489-494.
- Dupéron P., 1971. *Nature et comportement des stérols "libres" et esterifiés, au cours de la germination de divers types de semences. Hypothèses sur le rôle de ces substances chez les végétaux*. Physiol. Vég., 9: 373-399.
- Falsone G., 1972. *Inhaltsstoffe von Euphorbiaceae. Isolierung, Strukturaufklärung, biologische und pharmakologische Untersuchungen von Inhaltsstoffen des Milchsafte von Euphorbia biglandulosa*. Desf. Habilitationsschrift Mathematisch - Naturwissenschaftliche Fakultät, Universität Düsseldorf.
- Falsone G. & Schneider C., 1985. *Über Inhaltsstoffe von Euphorbiaceae, 8 Mitt. Triterpenester und Triterpenalkohole aus Euphorbia biglandulosa Desf.* Z. Naturforsch., 40b: 553-555.
- Farnsworth N.R. & Cordell G.A., 1976. *A review of some biologically active compounds isolated from plants as reported in the 1974-1975 literature*. Lloydia, 39: 420-455.
- Geuns J.M.C., 1975. *Regulation of sterol biosynthesis in etiolated mung bean hypocotyl sections*. Phytochemistry, 24: 1725-1727.
- Ghosh A., Misra S., Dutta A.K. & Choudhury A., 1985. *Pentacyclic triterpenoids and sterols from seven species of mangrove*. Phytochemistry, 24 (8): 1725-1727.
- Grunwald C., 1968. *Effect of sterols on the permeability of alcohol-treated red beet tissue*. Pl. Physiol., 43: 484-488.
- Grunwald C., 1971. *Effect of sterols, sterol ester and sterol glycoside in membrane permeability*. Pl. Physiol., 48: 653-655.
- Grunwald C., 1975. *Phytosterols in tobacco leaves at various stages of physiological maturity*. Phytochemistry, 14: 79-82.
- Gupta M.B., Nath R., Srivastava N., Shanker K., Kishor K. & Bhargava K.P., 1980. *Anti-inflammatory and antipyretic activities of β -sitosterol*. Pl. Med., 39: 157-163.
- Heftmann E., 1971. *Function of sterols in plants*. Lipids, 6: 128-133.
- Hultén E. & Fries M., 1986. *Atlas of North European Vascular Plants*. Königstein.
- Ingram D.S., Knights B.A., Mc Evoy I.J. & Mc Kay P., 1968. *Studies in the Cruciferae. Changes in the composition of the sterol fraction following germination*. Phytochemistry, 7: 1241-1245.
- Meusel H., Jäger E., Rauschert S. & Weinert E., 1978. *Vergleichende Chorologie der zentraleuropäischen Flora*. 2. Jena.
- Miles D.H., Kokpol U., Zalkow L.H., Steindel S.J. & Nabors J.B., 1974. *Tumor inhibitors I: preliminary investigation of antitumor activity of Sarracenia flava*. J. Pharm. Sci., 63: 613-615.
- Nes W.D. & Heftmann E., 1981. *A comparison of triterpenoids with steroids as membrane components*. J. Nat. Prod., 44: 377-400.
- Nielsen P.E., Nishimura H., Liang Y. & Calvin M., 1979. *Steroids from Euphorbia and other latex bearing plants*. Phytochemistry, 18: 103-104.
- Pakrashi S.C., Achari B. & Majdumar P.C., 1975. *Studies on Indian medicinal plants. Part XXXII- Constituents of Ananas comosus (Linn.) Merr. leaves*. Indian J. Chem., 13: 755.
- Pignatti S., 1982. *Flora d' Italia*. 2° vol. Edagricole, Bologna.
- Poldini L., 1980. *Catalogo floristico del Friuli-Venezia Giulia e dei territori adiacenti*. Studia Geobot., 1: 314-374.
- Poldini L., 1991. *Atlante delle piante vascolari nel Friuli-Venezia Giulia. Inventario floristico regionale*. Regione Autonoma Friuli-Venezia Giulia, Direzione Regionale Foreste e Parchi, Udine, & Dipartimento di Biologia, Università di Trieste.
- Poldini L., Martini F., Ganis P. & Vidali M., 1991. *Floristic databanks and the phytogeographic analysis of a territory. An example concerning Northeastern Italy*. In: Nimis P.L. & Crovello T.J. (eds.), *Quantitative approaches to Phytogeography*. Kluwer Academic Publishers, Dordrecht-Boston-London.
- Sekula B.C. & Nes W.R., 1980. *The identification of cholesterol and other sterols in Euphorbia pulcherrima*. Phytochemistry, 19: 1509-1512.
- Sharma R.K., 1970. *Biosynthesis of plant sterols and triterpenoids. The incorporation of (3RS)-[2- 14 C], (4R)-4- 14 H] mevalonate into α -spinasterol and β -amyrin in Camelia sinensis*. Phytochemistry, 9: 565-568.
- Smith P.F., 1969. *The role of lipids in membrane transport in Mycoplasma laidlawii*. Lipids, 4: 331-336.
- Waller G.R., Mangiafico S., Foster R.C. & Lawrence R.H. Jr., 1981. *Sterols of Delphinium ajacis; production and metabolic relationship in whole plants and callus tissue*. Pl. Med., 42: 344-355.

Received November 11, 1996

Accepted November 10, 1997

CHEMOSYSTEMATIC VALUE OF ARISTOLOCHIC ACIDS IN THE GENUS *ARISTOLOCHIA* L. IN FRIULI-VENEZIA GIULIA (NE ITALY)

Francesca CATENI¹⁾, Laura COASSINI LOKAR²⁾, and Fabrizio MARTINI²⁾

¹⁾ Dipartimento di Scienze Farmaceutiche, Università di Trieste, P.le Europa 1, I-34127 Trieste – ²⁾ Dipartimento di Biologia, Università di Trieste, Via L. Giorgieri 10, I-34127 Trieste

Keywords: *Aristolochia* L., aristolochic acids, chemotaxonomy, discriminant analysis.

Abstract: The composition of aristolochic acids from different populations of four *Aristolochia* species was analysed by HPLC; quantitative data were submitted to discriminant and statistical analyses. The interspecific variability was confirmed, and significant differences were detected by canonical discriminant functions. The aristolochic acid II and the ratio AAII/AI are significant in the discrimination of the examined species.

Introduction

Aristolochia L. (*Aristolochiaceae*) is a well-known genus since the ancient times for its officinal properties. In Italy, several *Aristolochia* species have been widely used as medicinal plants since Plinius up to the last century (Savi 1805, Targioni-Tozzetti 1824, Cassone 1850). Ten species occur in Italy (Nardi 1984), four of which are present in the Friuli-Venezia Giulia Region: *A. clematitis* L., *A. rotunda* L., *A. pallida* Willd. and *A. lutea* Desf. (Gortani 1906, Poldini 1980, 1991, Martini 1990).

Characters related to the hypogean apparatus and the flowers have been considered as the most discriminant taxonomically (Nardi 1984). Nardi's revision was mainly based both on the frequently disregarded flower morphological characters and on chromosome numbers. Some species, i.e. *A. pallida* and *A. lutea*, are morphologically similar, and their determination on morphological evidence alone is difficult.

A previous study (Cateni *et al.* 1993), reported the chemical composition of various populations of *A. clematitis*, and the distribution of aristolochic acids in different parts of the plants, also with regard to the seasonal stages.

In this paper we examine the aristolochic acid composition of the underground apparatus of the *Aristolochia* species occurring in the Friuli-Venezia Giulia Region, which were not yet phytochemically investigated. The distinct properties of aristolochic

acids among the species, and the evidence they provide on their mutual relationship were also considered.

The four species differ in their ecology. *A. clematitis* is the most autoapophytic, and grows in waste grounds along streams, ditches and sandy places near seashores, edges of cultivated fields, vineyards, hedges, rubble, margins of woods, coppices, on damp and aerated soils, rich in nitrates, up to 400 m. *Aristolochia rotunda* prefers damp grassy places along field margins, near ditches and streams, hedges, woods, shrubs, up to 400 m. *Aristolochia pallida* grows in woods, hedges, grassy places, mainly on plains and hills (up to 600 m). *Aristolochia lutea* is the taxon of the *A. pallida* complex with the widest distribution, and is also the most euritermic, growing mainly in *Ostrya carpinifolia*-woods (but also in beech woods), up to 1000 m. The distribution area of *A. pallida* includes the western part of the Region, while *A. lutea* is distributed in the eastern part; the northern populations of *A. lutea* co-exist with those of *A. pallida* and are often hardly distinguishable by morphological characters (Martini 1990).

The purpose of this study is to ascertain whether the four species can be distinguished on the basis of their aristolochic acids content, and to contribute further to the knowledge of aristolochic acids variation within the genus.

Materials and methods

Sampling concerned four populations of each species of *Aristolochia* in the following localities:

Aristolochia clematidis: 1) Gorizia, loc. Mainizza, 40 m; 2) Sgonico (TS), 255 m; 3) Polcenigo (PN), loc. S. Giovanni, 60 m; 4) Monfalcone (GO), loc. Alberoni, 2 m.

Aristolochia rotunda: 5) Muggia (TS), loc. Noghere, 5 m; 6) Muzzana del Turgnano (UD), loc. Bosco Coda di Manin, 6 m; 7) Latisana, Precenicco (UD), 3 m; 8) Monfalcone (GO), loc. Alberoni, 2 m.

Aristolochia pallida: 9) Polcenigo (PN), loc. Rui Bragam, 70 m; 10) Tramonti di Sotto (PN), loc. Stalle Spinespes, 590 m; 11) Peonis (UD), loc. Rio Sech, 400 m; 12) Jof di Maniago (PN), 450 m.

Aristolochia lutea: 13) M. S. Leonardo (TS), 350 m; 14) Ferneti (TS), 360 m; 15) Ajdovščina (Slovenia), Trnovski Gozd, M. Kucelj, 650 m; 16) Maniago (PN), 400 m.

The plants were collected at full anthesis to examine the quantitative variability in the hypogean apparatus. In each population, five samples of ten wild plants were randomly collected and voucher specimens were deposited in the Herbarium Universitatis Tergestinae (TSB).

The hypogean parts were treated with liquid nitrogen, finely grounded (60 mesh) and weighed (1 g dry weight estimated on another sample). The extraction was carried out with 150 ml of methanol in Soxhlet to exhaustion. The extracts were evaporated in vacuum to small volume. The residue was redissolved in 10 ml methanol. The extracts were analysed by high performance liquid chromatography (HPLC) as described in a previous paper (Cateni *et al.* 1992). HPLC analyses were performed using a Perkin Elmer ODS RP-column filled with bonded octadecylsilane on silica (10 µm), flow rate 1,8 ml/min, detection 310 nm (detector sensitivity 0,01 a.u.f.s.) and methanol/water/acetic acid (80:20:1) as mobile phase. All the determinations were performed at 22° C and at isocratic condition of elution.

The analytical data were submitted to elaboration with statistical methods (Zar 1984, Lagonegro & Feoli 1985a, 1985b).

Results and discussion

The study and the experiment were considered a good test for application of aristolochic acid data to

Tab. 1 - Variation and discriminant analysis in aristolochic acids I and II (% dry weight) in the examined populations of *Aristolochia*.

Species	Populations	Aristolochic acids I (%) II (%)		Ratio AAII/ AAI
<i>A. clematidis</i>	1	0.54	0.58	1.1
	2	0.25	0.24	1.0
	3	0.26	0.28	1.1
	4	0.42	0.54	1.3
	Average	0.37	0.41	1.1
	St. dev.	0.14	0.17	0.13
<i>A. rotunda</i>	5	0.09	0.09	1.0
	6	0.05	0.07	1.4
	7	0.07	0.09	1.2
	8	0.07	0.09	1.2
	Average	0.07	0.08	1.2
	St. dev.	0.02	0.01	0.17
<i>A. pallida</i>	9	0.05	0.16	3.1
	10	0.05	0.17	3.5
	11	0.06	0.19	3.4
	12	0.06	0.18	3.2
	Average	0.05	0.18	3.3
	St. dev.	0.003	0.02	0.2
<i>A. lutea</i>	13	0.04	0.1	2.4
	14	0.05	0.1	2.0
	15	0.04	0.1	2.2
	16	0.08	0.18	2.2
	Average	0.05	0.12	2.2
	St. dev.	0.017	0.04	0.16
Variable	Wilkslambda	F		Significance
AAI	0.1766	118.1		0
AAII	0.2697	68.6		0

taxonomic problems in the genus *Aristolochia* which is currently under revision. All the populations of the four examined species contain in the underground apparatus the same aristolochic acids (AAI and AAII) in different quantities. The quantitative composition in AAI and AAII and their ratio value in the various populations and species are reported in Tab. 1.

The quantitative acid composition of *A. clematidis* L. is clearly distinct from those of the other investigated species, and all its populations are more productive, about two to ten times higher; furthermore, in all the populations (except n. 2) AAII prevails. The quantity of AAII in the population n. 16 (*A. lutea*) is similar to that of *A. pallida*. These samples were derived from an isolated locality inside the distribution area of *A. pallida*. The ratio between the two aristolochic acids in the four species is very different for *A. pallida*, *A. lutea* and the remaining species: for *A. clematidis* the mean value is 1.1, for *A. rotunda* 1.2, for *A. pallida* 3.3 and for *A. lutea* 2.2.

The analysis of variance was used to display

- Chemosystematic value of aristolochic acids -

Tab. 2. - Analysis of variance of aristolochic acid I (left) and II (right) based on % dry weight in the populations of the four species of *Aristolochia* L. (Ac = *Aristolochia clematidis*; Ar = *A. rotunda*; Ap = *A. pallida*; Al = *A. lutea*). ($F_{0.05(3,16)}=3.24$; DF=degree of freedom; F= variance ratio test; Q= statistics for Cohrans's Q test).

AA I

Species	Source of variation	Sum of squares	DF	Mean squares	F	Q
<i>A. clematidis</i>	Treatment	0.293	3	0.098	285.691	
	Error	0.005	16	0		
	Total	0.299	19			
	Ac-Ar				208.015	24.981
	Ac-Ap				192.511	24.032
	Ac-Al				35.083	10.259
	Ar-Ap				0.300	0.949
	Ar-Al				72.244	14.722
	Ap-Al				63.230	13.775
<i>A. rotunda</i>	Treatment	0.004	3	0.001	326.077	
	Error	0.001	16	0		
	Total	0.005	19			
	Ac-Ar				325.167	31.233
	Ac-Ap				75.288	15.023
	Ac-Al				67.132	14.191
	Ar-Ap				87.591	16.210
	Ar-Al				96.805	17.024
	Ap-Al				0.230	0.831
<i>A. pallida</i>	Treatment	0	3	0	18.368	
	Error	0	16	0		
	Total	0	19			
	Ac-Ar				0.462	1.177
	Ac-Ap				8.297	4.989
	Ac-Al				5.678	4.127
	Ar-Ap				12.673	6.166
	Ar-Al				9.377	5.304
	Ap-Al				0.248	0.862
<i>A. lutea</i>	Treatment	0.004	3	0.001	342.908	
	Error	0	16	0		
	Total	0.004	19			
	Ac-Ar				11.837	5.959
	Ac-Ap				1.606	2.195
	Ac-Al				269.599	28.439
	Ar-Ap				4.724	3.746
	Ar-Al				168.453	22.480
	Ap-Al				229.395	26.245

AA II

Species	Source of variation	Sum of squares	DF	Mean squares	F	Q
<i>A. clematidis</i>	Treatment	0.453	3	0.151	487.210	
	Error	0.005	16	0		
	Total	0.458	19			
	Ac-Ar				308.661	30.430
	Ac-Ap				242.963	26.998
	Ac-Al				4.956	3.856
	Ar-Ap				3.926	3.432
	Ar-Al				235.395	26.574
	Ap-Al				178.519	23.142
<i>A. rotunda</i>	Treatment	0.002	3	0.001	23.966	
	Error	0	16	0		
	Total	0.002	19			
	Ac-Ar				20.363	7.816
	Ac-Ap				1.006	1.738
	Ac-Al				0.698	1.447
	Ar-Ap				12.315	6.078
	Ar-Al				13.521	6.369
	Ap-Al				0.028	0.291
<i>A. pallida</i>	Treatment	0	3	0.001	68.335	
	Error	0	16	0		
	Total	0	19			
	Ac-Ar				12.782	6.193
	Ac-Ap				67.065	14.184
	Ac-Al				22.086	8.140
	Ar-Ap				21.290	7.992
	Ar-Al				1.264	1.947
	Ap-Al				12.179	6.045
<i>A. lutea</i>	Treatment	0	3	0.008	920.049	
	Error	0	16	0		
	Total	0	19			
	Ac-Ar				0.043	0.360
	Ac-Ap				0.043	0.360
	Ac-Al				613.250	42.892
	Ar-Ap				0.173	0.721
	Ar-Al				602.990	42.532
	Ap-Al				623.598	42.253

Tab. 3.- T Student's results for regression lines: centroids slopes and intercepts values. Df= 5, $\alpha < 0.05$. (Ac= *Aristolochia clematidis*; Ar= *A. rotunda*; Ap= *A. pallida*; Al= *A. lutea*).

Species	Centroids		Slopes		Intercepts	
	T Student	Prob (%)	T Student	Prob (%)	T Student	Prob (%)
Ac - Ar	6.49	0.17	2.29	8.37	12.29	0.004
Ac- Ap	6.79	0.25	4.92	0.8	13.39	0.004
Ac- Al	7.09	0.21	2.78	4.97	13.36	0.004
Ar - Ap	11.51	0.03	13.66	0.02	9.24	0.02
Ar - Al	2.71	5.38	9.33	0.08	9.47	0.02
Ap - Al	3.28	3.04	6.36	0.31	11.20	0.001




Fig. 1. - Ellipses of equal concentration, regression lines and centroids for the examined *Aristolochia* species at the 5% level of confidence. The slopes and intercepts of the regression lines are significantly different ($p < 0.05$). 1: *A. clematitis*; 2: *A. rotunda*; 3: *A. pallida*; 4: *A. lutea*. The quantities are expressed as % dry weight.

the source of noticed variation among the four species. Statistically significant variation occurred for AAI and AAII. The most significant is AAI ($F = 118.1$) (Tab. 1). The results of the analysis of variance among populations are reported in Table 2. Because the critical value F is 3.14 at the 0.05 level of significance, all the values, which are smaller of 3.14, are not rejected. Therefore, *A. clematitis* populations are significantly distinct from those of the other species, in terms of both AAI and AAII. Statistically significant variations in the quantities of AAI occur among all the other populations of the various species. The populations of *A. lutea* significantly differ from those of *A. pallida*, although the two species are hardly distinguishable morphologically.

The four species can be discriminated by regression lines and centroid values of the aristolochic acids (Fig. 1). The values of centroid, slope and intercept are reported in Tab. 3. Most discriminant are the intercept and centroid values. The greatest disjunction is between *A. clematitis* and the other species.

Conclusions

The results demonstrate that aristolochic acids may be used for chemotaxonomical purposes in *Aristolochia*. The co-occurrence of structurally closely related compounds points to the chemical uniformity of the investigated species, differences being only quantitative. Aristolochic acid I and the ratio AAII/AAI are the more significant taxonomically: chemical and morphological data are highly congruent.

Several questions regarding systematic relationships can be addressed using chemical data. The four *Aristolochia* species may be distinguishable from one another, based on aristolochic acid II data. Quantitative differences were compared: (a) among populations of different species, and (b) by combining the populations of each species to increase the sample size. The two closely related species, *A. pallida* and *A. lutea*, can be also distinguished on a chemical basis. It is confirmed that, chemically, *A. clematitis* is more distant one from the other species.

These results have implications both for basic research on plant populations and for applied studies on plant secondary products as defensive or attractive agents. The different concentrations of the same compounds in the populations and in the species suggest that they may have a specific biological role, and could be related both to their possible function as allelopathic agents (Nishida & Fukami 1989) and to their ecological role, since the plant toxins can be sequestered from the diet and stored by insects for defence. Aristolochic acids are actually used for this purpose by the butterfly *Battus archidamas* Boisd. (larval stage, Harborne 1987, 1988). It is interesting to note that in Friuli-Venezia Giulia, another butterfly, *Zerynthia polyxena* Schiff. (= *Z. hysipyle* Schulz., *Papilionidae*) also feeds on *Aristolochia* (Morandini, *ex verbis*).

Riassunto

VALORE CHEMOSISTEMATICO DEGLI ACIDI ARISTOLOCHICI NEL GEN. *ARISTOLOCHIA* L. NEL FRIULI-VENEZIA GIULIA. La composizione in acidi aristolochici di differenti popolazioni appartenenti alle quattro specie di *Aristolochia* L. presenti nella regione Friuli - Venezia Giulia è stata analizzata tramite HPLC, analisi discriminanti e statistiche. La variabilità interspecifica è stata confermata e differenze significative sono state evidenziate tramite le funzioni canoniche discriminanti. Gli acidi aristolochici I e II e in particolare il rapporto AAII/AAI hanno valore di marcatori chemotassonomici data la congruenza fra dati chimici e morfologici.

Acknowledgments

The Authors would like to thank the Director of the Museo Friulano di Storia Naturale (Udine) Dr. C. Morandini, for information on the butterfly *Zerynthia polyxena*. This project was carried out thanks to the financial support of the Italian Ministry of Public Instruction (M.P.I. 40%).

References

- Cassone F., 1850. *Flora medico farmaceutica*. 5. Torino.
- Cateni F., Mamolo M.G. & Coassini Lokar L., 1992. *Rapid HPLC analysis for quantitative determination of aristolochic acids*. *Il Farmaco*, 47 (10): 1335-1342.
- Cateni F., Coassini Lokar L., Martini F. & Vrech E., 1993. *Aristolochic acids variation and distribution in some Aristolochia clematitis L. populations*. *Studia Geobot.*, 13: 299-312.
- Gortani L. & M., 1906. *Flora friulana con speciale riguardo alla Carnia*. 2. Udine.
- Harborne J.B., 1987. *Chemical signals in the ecosystem*. In: Dodge J.E. (ed.), *New Perspectives in Plant Science*. London, New York.
- Harborne J.B., 1988. *Introduction to Ecological Biochemistry*. New York.
- Lagonegro M. & Feoli E., 1985a. *The use of ellipses of equal concentration to analyse ordination vegetation patterns*. *Studia Geobot.*, 5: 143-165.
- Lagonegro M. & Feoli E., 1985b. *Analisi multivariata di dati. Manuale d'uso di programmi BASIC per personal computer*. Libreria Goliardica, Trieste.
- Martini F., 1990. *Il gruppo Aristolochia pallida nell'Italia nordorientale*. *Giorn. Bot. Ital.*, 124: 731-743.
- Nardi E., 1984. *The genus "Aristolochia" L. (Aristolochiaceae) in Italy*. *Webbia*, 38: 221-300.
- Nishida R. & Fukami H., 1989. *Ecological adaptation of an Aristolochiaceae-feeding swallowtail butterfly, Atrophaneura alcinous to aristolochic acid*. *J Chem. Ecol.*, 15: 2549-2563.
- Poldini L., 1980. *Catalogo floristico del Friuli-Venezia Giulia e dei territori adiacenti*. *Studia Geobot.*, 1: 313-474.
- Poldini L., 1991. *Atlante corologico delle piante vascolari nel Friuli-Venezia Giulia*. Regione Autonoma Friuli-Venezia Giulia, Direzione Regionale Foreste e Parchi, Udine, & Dipartimento di Biologia, Università di Trieste..
- Savi G., 1805. *Materia medica vegetabile toscana*. Firenze.
- Targioni-Tozzetti A., 1824. *Scelta di piante officinali più necessarie a conoscersi*. Firenze.
- Zar J.H., 1984. *Biostatistical analysis*, 2nd ed. Prentice- Hall Inc., New Jersey.

Received November 11, 1996

Accepted November 10, 1997

PHYTOSOCIOLOGICAL OBSERVATIONS ON THE BRYOLOGICAL VEGETATION OF MT. PALON (Mt. BONDONE GROUP, TRENTINO-ALTO ADIGE)

Michele ALEFFI¹, Maria PRIVITERA² and Marta PUGLISI²

¹ Dipartimento di Botanica e di Ecologia, Università di Camerino, Via Pontoni 5, I-62032 Camerino (MC)

² Dipartimento di Botanica, Università di Catania, Via A. Longo 19, I-95125 Catania.

Keywords: Alps, Bryophyte Communities, Mt. Palon, Trentino-Alto-Adige.

Abstract: Results of a study on bryophyte sociology in some localities of Mt. Bondone (eastern Italian Alps) are presented. Bryophyte associations observed in beech forests and *Pinus-mugo* scrubs are: *Orthotricho anomaligrimmietum pulvinatae*, *Pseudoleskeetum catenulatae*, *Ctenidietum mollusci*, *Solorino-Distichietum capillacei*, *Encalypto streptocarpace-Fissidentetum cristati* and *Pterigynandretum filiformis*, belonging to the *Tortulo-Homalothecietea sericei* class, and *Pleurozietum schreberi* of the *Hylocomietea splendentis* class.

Introduction

Although in the past Trentino-Alto Adige has been the focus of numerous bryological explorations and studies (Gerdol & Tomaselli 1991, Gerdol *et al.* 1991, Gerdol *et al.* 1994), very little is known about the bryovegetation of this region. Only fragmentary information is available on the following associations: *Barbuletea unguiculatae* Mohan 1978 (Cortini Pedrotti & Aleffi 1990), *Racomitrietea heterostici* Neumayr 1971 (Giacomini 1950), *Tortulo-Homalothecietea sericei* Hertel ex Mohan 1978 (Philippi 1983, Cortini Pedrotti 1994) and *Frullanio dilatatae-Leucodontetea sciuroidis* Mohan 1978 (Philippi 1983).

In particular, Monte Bondone was very little studied, information on its bryoflora being limited to publications by Venturi (1899) and Dalla Torre & Sarnthein (1904), while data on bryovegetation are given by only a single study on the Palinegra Bog near the "Le Viotte" Botanical Garden (Gerdol & Piccoli 1979).

In the summer of 1995, the Working Group for Bryology of the Italian Botanical Society conducted a floristic excursion in this area. The product of the excursion was a list of 142 species (29 liverworts and 113 mosses), 3 of which were new for Trentino-Alto Adige (Aleffi & Cortini Pedrotti 1996). The present work presents some aspects of the bryovegetation, sampled on that occasion.

Study area

The Monte Bondone Group extends along the orographic right of the Adige valley and reaches its highest point with the peak of Monte Cornetto (2179 m), that, together with the Doss d'Abramo and the Cima Verde, delimits to the south the wide Viotte valley, which is blanketed by a deep cover of morenic material. To the east of the Viotte is Monte Palon (2090 m) and to the west La Rosta (1832 m). All of these mountains are composed of sedimentary rocks, in particular Mesozoic limestones.

The climate of Mt. Bondone is of a pre-Alpine type; average annual precipitation, measured at the Viotte station at 1530 m, reaches 1167 mm and mean annual temperature is 5.9 °C. The rain distribution is equinoctial, with spring and autumn maxima.

The vegetation of the summit belongs to the montane and subalpine belts (Pedrotti 1981, Pedrotti *et al.* 1994). In the montane belt there is a lower montane horizon up to about 1000 m with beech forests of *Carici albae-Fagetum* Rübel 1930 ex Moor 1952 em. Lohm. 1953, and an upper montane horizon between 1000 and 1750 m, with beech forests of *Cardamino pentaphylli-Fagetum* Mayer 1974. The subalpine belt is characterized by *Pinus-mugo* scrubs prevalently of the *Sorbo chamaemespili-Pinetum*, an association which develops on calcareous substrata, but with humus-rich, acidic soils (Minghetti 1996); *Erico carnea-*

Pinetum prostratae Zöttl 1951 and *Rhododendro hirsuti-Pinetum prostratae* Zöttl 1951 grow on calcareous substrata with neutro-basic soils.

Material and methods

The phytosociological analysis has been carried out following the method of the Zürich-Montpellier School. Relevés were carried out on the slopes of Mt. Palon, on the east side, between ca. 1300 and 1500 m, corresponding to the *Cardamino pentaphylli-Fagetum*, and on the north-west side between ca. 1650 and 1800 m (La Cuna) in the *Sorbo chamaespili-Pinetum mugo*. They are from dry and sunny to moist and shaded calcareous rocks, from soil in *Pinus mugo* scrubs, and from trunks of beech trees.

The syntaxonomic arrangement follows Marstaller (1993); nomenclature follows Corley *et al.* (1981) and Corley & Crundwell (1991) for mosses, and Grolle (1983) for hepatics; author's names are according to Crosby (1992) and Grolle (1983).

Syntaxonomical scheme

TORTULO - HOMALOTHECIETEA SERICEI
Hertel ex Mohan 1978

GRIMMIETALIA ANODONTIS Smarda 1947

GRIMMION TERGESTINAE Smarda 1947

Orthotricho anomali-Grimmietum pulvinatae Stodiek 1937

Pseudoleskeelletum catenulatae Jezek & Vondráček 1962

CTENIDIETALIA MOLLUSCI Hadác & Smarda in Klika & Hadác 1944

CTENIDION MOLLUSCI Stefureac 1941

Ctenidietum mollusci Stodiek 1937

Solorino-Distichietum capillacei Reimers 1940

Encalypto streptocarpace-Fissidentetum cristati Neumayr 1971

NECKERETALIA COMPLANATAE Jezek & Vondráček 1962

NECKERION COMPLANATAE Smarda & Hadác in Klika & Hadác 1944

Pterigynandretum filiformis Hiliter 1925

HYLOCOMIETEA SPLENDENTIS Gillet ex Marstaller 1993

HYLOCOMIETALIA SPLENDENTIS Gillet ex Marstaller 1993

PLEUROZION SCHREBERI v. Krusentjerna 1945

Pleurozietum schreberi Wisniewski 1930

Results and discussion

Previous studies revealed the presence of some lithophilous and terricolous communities of the *Tortulo-Homalothecietea sericei* and *Hylocomietea splendentis* classes. Within the *Tortulo-Homalothecietea sericei*, widely distributed in central Europe, with many communities already known from Italy, associations of the *Grimmietalia anodontis* and *Ctenidietaalia mollusci* were identified. In addition, an epiphytic aspect of the bryological vegetation was observed, and referred to the *Neckeretalia complanatae* order, belonging to the same class.

Within the *Hylocomietea splendentis* class, the *Pleurozietum schreberi* association (*Pleurozion schreberi* alliance), was identified. This class includes montane and alpine terri-saxicolous meso- or hygrophytic communities at high altitude; however, some of them are found in different habitats (waterfalls, rocks along roadsides, etc.). The *Hylocomietea splendentis* class, with the single order *Hylocomietalia splendentis*, was recently recognized by Marstaller (1993). Well known in central and northern Europe, this class is mentioned for the first time from Italy in this article.

Orthotricho anomali-Grimmietum pulvinatae Stodiek 1937

A xero- and photophytic, pioneer community found on exposed calcareous rocks in beech woods at ca. 1500 m (Tab. 1, rel. group A). It is dominated by *Orthotrichum anomalum*, an association guide species, and *Schistidium apocarpum*, an alliance guide species; some class character species join them, such as *Homalothecium sericeum*, *Ctenidium molluscum* and *Rhynchostegium murale*. Syntaxonomically, it comes under the *Grimmion tergestinae* alliance of the *Grimmietalia anodontis* order, a syntaxon of the mostly xero- and photophytic *Tortulo-Homalothecietea sericei* class.

Widely distributed in Central Europe (Hübschmann 1986), it has also been reported for the Iberian peninsula (Guerra & Varo 1981, Gil &

Guerra 1985, Fuertes Lasala & Martinez Conde 1988, Moya *et al.* 1994), northern and central Italy (Giacomini 1950, Hébrard 1978) and Sicily (Lo Giudice & Privitera 1987, Lo Giudice *et al.* 1992).

***Pseudoleskeelletum catenulatae* Jezek & Vondráček 1962**

A montane, saxicolous, xero-mesophytic, photo-sciophytic association, found in *Cardamino pentaphylli-Fagetum* forests, on calcareous rocks with no or little inclination (Tab. 1, rel. group B). It is found in less xeric and sunny habitats than those colonized by *Orthotricho anomali-Grimmietum pulvinatae*. This community is physiologically heterogeneous, due to the presence of both pleurocarpous and acrocarpous mosses, with both a turf and cushion habit. Character species is *Pseudoleskeella catenulata*, a xerophilous and mesophilous species with a boreal-montane/dealpine distribution (Düll 1984-1985). Species of higher syntaxa are *Schistidium apocarpum* and *Orthotrichum anomalum*, *Tortella tortuosa* and *Homalothecium sericeum*. This community is included in the *Grimmion tergestinae* alliance of the *Grimmietalia anodontis*.

The association has often been reported for central Europe (Jezek & Vondráček 1962, Nörr 1970, Neumayr 1971, Hertel 1974, Marstaller 1980, 1992a, 1995). For southern Europe it is known from southeastern France (Hébrard 1971) and from the Iberian peninsula (Guerra 1985, Fuertes Lasala & Martinez Conde 1988). The association is new to Italy.

***Solorino-Distichietum capillacei* Reimers 1940**

This association (Tab. 1, rel. group C) was found on the eastern and northwestern slopes of M. Palon, on fairly humid and shaded rocks; it prefers northern exposure. It substitutes *Pseudoleskeelletum catenulatae* in more mesic and shaded conditions.

Distichium capillaceum, a faithful species, and *D. inclinatum*, a boreal-montane species, are characteristic of the association; they are accompanied by *Ctenidium molluscum*, *Campylium halleri*, which are alliance character species, and by several order species such as *Tortella tortuosa*, *Rhynchostegium murale*, *Encalypta streptocarpa*, *Schistidium apocarpum*, *Mnium thomsonii* and *Scapania aequiloba*. Floristically and ecologically,

the association should be placed under the *Ctenidion mollusci* alliance of the *Ctenidietalia mollusci* order, a syntaxon of the *Tortulo-Homalothecietea sericeae* class, with scio- and mesophytic characters.

The distribution prevalently extends along the mountains of central Europe (von Hübschmann 1986, Marstaller 1987, 1991, 1992b); from southern Europe, this association was recently reported for some mountains of Sicily (Lo Giudice *et al.* 1992).

***Ctenidietum mollusci* Stodiek 1937**

This community (Tab. 1, rel. group D) colonizes fairly shaded and humid calcareous rocks covered by a soil layer. Quite common in the survey area, it behaves as an exocomophytic, clearly mesophilous and sciophilous association. It is characterized by *Ctenidium molluscum*, a circumboreal guide species, joined by many alliance and order character species. This association belongs in the *Ctenidion mollusci* alliance of the *Ctenidietalia mollusci* order. It is widespread in Central Europe (Hübschmann 1986), and has also been reported from former Yugoslavia (Hébrard 1975), the Iberian peninsula (Guerra & Varo 1981, Gil & Guerra 1985, Fuertes Lasala & Martinez Conde 1988) and Sicily (Lo Giudice *et al.* 1992, Privitera & Puglisi 1996).

***Encalypto streptocarpha-Fissidentetum cristati* Neumayr 1971**

This chasmocomophytic, meso- and sciophytic community (Tab. 1, rel. group E) is frequently found both in beech woods and in *Pinus mugo* scrubs, where it colonizes fissures and concavities of the rocks, mostly covering small surfaces. *Fissidens dubius* (= *F. cristatus*) is an association character species. *Ctenidium molluscum*, an alliance character species, accompanies it, as some order and class character species, among which *Encalypta streptocarpa* and *Tortella tortuosa* are striking for their constancy and abundance. The association is included in the *Ctenidion mollusci* alliance of the *Ctenidietalia mollusci* order. It is known from Germany, Czechoslovakia, France, Spain (Vadani 1983, Ros & Guerra 1987) and, recently, from Sicily (Privitera & Puglisi 1996).

***Pterigynandretum filiformis* Hilitzer 1925**

This association (Tab. 1, rel. group F) was found

54

Studia Geobot. 16 (1998)

sporadically on the trunks of beech trees. *Pseudoleskeella nervosa* and *Pterigynandrum filiforme*, respectively association character and differential species, dominate the community. *Plagiomnium cuspidatum*, is a *Neckerion complanatae* character species, joined by several corticolous species of the *Frullanio dilatatae-Leucodontetea sciurioidis* class, such as *Frullania dilatata*, *Radula complanata*, *Orthotrichum pumilum*, *O. striatum*. Synsystematically, Marstaller (1984) placed the association in the *Neckerion complanatae* of the *Neckeretalia complanatae*, a *syntaxon* in which he joined cortico-saxicolous and epiphytic associations. Even though our relevés would belong more to the *Frullanio dilatatae-Leucodontetea sciurioidis* class, we feel that for the moment Marstaller's suggested assignment should be followed. The community is distributed in many localities of central Europe, with maximum diffusion in the eastern and north-eastern parts; it extends to the Carpatians and the southern Alps, and, in northern Europe, as far north as southern Scandinavia (Marstaller 1992c). In Italy there are repots from Lombardy, Veneto and Trentino (Philippi 1983).

Pleurozietum schreberi Wisniewski 1930

This montane association (Tab. 1, rel. group G) was found on rocks covered by a thick stratum of humus-rich soil in fairly humid and shaded sites in the *Pinus mugo* scrubs, where it is quite common. It is composed of only a few mosses, almost all of which are of considerable size, its low biodiversity being counterbalanced by the high biomass, forming a more or less continuous blanket on wide surfaces. It could represent the most evolved stage among bryophyte communities of the understory of beechwoods and conifer woods of mountains with an alpine climate.

Hylocomium splendens, a sub-boreal moss, is considered as the association guide species; it is accompanied by the alliance character species *Pleurozium schreberi* and *Dicranum scoparium*, and by the class character species *Rhy-tidiadelphus triquetrus*. The few companion species are *Ctenidium molluscum* and *Tortella tor-tuosa*. The community belongs to the *Pleurozion schreberi* alliance of the *Hylocomietalia splen-dentis* order, in the *Hylocomietea splendentis* class. Described by Wisniewski (1930) for Poland, this association is

also known from Thuringia (Marstaller 1994) and here it is reported for the first time for Italy. We also found it on M. Baldo above Rubiana (Verona) at about 850 m, within beech woods. In addition, some vegetational aspects identified by Giacomini (1939) in the Lombard Alps and cited generically as "*Hylocomietum*", could be attributed to the *Pleurozietum schreberi*.

Acknowledgments

This study has been supported by funds from MURST 60%.

References

- Aleffi M. & Cortini Pedrotti C., 1996. *Contributo alla Flora Briologica del Gruppo del Monte Bondone (Trentino-Alto Adige)*. Inform. Bot. Ital., 28: 161-170.
- Corley M.F.V. & Crundwell A.C., 1991. *Additions and amendments to the mosses of Europe and the Azores*. J. Bryol., 16: 337-356.
- Corley M.F.V., Crundwell A.C., Düll R., Hill M.O., Smith A.J.E., 1981. *Mosses of Europe and the Azores; an annotated list of species, with synonyms from the recent literature*. J. Bryol., 11: 609-689.
- Cortini Pedrotti C., 1994. *Briofite delle Marocche di Dro*. In: Pedrotti F. (ed.), *Guida all'escursione della Società Italiana di Fitosociologia in Trentino (1-5 luglio 1994)*. Dipartimento di Botanica e di Ecologia, Università degli Studi di Camerino, 66-67.
- Cortini Pedrotti C. & Aleffi M., 1990. *Associazioni di briofite e di alghe dei laghi Idro e Terlago (Italia settentrionale)*. Doc. Phytosoc., 12: 265-272.
- Crosby M.R., 1992. *Bryophytes*. In: Brummit R.K. & Powell C.E. (eds.), *Authors of plant names*. Royal Botanical Gardens, Kew.
- Dalla Torre K.W. & Samthain L.G., 1904. *Die Moose von Tirol, Vorarlberg und Liechtenstein*. 5. Wagner, Innsbruck, LIV + 671 pp.
- Düll R., 1984. *Distribution of European and Macaronesian mosses (Bryophytina). Part I*. Bryol. Beitr., 4: 1-109.
- Düll R., 1985. *Distribution of European and Macaronesian mosses (Bryophytina). Part II*. Bryol. Beitr., 5: 110-232.
- Fuertes Lasala E. & Martinez-Conde E., 1988. *Vegetación briofítica del macizo oriental de los picos de Europa en Cantabria (España). I. Comunidades saxícolas y acuáticas*. Cryptogamie, Bryol. Lichénol., 9: 109-127.
- Gerdol R. & Piccoli F., 1979. *La vegetazione della Torbiera Palinestra (Monte Bondone - Trento)*. Arch. Bot. Biogeogr. Ital., 55: 25-33.
- Gerdol R. & Tomaselli M., 1991. *Tipologia e caratterizzazione ecologica della vegetazione di alcune torbiere a sfagni delle Dolomiti alto-atesine*. Ann. Lab. Prov. Aut. Bolzano, 6: 153-173.
- Gerdol R., Tomaselli M. & Boiti I., 1991. *Bryophyte flora and vegetation of the glaciokarst plains in the Dolomites (S. Alps, Italy)*. J. Bryol., 16: 413-427.
- Gerdol R., Tomaselli M. & Bragazza L., 1994. *A Floristic-Ecologic Classification of Five Mire Sites in the Montane-*

- Subalpine Belt of South Tyrol (S Alps, Italy)*. *Phyton*, 34: 35-56.
- Giacomini V., 1939. *Studi briogeografici. Associazioni di briofite in Alta Valcamonica e in Valfurva (Alpi Retiche di Lombardia)*. *Atti Ist. Bot. "Giovanni Briosi"*, 12: 1-139.
- Giacomini V., 1950. *Ricerche sulla flora briologica xeroterminica delle Alpi italiane*. *Vegetatio*, 3: 1-123.
- Gil J.A. & Guerra J., 1985. *Estudio briosociológico de las sierras de la Demanda y Urbión (España)*. *Cryptogamie, Bryol. Lichénol.*, 6: 219-258.
- Grolle R., 1983. *Hepatics of Europe including the Azores: an annotated list of species, with synonyms from the recent literature*. *J. Bryol.*, 12: 403-459.
- Guerra J., 1985. *Nuevos datos sobre la clase Tortulo-Homalothecieta sericea en las Sierras Béticas (sur de España)*. *Lazaroa*, 8: 323-331.
- Guerra J. & Varo J., 1981. *Contribución al conocimiento de la vegetación briofítica terrícola e basófila del sur de la Península Ibérica*. *An. Jard. Bot. Madrid*, 37: 693-701.
- Hébrard J.P., 1971. *Contribution à l'étude des Bryoassociations rupicoles de l'étage alpin dans le Sud-Est de la France*. *Ext. Ann. L'Univ. Provence-Sciences*, 46: 117-149.
- Hébrard J.P., 1975. *Remarques sur la flore et la végétation Bryologique de Bosnie-Herzégovine (Yougoslavie)*. *Rev. Bryol. Lichénol.*, 41: 397-448.
- Hébrard J.P., 1978. *Aperçu sur la végétation muscinales des montagnes calcaires dans la province italienne des Abruzzes*. *Rev. Bryol. Lichénol.*, 44: 7-45.
- Hertel E., 1974. *Epilithische Moose und Moosgesellschaften im nordöstlichen Bayern*. *Ber. Naturwiss. Ges. Bayreuth*, 1: 1-498.
- Hübschmann von A., 1986. *Prodromus der Moosgesellschaften Zentraleuropas*. *Bryophytorum Bibliotheca*, 32: 1-413.
- Jezek V. & Vondráček M., 1962. *Společensva mechorostu doliny Siedmich Pramenov v Belanských Tatrách*. *Biol. Práce Slor. Akad. Vied.*, 8: 1-48.
- Lo Giudice R. & Privitera M., 1987. *Sulla briovegetazione del Vallone Murapane (Piazza Armerina)*. *Boll. Acc. Gioenia Sci. Nat.*, 20: 149-163.
- Lo Giudice R., Privitera M. & Puglisi M., 1992. *Sulla classe Tortulo-Homalothecieta sericea Hertel 1974 in Sicilia*. *Arch. Bot. Ital.*, 68: 153-172.
- Marstaller R., 1980. *Die Moosgesellschaften des Verbandes Schistidion apocarpi Jezek und Vondracek*. *Feddes Repert.*, 91: 337-361.
- Marstaller R., 1984. *Die moosgesellschaften des Naturschutzgebietes "Steinklöße" bei Nebra, Bezirk Halle*. *Arch. Naturschutz u. Landsch.-Forsch.*, 24: 1-15.
- Marstaller R., 1987. *Die Moosvegetation des Binsenberges bei Krölpa, Kreis Pöneck*. 27 *Beitrag zur Moosvegetation Thüringens*. *Gleditschia*, 15: 139-156.
- Marstaller R., 1991. *Zur Kenntnis der Moosvegetation des Clydenfelsens bei Ölsen, Kreis Pöneck*. 55. *Beitrag zur Moosvegetation Thüringens*. *Gleditschia*, 19: 269-283.
- Marstaller R., 1992a. *Photophytische Moosgesellschaften im Zechsteingebiet zwischen Waldfisch und Schweina, Kreis Bad Salzungen*. *Gleditschia*, 20: 107-122.
- Marstaller R., 1992b. *Bryosozologische Untersuchungen im Zittauer Gebirge*. *Abh. Ber. Naturkundemus. Görlitz*, 66: 1-41.
- Marstaller R., 1992c. *Die Moosgesellschaften des Verbandes Neckerion complanatae Sm. et Had. in Kl. et Had. 1944*. 54. *Beitrag zur Moosvegetation Thüringens*. *Herzogia*, 9: 257-318.
- Marstaller R., 1993. *Synsystematische Übersicht über die Moosgesellschaften Zentraleuropas*. *Herzogia*, 9: 513-541.
- Marstaller R., 1994. *Die Moosgesellschaften des Naturschutzgebietes Kleiner Gleichberg bei Römheld, Grabfeld*. *Ber. Bayer. Bot. Ges.*, 64: 87-113.
- Marstaller R., 1995. *Die Moosgesellschaften des Naturschutzgebietes Bohlen bei Saalfeld*. 57. *Beitrag zur Moosvegetation Thüringens*. *Gleditschia*, 23: 181-199.
- Minghetti P., 1996. *Analisi fitosociologica delle pinete a Pinus mugo Turra del Trentino (Italia)*. *Doc. Phytosoc.*, 16: 461-503.
- Moya J.J., Ros R.M. & Guerra J., 1994. *Bryophyte flora and vegetation of the Sierra del Maigmo (Alicante, Spain)*. *Cryptogamie, Bryol. Lichénol.*, 15: 43-65.
- Neumayr L., 1971. *Moosgesellschaften der südöstlichen Frankenalb und des Vorderen Bayrischen Waldes*. *Hoppea*, 29: 1-364.
- Nörr M., 1970. *Die Moosvegetation des Rübeländer Kalkgebietes*. *Hercynia*, 7: 13-52.
- Pedrotti F., 1981. *Carta della vegetazione del foglio Trento*. C.N.R., Collana programma finalizzato "Promozione qualità ambiente", Roma, AQ/1/17: 1-38.
- Pedrotti F., Gafta D. & Minghetti P., 1994. *Il Monte Bondone*. In: Pedrotti F. (ed.), *Guida all'escursione della Società Italiana di Fitosociologia in Trentino (1 - 5 luglio 1994)*. Dipartimento di Botanica e di Ecologia, Università degli Studi di Camerino, 5-19.
- Philippi G., 1983. *Epiphytische Moosvegetation des Gardasee-Gebietes*. *Andrias*, 2: 23-52.
- Privitera M. & Puglisi M., 1996. *La vegetazione briofítica dell'Etna (Sicilia, Italia)*. *Braun-Blanquetia*, 19: 1-59.
- Ros R. & Guerra J., 1987. *Vegetación briofítica terrícola de la Región de Murcia (sureste de España)*. *Phytocoenologia*, 15: 505-567.
- Vadam J.C., 1983. *L' Encalypto streptocarpe-Fissidentetum cristati Neumayr 1971 dans les environs de Montbéliard (Doubs)*. *Cryptogamie, Bryol. Lichénol.*, 4: 145-153.
- Venturi G., 1899. *Le Muscinee del Trentino*. G. Zippel, Trento.
- Wisniewski T., 1930. *Zespoły mszaków epifitowych Polski ze szczególnem uwzględnieniem puszczy Białowieskiej. (Les associations des Muscinées épiphytes de la Pologne)*. *Bull. Int. Acad. Pol. Sc. Lettr. Sér. B.*, I (1929): 293-343.

Received December 5, 1996

Accepted December 6, 1997

CATALOGUE OF LICHENIZED AND LICHENICOLOUS FUNGI OF CYPRUS

Birgit LITTERSKI¹⁾ and Helmut MAYRHOFER²⁾

¹⁾ Botanisches Institut, Ernst-Moritz-Arndt-Universität Greifswald, Grimmer Str. 88, D-17489 Greifswald, Germany. ²⁾ Institut für Botanik, Karl-Franzens-Universität Graz, Holteigasse 6, A-8010 Graz, Austria

Keywords: altitudinal zones, biodiversity, checklist, Cyprus, flora, lichenicolous fungi, lichenized fungi, lichens.

Abstract: This annotated checklist of lichens and lichenicolous fungi of Cyprus is based mainly on literature records. The currently known lichen flora is composed of 365 taxa (352 species of lichenized fungi, 2 subspecies, 3 varieties and 8 lichenicolous fungi). The Troodos Region, with about 230 taxa, is the part of the island with the highest number of known species. For each lichen species references, regions, substratum and altitudinal range are reported.

Introduction

The annotated catalogue of lichens and lichenicolous fungi of Cyprus provided here is based mainly on literature records, and is a first-approximation checklist compiled within the project of the OPTIMA Commission of Lichens for a general biodiversity inventory of the lichens of the Mediterranean area (Nimis 1996). The first lichen records from Cyprus were published by Flotow (1843), Koerber (1862) and Krempelhuber (1865). The records of Koerber (1862) and Krempelhuber (1865) were based on material collected by F. Unger. Recently, the lichen flora of Cyprus, especially in the Greek part, has received more attention (Aptroot 1996, Litterski 1997, Litterski & Mayrhofer 1998). Some regions in the Turkish part were investigated by Güvenc & Öztürk (1994, 1995).

The island of Cyprus is 9251 km² in size, and is situated in the south-eastern part of the Mediterranean Sea. The vegetation is of eu-Mediterranean type.

The currently known lichen flora comprises 365 taxa (352 species of lichenized fungi, 2 subspecies and 3 varieties of lichenized fungi, and 8 lichenicolous fungi). The Troodos Region, with about 230 taxa, and the Akamas-Pafos Region, with about 110 taxa, are the two parts of the island with the highest number of known species. Of the known species, 8% have *Trentepohliaceae* as phycobiont and occur mostly at lower altitudes, 84% have other green algae, and the remaining 8%

have cyanobionts. The 352 species of lichenized fungi and 8 species of lichenicolous fungi belong to different orders, families and genera. Most of the species belong to the large order *Lecanorales* (222 species). Within the *Lecanorales* the *Physciaceae* (47 species), *Lecanoraceae* (35 species) and *Parmeliaceae* (36 species) are the families with the highest number of known species. Other important orders are the *Teloschistales* (43 species) and *Arthoniales* (22 species).

Methods and arrangement of the checklist

In the following checklist, species are listed alphabetically. For each lichen species, references, regions, substratum and altitudinal range are provided, so far as is possible. The references are listed in temporal order.

The following regions (Fig. 1) are reported in brackets:

- I Akamas-Pafos Region
- II Mamonia Region
- III Troodos Region
- IV Circum Troodos-Mesaoria Region
- V Pentadictolos Region

This delimitation corresponds with geological features (Geological Survey Department 1995). When it is not known where the locality mentioned in the literature is situated, the abbreviation ? is used.

The altitudinal zones are delimited as follows:

- 1996 (II), Litterski 1997 (I, III), Litterski & Mayrhofer 1998 (I, IV); pl/co, mo; ca, ro, si.
- Caloplaca marmorata* (Bagl.) Jatta - Litterski 1997 (III), Litterski & Mayrhofer 1998 (I); pl/co, salp; ca, si.
- Caloplaca navasiana* Nav.-Ros. & Cl. Roux - Navarro-Rosinés & Roux 1993 (IV), Navarro-Rosinés & Roux 1994 (fig. 1), Navarro-Rosinés & Roux 1995 (IV); pl/co; ca.
- Caloplaca saxicola* (Hoffm.) Nordin - Flotow 1843 (*Lecanora murorum* y. *callopisma*), Güvenc & Öztürk 1995 (V), Litterski 1997 (I, II); pl/co; ca, li.
- Caloplaca subpallida* H. Magn. - Aptroot 1996 (II, III); [pl/co, mo]; ro.
- Caloplaca tavaresiana* Nav.-Ros. & Cl. Roux - Navarro-Rosinés & Roux 1993 (IV), Navarro-Rosinés & Roux 1994 (fig. 1); pl/co; ca.
- Caloplaca teicholyta* (Ach.) J. Steiner - Aptroot 1996 (II), Litterski 1997 (I); pl/co; ro.
- Caloplaca variabilis* (Pers.) Müll. Arg. - Litterski & Mayrhofer 1998 (I); pl/co; ca.
- Caloplaca veneris* Cl. Roux & Nav.-Ros. - Roux & Egea 1992 (IV), Roux & Navarro-Rosinés 1992 (IV), Navarro-Rosinés & Roux 1993 (IV), Navarro-Rosinés & Roux 1994 (fig. 1); pl/co; ca.
- Candelariella aurella* (Hoffm.) Zahlbr. - Güvenc & Öztürk 1995 (V), Litterski 1997 (III), Litterski & Mayrhofer 1998 (I); pl/co, submo, salp; ca, Jn, Ps, si.
- Candelariella vitellina* (Hoffm.) Müll. Arg. - Aptroot 1996 (III), Litterski 1997 (III), Litterski & Mayrhofer 1998 (III); mo, salp; ro, si.
- Candelariella xanthostigma* (Ach.) Lettau - Aptroot 1996 (III); [mo]; Pi.
- **Carbonea vitellinaria* (Nyl.) Hertel - Litterski & Mayrhofer 1998 (III); salp; si, on *Candelariella vitellina*.
- Catapyrenium daedaleum* (Kremp.) Stein - Aptroot 1996 (II, III), Litterski 1997 (I, III); pl/co, submo; so.
- Catapyrenium psoromoides* (Borrer) R. Sant. - Litterski & Mayrhofer 1998 (III); submo; Qu.
- Catillaria chalybeia* (Borrer) A. Massal. - Aptroot 1996 (III), Litterski & Mayrhofer 1998 (I); pl/co, [mo]; sa, ro.
- Catillaria lenticularis* (Ach.) Th. Fr. - Litterski & Mayrhofer 1998 (I); pl/co; ca.
- **Catillaria mediterranea* Hafellner - Litterski & Mayrhofer 1998 (III); mo; Cd, on *Physcia semipinnata* and *Physcia stellaris*.
- Catillaria nigroclavata* (Nyl.) Schuler - Litterski 1997 (III); mo; cone (Pi).
- Chromatochlamys muscorum* (Fr.) H. Mayrhofer & Poelt - Litterski & Mayrhofer 1998 (III); mo; mos.
- Cladonia cariosa* (Ach.) Spreng. - Litterski 1997 (III); mo; Pi.
- Cladonia cervicornis* (Ach.) Flot. - Krempelhuber 1865 (IV), Litterski & Mayrhofer 1998 (I, III); pl/co; so.
- Cladonia convoluta* (Lam.) Anders - Flotow 1843 (*C. endiviaefolia*), Krempelhuber 1865 (IV, V, *C. endivifolia*), Güvenc & Öztürk 1994 (IV, V), Litterski 1997 (I, III); pl/co, submo; so.
- Cladonia fimbriata* (L.) Fr. - Güvenc & Öztürk 1994 (IV), Litterski 1997 (III); pl/co, mo; Pi.
- Cladonia foliacea* (Huds.) Willd. - Güvenc & Öztürk 1994 (IV, V); pl/co; so.
- Cladonia furcata* (Huds.) Schrad. - Flotow 1843; -;
- Cladonia pocillum* (Ach.) O. J. Rich. - Flotow 1843 (*C. pyxidata* c. *p.*), Güvenc & Öztürk 1994 (IV, V), Litterski 1997 (I, III); pl/co, submo, mo; so.
- Cladonia pyxidata* (L.) Hoffm. - Krempelhuber 1865 (IV, V), Güvenc & Öztürk 1994 (V, *C. chlorophaea*, *C. pyxidata*), Aptroot 1996 (III); pl/co, [mo]; Pi, so.
- Cladonia ramulosa* (With.) J. R. Laundon - Litterski & Mayrhofer 1998 (III); mo; so.
- Cladonia rangiformis* Hoffm. - Krempelhuber 1865 (IV), Güvenc & Öztürk 1994 (IV, V), Litterski 1997 (III); pl/co, submo, mo; so.
- Cladonia squamosa* Hoffm. - Litterski 1997 (III); mo; so.
- Clauzadea immersa* (Hoffm.) Hafellner & Bellem. - Litterski & Mayrhofer 1998 (I); pl/co; ca.
- Clauzadea metzleri* (Körb.) Clauzade & Cl. Roux - Aptroot 1996 (II), Litterski 1997 (I); pl/co; li, ro.
- Collema auriforme* (With.) Coppins & J. R. Laundon - Litterski 1997 (III), Litterski & Mayrhofer 1998 (III); submo, salp; mos, si.
- Collema crispum* (Huds.) Weber - Flotow 1843, Güvenc & Öztürk 1994 (V), Aptroot 1996 (II), Litterski 1997 (III), Litterski & Mayrhofer 1998 (III); pl/co, mo; ca, ro, si, so.
- Collema cristatum* (L.) Weber - Krempelhuber 1865 (IV, V), Degelius 1954 (II), Güvenc & Öztürk 1994 (V), Aptroot 1996 (II), Litterski 1997 (III), Litterski & Mayrhofer 1998 (I, III); pl/co, submo; ca, mos, ro, si, so.
- Collema limosum* (Ach.) Ach. - Aptroot 1996 (II);

- [pl/co]; so.
- Collema occultatum*** Bagl. - Litterski & Mayrhofer 1998 (III); submo; Al.
- Collema ryssoleum*** (Tuck.) A. Schneid. - Litterski & Mayrhofer 1998 (I); pl/co; sa.
- Collema subflaccidum*** Degel. - Litterski 1997 (III); submo, mo; Cd, Ce, Cu.
- Collema subnigrescens*** Degel. - Litterski 1997 (III); submo, mo; Ac, Cu, Qu.
- Collema tenax*** (Sw.) Ach. - Aptroot 1996 (II, III), Litterski 1997 (I, III); pl/co, submo; so.
- Cyphelium tigillare*** (Ach.) Ach. - Litterski 1997 (III); salp; Pi.
- **Dactylospora attendenda*** (Nyl.) Arnold - Litterski & Mayrhofer 1998 (III); salp; wo, on *Lecanora* sp.
- Dendroscocaulon umhausense*** (Auersw.) Degel. - Litterski & Mayrhofer 1998 (III); mo; Cd, Qu.
- Degelia plumbea*** (Lightf.) P. M. Jørg. & P. James - Litterski 1997 (III); mo; Cd, Qu.
- Dermatocarpon miniatum*** (L.) W. Mann - Krempelhuber 1865 (III, *Endocarpon m.*), Litterski 1997 (III); mo, [hmo]; si.
- Dermatocarpon nodulosum*** (Kremp.) Zahlbr. - Krempelhuber 1865 (III, *Endocarpon n.*), typus in M, teste: O. Breuß; [hmo]; ro.
- Dimerella tavaresiana*** Vězda - Litterski & Mayrhofer 1998 (I); pl/co; Pi.
- Diploicia canescens*** (Dicks.) A. Massal. - Güvenc & Öztürk 1995 (V), Litterski 1997 (I), Litterski & Mayrhofer 1998 (I); pl/co; ca, Ce, li, Ol, Pi, Pr, sa.
- Diploschistes actinostomus*** (Pers. ex Ach.) Zahlbr. - Litterski & Mayrhofer 1998 (I); pl/co; ca.
- Diploschistes candidissimus*** (Kremp.) Zahlbr. - Krempelhuber 1865 (V, *Limboria candidissima*), Lettau 1932 (*D. calcarius*), Lumbsch 1989, Litterski & Mayrhofer 1998 (I); pl/co; ca.
- Diploschistes diacapsis*** (Ach.) Lumbsch - Litterski 1997 (III); submo; so.
- Diploschistes gypsaceus*** (Ach.) Zahlbr. - Krempelhuber 1865 (V, *Urceolaria scruposa* var. *gypsacea*, var. *cretacea*), Aptroot 1996 (II); [pl/co]; so.
- Diploschistes muscorum*** (Scop.) R. Sant. - Flotow 1843 (*Urceolaria villarsii* var. *macrior*), Krempelhuber 1865 (V, *Urceolaria scruposa* var. *bryophila*, var. *arenaria*), Lumbsch 1989, Litterski 1997 (III); submo; so.
- Diploschistes ocellatus*** (Vill.) Norman - Flotow 1843 (*Urceolaria villarsii*), Krempelhuber 1865 (IV, *Urceolaria ocellata*), Lumbsch 1989, Litterski & Mayrhofer 1998 (III); pl/co; sa.
- Diploschistes scruposus*** (Vill.) Norman - Güvenc & Öztürk 1995 (IV, V); pl/co; ca.
- Diplotomma alboatrum*** (Hoffm.) Flot. - Krempelhuber 1865 (IV, V), Güvenc & Öztürk 1995 (IV, V, *D. alboatrum*, *D. epipolium*), Litterski 1997 (I, II, III, *D. alboatrum*, *D. epipolium*), Litterski & Mayrhofer 1998 (III); pl/co, mo; ca, Ce, li, Ol, ca, cone (Pi).
- Diplotomma venustum*** (Körb.) Körb. - Güvenc & Öztürk 1995 (V, *Buellia venusta*); pl/co; ca.
- Dirina ceratoniae*** (Ach.) Fr. - Tehler 1983, Litterski 1997 (I); pl/co; Ce.
- Dirina cretacea*** (Zahlbr.) Tehler - Tehler 1983 (I, IV), Roux 1991 (map 3); [pl/co]; ca.
- Dirina massiliensis*** Durieu & Mont. f. *massiliensis* - Krempelhuber 1865 (IV, *D. repanda*), Tehler 1983, Roux 1991 (map 2), Litterski & Mayrhofer 1998 (I); pl/co; ca, sa.
- Dirina massiliensis*** Durieu & Mont. f. *sorediata* (Müll. Arg.) Tehler - Tehler 1983, Roux 1991 (map 2), Follmann 1993 (*D. stenhammari*, map 7), Litterski & Mayrhofer 1998 (I); pl/co; ca, sa.
- Epiphloeia terrena*** (Nyl.) Trevis. - Aptroot 1996 (III); [mo]; so.
- Evernia prunastri*** (L.) Ach. - Krempelhuber 1865 (III), Güvenc & Öztürk 1994 (V), Aptroot 1996 (III), Litterski 1997 (III); submo, mo, [hmo]; Cd, Pi, Qu.
- Fulgensia bracteata*** (Hoffm.) Räsänen - Krempelhuber 1865 (IV, *Gyalolechia b.*); pl/co; sa.
- Fulgensia fulgens*** (Sw.) Elenkin - Krempelhuber 1865 (V, *Psoroma f.*), Güvenc & Öztürk 1995 (IV, V), Aptroot 1996 (II), Litterski 1997 (III); pl/co, submo; ca, so.
- Fulgensia fulgida*** (Nyl.) Szatala - Litterski 1997 (I, III); pl/co, submo; so.
- Fulgensia subbracteata*** (Nyl.) Poelt - Vězda 1988a (IV, No. 2191), Aptroot 1996 (II), Litterski & Mayrhofer 1998 (I, IV); pl/co; ro, so.
- Fuscopannaria olivacea*** (P. M. Jørg.) P. M. Jørg. - Litterski 1997 (III, *Pannaria o.*); mo; Cd, Qu.
- Fuscopannaria saubinetii*** (MONT.) P. M. Jørg. - Litterski & Mayrhofer 1998 (III); submo; Qu.
- Gyalecta jenensis*** (Batsch) Zahlbr. - Litterski 1997 (III); mo; si.
- Haematomma nemetzii*** J. Steiner - Litterski & Mayrhofer 1998 (I); pl/co; sa.
- Heterodermia speciosa*** (Wulfen) Trevis. - Øvstedal 1987 (III); salp; Pi.

- Hymenelia prevostii* (Duby) Kremp. – Aptroot 1996 (II); [pl/co]; ro.
- Hypocenomysce scalaris* (Lilj.) M. Choisy - Krempelhuber 1865 (III, *Psora ostreata*), Litterski 1997 (III); [hmo], salp; Pi.
- Hypogymnia farinacea* Zopf - Litterski 1997 (III); salp; Pi.
- Hypogymnia tubulosa* (Schaer.) Hav. - Krempelhuber 1865 (III, *Parmelia ceratophylla* var. *t.*), Øvstedal 1987 (III), Litterski 1997 (III); mo, salp; Pi.
- Immersaria athrocarpa* (Ach.) Rambold & Pietschm. - Krempelhuber 1865 (III, *Lecidea conformis*); [mo]; si.
- **Kirschsteiniethelia aethiops* (Berk. & M. A. Curtis) D. Hawksw. - Aptroot 1996 (III); mo; -.
- Lasallia pustulata* (L.) Mérat - Litterski & Mayrhofer 1998 (I); pl/co; sa.
- Lecania cyrtella* (Ach.) Th. Fr. - Litterski & Mayrhofer 1998 (I); pl/co; br.
- Lecania inundata* (Hepp ex Körb.) M. Mayrhofer - Litterski 1997 (I); pl/co; sa.
- Lecania spadicea* (Flot.) Zahlbr. - Aptroot 1996 (II), Litterski & Mayrhofer 1998 (I); pl/co; ca, ro.
- Lecania sylvestris* (Arnold) Arnold - Aptroot 1996 (II); [pl/co]; ro.
- Lecania turicensis* (Hepp) Müll. Arg. - Litterski & Mayrhofer 1998 (I); pl/co; ca, sa.
- Lecanographa grumulosa* (Dufour) Egea & Torrente - Krempelhuber 1865 (V, *Lecanactis* g.), Litterski & Mayrhofer 1998 (I); pl/co; ca.
- Lecanographa lyncea* (Sm.) Egea & Torrente - Litterski & Mayrhofer 1998 (I); pl/co; Ce, Pi.
- Lecanora achariana* A. L. Sm. - Aptroot 1996 (II); [pl/co]; ro.
- Lecanora allophana* Nyl. - Krempelhuber 1865 (III, *Lecanora subfusca* var. *distans*), Güvenc & Öztürk 1995 (IV, V); pl/co; Cr, Cu.
- Lecanora argentata* (Ach.) Malme - Aptroot 1996 (III), Litterski 1997 (I, III); mo; Ac, Ce, Pi.
- Lecanora bolcana* (Pollini) Poelt - Krempelhuber 1865 (? , III, *Placodium saxicolum* var. *diffRACTUM*), Litterski 1997 (III), Litterski & Mayrhofer 1998 (I, III); pl/co, mo; sa, si.
- Lecanora campestris* (Schaer.) Hue var. *campestris* - Litterski & Mayrhofer 1998 (I); pl/co; ca.
- Lecanora campestris* (Schaer.) Hue var. *alba* de Lesd. - Vězda 1988a (IV, No. 2188); pl/co; ca.
- Lecanora chlarotera* Nyl. - Güvenc & Öztürk 1995 (IV, V, *L. chlarona*, *L. chlarotera*), Kondratyuk & Poelt 1997 (III); pl/co, mo; Ce, Cu, Ol, Pi, Ps, Qu.
- Lecanora dispersa* (L.) Sommerf. - Aptroot 1996 (III), Litterski 1997 (II); pl/co, [mo]; li, ro.
- Lecanora expallens* Ach. - Aptroot 1996 (III); [mo]; Pi.
- Lecanora hagenii* (Ach.) Ach. - Güvenc & Öztürk 1995 (V), Aptroot 1996 (III); pl/co, [mo]; Ps, Vi.
- Lecanora horiza* (Ach.) Linds. - Güvenc & Öztürk 1995 (IV, *L. sienae*), Litterski 1997 (I, III); pl/co, mo, salp; Ci, Ol, Pi, Qu.
- Lecanora intumescens* (Rebent.) Rabenh. - Aptroot 1996 (III); [mo]; Qu.
- Lecanora marginata* (Schaer.) Hertel & Rambold - Krempelhuber 1865 (IV, *Lecidea elata*); [pl/co]; -.
- Lecanora meridionalis* H. Magn. - Litterski & Mayrhofer 1998 (I); pl/co; Ce, Cs.
- Lecanora muralis* (Schreb.) Rabenh. - Krempelhuber 1865 (III, *Placodium saxicolum*), Güvenc & Öztürk 1995 (V), Litterski 1997 (II, III); pl/co, mo, salp; li, si.
- Lecanora piniperda* Körb. - Vězda 1988a (III, No. 2189), Aptroot 1996 (III); [mo], salp; Pi.
- Lecanora polytropa* (Hoffm.) Rabenh. - Aptroot 1996 (III); [mo]; ro.
- Lecanora pruinosa* Chaub. - Güvenc & Öztürk 1995 (V); pl/co; ro.
- Lecanora rupicola* (L.) Zahlbr. - Litterski 1997 (III), Litterski & Mayrhofer 1998 (III); submo, mo; si.
- Lecanora saligna* (Schrad.) Zahlbr. var. *sarcopsis* Ach. - Litterski & Mayrhofer 1998 (III); salp; Pi.
- Lecanora subintricata* (Nyl.) Th. Fr. - Litterski 1997 (III); salp; Pi.
- Lecanora sulphurea* (Hoffm.) Ach. - Litterski 1997 (III), Litterski & Mayrhofer 1998 (I); pl/co, mo; sa, si.
- Lecanora symmicta* (Ach.) Ach. - Litterski & Mayrhofer 1998 (I); pl/co; Cs.
- Lecanora umbrina* (Ach.) A. Massal. - Litterski 1997 (III); mo; Pi.
- Lecanora varia* (Hoffm.) Ach. - Krempelhuber 1865 (III), Aptroot 1996 (III), Litterski 1997 (III); mo; Pi.
- Lecidea atrobrunnea* (Lam. & DC.) Schaer. - Krempelhuber 1865 (III); [hmo]; -.
- Lecidea fuscoatra* (L.) Ach. - Krempelhuber 1865 (? , *L. fumosa*), Aptroot 1996 (III), Litterski 1997 (III); submo, mo; ro, si.
- Lecidea hypopta* Ach. - Øvstedal 1987 (III, cf.), Aptroot 1996 (III); [mo], salp (cf.); Pi.
- Lecidea lapicida* (Ach.) Ach. var. *pantherina* Ach.

- [pl/co]; so.
- Collema occultatum* Bagl. - Litterski & Mayrhofer 1998 (III); submo; Al.
- Collema rysssoleum* (Tuck.) A. Schneid. - Litterski & Mayrhofer 1998 (I); pl/co; sa.
- Collema subflaccidum* Degel. - Litterski 1997 (III); submo, mo; Cd, Ce, Cu.
- Collema subnigrescens* Degel. - Litterski 1997 (III); submo, mo; Ac, Cu, Qu.
- Collema tenax* (Sw.) Ach. - Aptroot 1996 (II, III), Litterski 1997 (I, III); pl/co, submo; so.
- Cyphelium tigillare* (Ach.) Ach. - Litterski 1997 (III); salp; Pi.
- **Dactylospora attendenda* (Nyl.) Arnold - Litterski & Mayrhofer 1998 (III); salp; wo, on *Lecanora* sp.
- Dendriscoaulon umhausense* (Auersw.) Degel. - Litterski & Mayrhofer 1998 (III); mo; Cd, Qu.
- Degelia plumbea* (Lightf.) P. M. Jørg. & P. James - Litterski 1997 (III); mo; Cd, Qu.
- Dermatocarpon miniatum* (L.) W. Mann - Krempelhuber 1865 (III, *Endocarpon m.*), Litterski 1997 (III); mo, [hmo]; si.
- Dermatocarpon nodulosum* (Kremp.) Zahlbr. - Krempelhuber 1865 (III, *Endocarpon n.*), typus in M, teste: O. Breuß; [hmo]; ro.
- Dimerella tavaresiana* Vězda - Litterski & Mayrhofer 1998 (I); pl/co; Pi.
- Diploicia canescens* (Dicks.) A. Massal. - Güvenc & Öztürk 1995 (V), Litterski 1997 (I), Litterski & Mayrhofer 1998 (I); pl/co; ca, Ce, li, Ol, Pi, Pr, sa.
- Diploschistes actinostomus* (Pers. ex Ach.) Zahlbr. - Litterski & Mayrhofer 1998 (I); pl/co; ca.
- Diploschistes candidissimus* (Kremp.) Zahlbr. - Krempelhuber 1865 (V, *Limboria candidissima*), Lettau 1932 (*D. calcarius*), Lumbsch 1989, Litterski & Mayrhofer 1998 (I); pl/co; ca.
- Diploschistes diacapsis* (Ach.) Lumbsch - Litterski 1997 (III); submo; so.
- Diploschistes gypsaceus* (Ach.) Zahlbr. - Krempelhuber 1865 (V, *Urceolaria scruposa* var. *gypsacea*, var. *cretacea*), Aptroot 1996 (II); [pl/co]; so.
- Diploschistes muscorum* (Scop.) R. Sant. - Flotow 1843 (*Urceolaria villarsii* var. *macrior*), Krempelhuber 1865 (V, *Urceolaria scruposa* var. *bryophila*, var. *arenaria*), Lumbsch 1989, Litterski 1997 (III); submo; so.
- Diploschistes ocellatus* (Vill.) Norman - Flotow 1843 (*Urceolaria villarsii*), Krempelhuber 1865 (IV, *Urceolaria ocellata*), Lumbsch 1989, Litterski & Mayrhofer 1998 (III); pl/co; sa.
- Diploschistes scruposus* (Vill.) Norman - Güvenc & Öztürk 1995 (IV, V); pl/co; ca.
- Diplotomma alboatrum* (Hoffm.) Flot. - Krempelhuber 1865 (IV, V), Güvenc & Öztürk 1995 (IV, V, *D. alboatrum*, *D. epipolium*), Litterski 1997 (I, II, III, *D. alboatrum*, *D. epipolium*), Litterski & Mayrhofer 1998 (III); pl/co, mo; ca, Ce, li, Ol, ca, cone (Pi).
- Diplotomma venustum* (Körb.) Körb. - Güvenc & Öztürk 1995 (V, *Buellia venusta*); pl/co; ca.
- Dirina ceratoniae* (Ach.) Fr. - Tehler 1983, Litterski 1997 (I); pl/co; Ce.
- Dirina cretacea* (Zahlbr.) Tehler - Tehler 1983 (I, IV), Roux 1991 (map 3); [pl/co]; ca.
- Dirina massiliensis* Durieu & Mont. f. *massiliensis* - Krempelhuber 1865 (IV, *D. repanda*), Tehler 1983, Roux 1991 (map 2), Litterski & Mayrhofer 1998 (I); pl/co; ca, sa.
- Dirina massiliensis* Durieu & Mont. f. *sorediata* (Müll. Arg.) Tehler - Tehler 1983, Roux 1991 (map 2), Follmann 1993 (*D. stenhammari*, map 7), Litterski & Mayrhofer 1998 (I); pl/co; ca, sa.
- Epiphloeia terrena* (Nyl.) Trevis. - Aptroot 1996 (III); [mo]; so.
- Evernia prunastri* (L.) Ach. - Krempelhuber 1865 (III), Güvenc & Öztürk 1994 (V), Aptroot 1996 (III), Litterski 1997 (III); submo, mo, [hmo]; Cd, Pi, Qu.
- Fulgensia bracteata* (Hoffm.) Räsänen - Krempelhuber 1865 (IV, *Gyalolechia b.*); pl/co; -.
- Fulgensia fulgens* (Sw.) Elenkin - Krempelhuber 1865 (V, *Psoroma f.*), Güvenc & Öztürk 1995 (IV, V), Aptroot 1996 (II), Litterski 1997 (III); pl/co, submo; ca, so.
- Fulgensia fulgida* (Nyl.) Szatala - Litterski 1997 (I, III); pl/co, submo; so.
- Fulgensia subbracteata* (Nyl.) Poelt - Vězda 1988a (IV, No. 2191), Aptroot 1996 (II), Litterski & Mayrhofer 1998 (I, IV); pl/co; ro, so.
- Fuscopannaria olivacea* (P. M. Jørg.) P. M. Jørg. - Litterski 1997 (III, *Pannaria o.*); mo; Cd, Qu.
- Fuscopannaria saubinetii* (MONT.) P. M. Jørg. - Litterski & Mayrhofer 1998 (III); submo; Qu.
- Gyalecta jenensis* (Batsch) Zahlbr. - Litterski 1997 (III); mo; si.
- Haematomma nemetzii* J. Steiner - Litterski & Mayrhofer 1998 (I); pl/co; sa.
- Heterodermia speciosa* (Wulfen) Trevis. - Øvstedal 1987 (III); salp; Pi.

- Hymenelia prevostii* (Duby) Kremp. – Aptroot 1996 (II); [pl/co]; ro.
- Hypocenomyce scalaris* (Lilj.) M. Choisy - Krempelhuber 1865 (III, *Psora ostreata*), Litterski 1997 (III); [hmo], salp; Pi.
- Hypogymnia farinacea* Zopf - Litterski 1997 (III); salp; Pi.
- Hypogymnia tubulosa* (Schaer.) Hav. - Krempelhuber 1865 (III, *Parmelia ceratophylla* var. *t.*), Øvstedal 1987 (III), Litterski 1997 (III); mo, salp; Pi.
- Immersaria athrocarpa* (Ach.) Rambold & Pietschm. - Krempelhuber 1865 (III, *Lecidea conformis*); [mo]; si.
- **Kirschsteiniothelia aethiops* (Berk. & M. A. Curtis) D. Hawksw. - Aptroot 1996 (III); mo; -.
- Lasallia pustulata* (L.) Mérat - Litterski & Mayrhofer 1998 (I); pl/co; sa.
- Lecania cyrtella* (Ach.) Th. Fr. - Litterski & Mayrhofer 1998 (I); pl/co; br.
- Lecania inundata* (Hepp ex Körb.) M. Mayrhofer - Litterski 1997 (I); pl/co; sa.
- Lecania spadicea* (Flot.) Zahlbr. - Aptroot 1996 (II), Litterski & Mayrhofer 1998 (I); pl/co; ca, ro.
- Lecania sylvestris* (Arnold) Arnold - Aptroot 1996 (II); [pl/co]; ro.
- Lecania turicensis* (Hepp) Müll. Arg. - Litterski & Mayrhofer 1998 (I); pl/co; ca, sa.
- Lecanographa grumulosa* (Dufour) Egea & Torrente - Krempelhuber 1865 (V, *Lecanactis* g.), Litterski & Mayrhofer 1998 (I); pl/co; ca.
- Lecanographa lyncea* (Sm.) Egea & Torrente - Litterski & Mayrhofer 1998 (I); pl/co; Ce, Pi.
- Lecanora achariana* A. L. Sm. - Aptroot 1996 (II); [pl/co]; ro.
- Lecanora allophana* Nyl. - Krempelhuber 1865 (III, *Lecanora subfusca* var. *distans*), Güvenc & Öztürk 1995 (IV, V); pl/co; Cr, Cu.
- Lecanora argentata* (Ach.) Malme - Aptroot 1996 (III), Litterski 1997 (I, III); mo; Ac, Ce, Pi.
- Lecanora bolcana* (Pollini) Poelt - Krempelhuber 1865 (? , III, *Placodium saxicolum* var. *diffRACTUM*), Litterski 1997 (III), Litterski & Mayrhofer 1998 (I, III); pl/co, mo; sa, si.
- Lecanora campestris* (Schaer.) Hue var. *campestris* - Litterski & Mayrhofer 1998 (I); pl/co; ca.
- Lecanora campestris* (Schaer.) Hue var. *alba* de Lesd. - Vězda 1988a (IV, No. 2188); pl/co; ca.
- Lecanora chlarotera* Nyl. - Güvenc & Öztürk 1995 (IV, V, *L. chlarona*, *L. chlarotera*), Kondratyuk & Poelt 1997 (III); pl/co, mo; Ce, Cu, Ol, Pi, Ps, Qu.
- Lecanora dispersa* (L.) Sommerf. - Aptroot 1996 (III), Litterski 1997 (II); pl/co, [mo]; li, ro.
- Lecanora expallens* Ach. - Aptroot 1996 (III); [mo]; Pi.
- Lecanora hagenii* (Ach.) Ach. - Güvenc & Öztürk 1995 (V), Aptroot 1996 (III); pl/co, [mo]; Ps, Vi.
- Lecanora horiza* (Ach.) Linds. - Güvenc & Öztürk 1995 (IV, *L. sienae*), Litterski 1997 (I, III); pl/co, mo, salp; Ci, Ol, Pi, Qu.
- Lecanora intumescens* (Rebent.) Rabenh. - Aptroot 1996 (III); [mo]; Qu.
- Lecanora marginata* (Schaer.) Hertel & Rambold - Krempelhuber 1865 (IV, *Lecidea elata*); [pl/co]; -.
- Lecanora meridionalis* H. Magn. - Litterski & Mayrhofer 1998 (I); pl/co; Ce, Cs.
- Lecanora muralis* (Schreb.) Rabenh. - Krempelhuber 1865 (III, *Placodium saxicolum*), Güvenc & Öztürk 1995 (V), Litterski 1997 (II, III); pl/co, mo, salp; li, si.
- Lecanora piniperda* Körb. - Vězda 1988a (III, No. 2189), Aptroot 1996 (III); [mo], salp; Pi.
- Lecanora polytropa* (Hoffm.) Rabenh. - Aptroot 1996 (III); [mo]; ro.
- Lecanora pruinosa* Chaub. - Güvenc & Öztürk 1995 (V); pl/co; ro.
- Lecanora rupicola* (L.) Zahlbr. - Litterski 1997 (III), Litterski & Mayrhofer 1998 (III); submo, mo; si.
- Lecanora saligna* (Schr.) Zahlbr. var. *sarcopsis* Ach. - Litterski & Mayrhofer 1998 (III); salp; Pi.
- Lecanora subintricata* (Nyl.) Th. Fr. - Litterski 1997 (III); salp; Pi.
- Lecanora sulphurea* (Hoffm.) Ach. - Litterski 1997 (III), Litterski & Mayrhofer 1998 (I); pl/co, mo; sa, si.
- Lecanora symmicta* (Ach.) Ach. - Litterski & Mayrhofer 1998 (I); pl/co; Cs.
- Lecanora umbrina* (Ach.) A. Massal. - Litterski 1997 (III); mo; Pi.
- Lecanora varia* (Hoffm.) Ach. - Krempelhuber 1865 (III), Aptroot 1996 (III), Litterski 1997 (III); mo; Pi.
- Lecidea atrobrunnea* (Lam. & DC.) Schaer. - Krempelhuber 1865 (III); [hmo]; -.
- Lecidea fuscoatra* (L.) Ach. - Krempelhuber 1865 (? , *L. fumosa*), Aptroot 1996 (III), Litterski 1997 (III); submo, mo; ro, si.
- Lecidea hypopta* Ach. - Øvstedal 1987 (III, cf.), Aptroot 1996 (III); [mo], salp (cf.); Pi.
- Lecidea lapicida* (Ach.) Ach. var. *pantherina* Ach.

- Krempelhuber 1865 (III, *L. polycarpa*); [mo, hmo]; -.
- Lecidea lurida*** Ach. - Güvenc & Öztürk 1995 (V, *Psora l.*); submo; ca.
- Lecidea sarcogynoides*** Körb. - Litterski & Mayrhofer 1998 (I); pl/co; sa.
- Lecidella achristotera*** (Nyl.) Hertel & Leuckert - Güvenc & Öztürk 1995 (V); pl/co; Cr, Ps.
- Lecidella asema*** (Nyl.) Knoph & Hertel - Güvenc & Öztürk 1995 (V); pl/co; ca.
- Lecidella carpathica*** Körb. - Aptroot 1996 (III), Litterski 1997 (III); [mo], salp; ro, si.
- Lecidella elaeochroma*** (Ach.) M. Choisy - Krempelhuber 1865 (III, *Lecidea enteroleuca* var. *rugulosa*), Güvenc & Öztürk 1995 (IV, V), Litterski 1997 (III), Litterski & Mayrhofer 1998 (III); pl/co, mo, [hmo], salp; Ac, Ce, Ci, Cu, Ol, Pi, Ps, Qu.
- Lecidella euphorea*** (Flörke) Hertel - Krempelhuber 1865 (V, *Lecidea sabuletorum* var. *e.*), Güvenc & Öztürk 1995 (IV), Aptroot 1996 (III), Kondratyuk & Poelt 1997 (III); pl/co, mo; Ce, Pi, Qu.
- Lecidella stigmathea*** (Ach.) Hertel & Leuckert - Aptroot 1996 (III); [mo]; ro.
- Lempholemma chalazanum*** (Ach.) de Lesd. - Krempelhuber 1865 (III, *Physma compactum* - *Collema ch.*); [mo]; -.
- Lempholemma elveloideum*** (Ach.) Zahlbr. - Litterski & Mayrhofer 1998 (I); pl/co; ca.
- Lepraria crassissima*** (Hue) Lettau var. *isidiata* Llimona - Vězda 1988a (IV, No. 2192), Güvenc & Öztürk 1995 (V); pl/co; ca.
- Lepraria incana*** (L.) Ach. - Litterski 1997 (III); mo; Pi.
- Lepraria nivalis*** J. R. Laundon - Litterski 1997 (III), Litterski & Mayrhofer 1998 (I); pl/co, submo, mo; ca, si, so.
- Leprocaulon microscopicum*** (Vill.) D. Hawksw. - Litterski 1997 (III); submo, mo; so (among si).
- Leproloma diffusum*** J. R. Laundon - Aptroot 1996 (III); [mo]; ro.
- Leptochidium albociliatum*** (Desm.) M. Choisy - Krempelhuber 1865 (III, *Collema a.*), Aptroot 1996 (III), Litterski 1997 (III); mo, [hmo]; mos, si, so.
- Leptogium biatorinum*** (Nyl.) Leight. - Aptroot 1996 (II); [pl/co]; ro, so.
- Leptogium corniculatum*** (Hoffm.) Minks - Aptroot 1996 (III); [mo]; so.
- Leptogium gelatinosum*** (With.) J. R. Laundon - Litterski 1997 (III), Litterski & Mayrhofer 1998 (III); submo, mo; mos.
- Leptogium lichenoides*** (L.) Zahlbr. - Krempelhuber 1865 (III, V, *L. atrocaeruleum*), Aptroot 1996 (III), Litterski 1997 (III); mo; Cd, ro, wo.
- Leptogium subtile*** (Schrader) Torss. - Litterski & Mayrhofer 1998 (III); submo, mo; Qu.
- Leptogium teretiusculum*** (Wallr.) Arnold - Litterski & Mayrhofer 1998 (III); pl/co; so.
- Leptogium turgidum*** (Ach.) Cromb. - Krempelhuber 1865 (I, *Collema t.*); [pl/co]; -.
- Letharia vulpina*** (L.) Hue - Krempelhuber 1865 (III, *Evernia v.*), Petrak *et al.* 1941, Schade 1954 (III), Gams 1955, Hafellner 1987, Øvstedal 1987 (III), Vězda 1988b (III, No. 2250), Litterski 1997 (III); [hmo], salp; Pi.
- **Lichenodiplis lecanorae*** (Vouaux) Dyko & D. Hawksw. - Aptroot 1996 (III); [mo]; on *Caloplaca cerina*.
- **Lichenostigma elongata*** Nav.-Ros. & Hafellner - Navarro-Rosinés & Hafellner 1996 (IV); pl/co; on *Lobothallia radiosa*.
- Lobothallia prae-radiosa*** (Nyl.) Hafellner - Litterski 1997 (III, *Aspicilia p.*); salp; si.
- Lobothallia radiosa*** (Hoffm.) Hafellner - Krempelhuber 1865 (II, *Placodium radiosum*), Hermann *et al.* 1973 (II, *Lecanora r.*), Litterski & Mayrhofer 1998 (III), Navarro-Rosinés & Hafellner 1996 (IV); pl/co, salp; si.
- Megaspora verrucosa*** (Ach.) Hafellner & V. Wirth var. *mutabilis* Nimis & Cl. Roux - Øvstedal 1987 (III, *Aspicilia m.*), Litterski 1997 (III); salp; Jn, Pi.
- Melanelia elegantula*** (Zahlbr.) Essl. - Litterski 1997 (III); salp; Pi.
- Melanelia exasperata*** (De Not.) Essl. - Krempelhuber 1865 (III, *P. aspera*), Esslinger 1977 (*Parmelia e.*, fig. 15), Litterski 1997 (III); mo, [hmo], salp; Pi, Qu.
- Melanelia exasperatula*** (Nyl.) Essl. - Øvstedal 1987 (III); salp; Pi.
- Melanelia glabra*** (Schaer.) Essl. - Litterski 1997 (III); mo; Ce, Cu, Qu.
- Melanelia glabratula*** (Lamy) Essl. - Litterski 1997 (III); mo; Pi.
- Melanelia laciniatula*** (H. Olivier) Essl. - Litterski 1997 (III); mo; Cd, Pi, Qu.
- Micarea erratica*** (Körb.) Hertel, Rambold & Pietschm. - Aptroot 1996 (III); [mo]; ro.
- Micarea misella*** (Nyl.) Hedl. - Litterski & Mayrhofer 1998 (III); pl/co; wo/Pi.
- Micarea prasina*** Fr. - Litterski 1997 (III); mo; cone (Pi).
- Moelleropsis nebulosa*** (Hoffm.) Gyeln. - Litterski & Mayrhofer 1998 (III); submo; so.

- Mycobilimbia hypnorum* (Lib.) Kalb & Hafellner - Litterski & Mayrhofer 1998 (III); pl/co; mo/so.
- Mycobilimbia sphaeroides* (Dicks.) ined. - Flotow 1843 (*Lecidea sabuletorum*, *pilularis*); -, br.
- Naetrocymbe punctiformis* A. Massal. - Aptroot 1996 (III, *Arthopyrenia* p.), Litterski & Mayrhofer 1998 (I, *Arthopyrenia* p.); pl/co, [mo]; Ne, Qu.
- Neofuscelia luteonotata* (J. Steiner) Essl. - Vězda 1988a (IV, No. 2195, *Parmelia* l.); pl/co; si.
- Neofuscelia pulla* (Ach.) Essl. - Güvenc & Öztürk 1994 (V, *Parmelia* p.), Aptroot 1996 (III), Litterski 1997 (III), Litterski & Mayrhofer 1998 (I, III); pl/co, submo, mo; ro, si.
- Nephroma laevigatum* Ach. - Litterski 1997 (III); mo; Cd, Qu.
- Ochrolechia balcanica* Verseghe - Litterski 1997 (III); mo; Qu.
- Ochrolechia pallescens* (L.) A. Massal. - Litterski 1997 (III); mo; Cd, Pi.
- Ochrolechia parella* (L.) A. Massal. - Litterski 1997 (III); submo; si.
- Opegrapha calcarea* Turner ex Sm. & Sowerby - Litterski & Mayrhofer 1998 (I); pl/co; ca.
- Opegrapha durieui* Mont. - Roux & Egea 1992 (IV); pl/co; ca.
- **Opegrapha physciaria* (Nyl.) D. Hawksw. & Coppins - Litterski & Mayrhofer 1998 (I); pl/co; on *Xanthoria parietina* (thallus).
- Opegrapha rupestris* Pers. - Krempelhuber 1865 (III); [submo]; sa.
- Opegrapha varia* Pers. - Litterski 1997 (I), Litterski & Mayrhofer 1998 (*O. v.* var. *herbarum*, var. *varia*, I, III); pl/co, submo; Ce, Jn, Qu.
- Opegrapha variaeformis* Anzi - Roux & Egea 1992 (IV); pl/co; ca.
- Parmelia saxatilis* (L.) Ach. - Krempelhuber 1865 (III), Litterski 1997 (III); mo, [hmo]; Pi, si.
- Parmelia submontana* Nadv. ex Hale - Litterski 1997 (III); mo; Cd, Pi.
- Parmelina carporrhizans* (Taylor) Poelt & Vězda - Krempelhuber 1865 (III, *Parmelia* c.); [mo]; -
- Parmelina pastillifera* (Harm.) Hale - Litterski 1997 (III); mo; si.
- Parmelina quercina* (Willd.) Hale - Aptroot 1996 (III); [mo]; Qu.
- Parmelina tiliacea* (Hoffm.) Hale - Krempelhuber 1865 (? , III, *Parmelia quercifolia* a t.), Güvenc & Öztürk 1994 (V, *Parmelia* t.), Litterski 1997 (III); submo, mo; Cd, Ce, Pi.
- Parmeliella triptophylla* (Ach.) Müll. Arg. - Flotow 1843 (*Biatora t.* var. *corallinoides*); -, -.
- Parmeliopsis ambigua* (Wulfen) Nyl. - Litterski 1997 (III); salp; Pi.
- Peltigera canina* (L.) Willd. - Litterski 1997 (III); mo; so.
- Peltigera neckeri* Müll. Arg. - Krempelhuber 1865 (I, as *P. polydactyla*, WU, teste), Litterski 1997 (III); [pl/co], mo; so.
- Peltigera rufescens* (Weiss) Humb. - Aptroot 1996 (III), Litterski 1997 (III); mo; so.
- Pertusaria albescens* (Huds.) M. Choisy & Werner - Güvenc & Öztürk 1995 (V), Litterski 1997 (III); pl/co, submo, mo; ca, Cd, Cu, Pi, Qu, mos, si.
- Pertusaria amara* (Ach.) Nyl. - Güvenc & Öztürk 1995 (V); pl/co; Cu, so.
- Pertusaria coccodes* (Ach.) Nyl. - Litterski 1997 (III); mo; Cd, Pi.
- Pertusaria flavicans* Lamy - Litterski & Mayrhofer 1998 (I); pl/co; sa.
- Pertusaria hemisphaerica* (Flörke) Erichsen - Litterski & Mayrhofer 1998 (III); mo; Cd, Qu.
- Pertusaria hymenea* (Ach.) Schaer. - Litterski 1997 (III); mo; Ce, Qu.
- Pertusaria leucostoma* (Bernh.) A. Massal. - Güvenc & Öztürk 1995 (V); pl/co; Ps.
- Pertusaria maximiliana* Klem. - Litterski & Mayrhofer 1998 (I); pl/co; Jn.
- Pertusaria pertusa* auct. - Güvenc & Öztürk 1995 (V), Litterski 1997 (III); submo, mo; Cd, Ce, Cu, Qu.
- Pertusaria rupicola* (Fr.) Harm. var. *coralloidea* (Anzi) Croz. - Litterski & Mayrhofer 1998 (III); mo; si.
- **Phacopsis vulpina* Tul. - Hafellner 1987, Vězda 1988b (III, No. 2250); salp; on thallus of *Letharia vulpina*.
- Phaeophyscia orbicularis* (Neck.) Moberg - Krempelhuber 1865 (III, *Anaptychia obscura* var. *o.*), Kondratyuk & Poelt 1997 (III); mo; Qu.
- Phaeophyscia poeltii* (Frey) Nimis - Aptroot 1996 (III); [mo]; Qu.
- Physcia adscendens* (Fr.) H. Olivier - Krempelhuber 1865 (III, *Anaptychia stellaris* v. *hispida*), Güvenc & Öztürk 1994 (IV, V), Aptroot 1996 (III), Litterski 1997 (III), Litterski & Mayrhofer 1998 (III); pl/co, submo, mo, salp; Ce, Cr, Cu, Ol, Pi, Ps, Qu, ro.
- Physcia aipolia* (Ehrh. ex. Humb.) Fürnr. - Krempelhuber 1865 (III, *Anaptychia stellaris* a. a.), Güvenc & Öztürk 1994 (V), Øvstedal 1987 (III), Aptroot 1996 (III), Kondratyuk & Poelt 1997 (III); pl/co, mo, [hmo], salp; Ce, Pi, Qu.

- Physcia biziana* (A. Massal.) Zahlbr. - Aptroot 1996 (III), Litterski 1997 (III), Litterski & Mayrhofer 1998 (I); pl/co, [mo], salp; ca, Jn, Pi
- Physcia caesia* (Hoffm.) Fürnr. - Litterski 1997 (I, III); pl/co, mo; li, si.
- Physcia dimidiata* (Arnold) Nyl. - Litterski & Mayrhofer 1998 (III); pl/co; si.
- Physcia dubia* (Hoffm.) Lettau - Krempelhuber 1865 (III, *Anaptychia pulchella* var. *d.*); [mo, hmo]; ro.
- Physcia scopulorum* (Lambinon & Vězda) Poelt & Nimis - Litterski & Mayrhofer 1998 (I); pl/co; ca.
- Physcia semipinnata* (J. F. Gmel.) Moberg - Güvenc & Öztürk 1994 (IV, V), Aptroot 1996 (III), Kondratyuk & Poelt 1997 (III), Litterski & Mayrhofer 1998 (III); pl/co, submo, mo, salp; br, Ce, Cd, Cr, Cu, Ol, Pi, Ps, Qu.
- Physcia stellaris* (L.) Nyl. - Krempelhuber 1865 (III, *Anaptychia* st. *B. ambigua*), Aptroot 1996 (III), Litterski 1997 (III), Litterski & Mayrhofer 1998 (III); pl/co, mo, [hmo], salp; mos, si, Cd, Ol, Qu.
- Physcia tenella* (Scop.) DC. - Güvenc & Öztürk 1994 (IV, V), Aptroot 1996 (III); pl/co, submo, mo; Ce, Cr, Pi, Ps.
- Physconia americana* Essl. - GZU (III) (Troodos, Cedar Valley, on *Cedrus*, ca. 1000 m, leg. H. Mayrhofer 1987, det./rev. Th. Esslinger 1997); mo; Cd.
- Physconia distorta* (With.) J. R. Laundon - Krempelhuber 1865 (III, *Anaptychia pulverulenta* var. *musciicola*), Güvenc & Öztürk 1994 (V), Aptroot 1996 (III), Litterski 1997 (III), Kondratyuk & Poelt 1997 (III); submo, mo, hmo; Cd, Ce, Cu, Pi, Qu.
- Physconia enteroxantha* (Nyl.) Poelt - Litterski 1997 (III), Kondratyuk & Poelt 1997 (III); mo; Qu, si.
- Physconia grisea* (Lam.) Poelt - Krempelhuber 1865 (III, *Anaptychia pulverulenta* var. *g.*), Øvstedal 1987 (III); salp; Pi.
- Physconia muscigena* (Ach.) Poelt - Krempelhuber 1865 (III, *Anaptychia pulverulenta* var. *angustata*); [mo, hmo]; Jn.
- Physconia perisidiosa* (Erichsen) Moberg - Litterski 1997 (III); mo; Qu.
- Physconia subpulverulenta* (Szatala) Poelt - Litterski 1997 (III); mo; Qu.
- Physconia venusta* (Ach.) Poelt - Krempelhuber 1865 (III, *Anaptychia pulverulenta* var. *v.*); [mo]; -.
- Placidium adami-borosi* Szatala - Breuss 1990 (III, *Catapyrenium a.*); [hmo]; -.
- Placidium michelii* A. Massal. - Aptroot 1996 (II, *Catapyrenium m.*); [pl/co]; ro.
- Placidium rufescens* (Ach.) A. Massal. - Krempelhuber 1865 (III); [hmo]; -.
- Placidium semaforonense* (Breuss) Breuss - Breuss 1990 (I, *Catapyrenium s.*), Aptroot 1996 (II, *Catapyrenium s.*); [pl/co]; so.
- Placidium tenellum* (Breuss) Breuss - Breuss 1990 (V, *Catapyrenium t.*); [pl/co]; so.
- Placynthiella icmalea* (Ach.) Coppins & P. James - Aptroot 1996 (III); [mo]; Pi.
- Placynthium nigrum* (Huds.) Gray - Litterski & Mayrhofer 1998 (I, III); pl/co; sa, si.
- Platismatia glauca* (L.) W. L. Culb. et C. F. Culb. - Krempelhuber 1865 (III, *Cetraria g.*), Litterski 1997 (III); mo, [hmo]; Cd, Pi, Qu.
- Pleurosticta acetabulum* (Neck.) Elix & Lumbsch - Güvenc & Öztürk 1994 (V, *Parmelia a.*); submo; Cu.
- Porina linearis* (Leight.) Zahlbr. - Litterski & Mayrhofer 1998 (I); pl/co; ca.
- Porpidia melinodes* (Körb.) Gowan & P. James - Aptroot 1996 (III); [mo]; ro.
- Protoparmelia montagnei* (Fr.) Poelt & Nimis - Krempelhuber 1865 (I, II, *Placodium m.* var. *calcareo*), Litterski & Mayrhofer 1998 (I); pl/co; sa.
- Pseudevernia furfuracea* (L.) Zopf - Krempelhuber 1865 (III, *Evernia f.*), Øvstedal 1987 (III), Litterski 1997 (III); mo, [hmo], salp; Pi, Qu.
- Psora decipiens* (Hedw.) Hoffm. - Flotow 1843 (*Biatora d.*), Krempelhuber 1865 (?), Güvenc & Öztürk 1995 (IV, V), Aptroot 1996 (II), Litterski 1997 (II, III); pl/co, submo; ca, so.
- Psora testacea* Hoffm. - Krempelhuber 1865 (III, *Psora t.* var. *turgida*); [mo]; -.
- Pyrenula chlorospila* Arnold - Litterski & Mayrhofer 1998 (I); pl/co; Ce.
- Pyrhospora quernea* (Dicks.) Körb. - Litterski & Mayrhofer 1998 (I); pl/co; Cs, Jn.
- Ramalina breviscula* Nyl. - Litterski & Mayrhofer 1998 (I); pl/co; sa, si.
- Ramalina calicaris* (L.) Fr. - Litterski 1997 (III), Litterski & Mayrhofer 1998 (I); pl/co, mo; Cd, Pr, Qu.
- Ramalina canariensis* J. Steiner - Güvenc & Öztürk 1994 (IV), Litterski 1997 (I), Litterski & Mayrhofer 1998 (I); pl/co; Ce, Pi, Ps, Pr.
- Ramalina capitata* (Ach.) Nyl. var. *digitellata* (Nyl.) Nimis - Litterski & Mayrhofer 1998 (I); pl/co; sa.
- Ramalina farinacea* (L.) Ach. - Krempelhuber

- 1865 (III), Güvenc & Öztürk 1994 (IV, V), Litterski 1997 (III), Litterski & Mayrhofer 1998 (I); pl/co, submo, mo, [hmo]; Pi, Pr, Qu.
- Ramalina fastigiata* (Pers.) Ach. - Güvenc & Öztürk 1994 (V), Litterski & Mayrhofer 1998 (III); pl/co, submo, mo; Cd, Cr, Ol, Pi, Ps, Qu.
- Ramalina fraxinea* (L.) Ach. - Krempelhuber 1865 (?), Güvenc & Öztürk 1994 (V); pl/co, submo; Cr, Pi, Ps.
- Ramalina obtusata* (Arnold) Bitter - Güvenc & Öztürk 1994 (V); pl/co; Ce, Cu, Ol, Pi, Ps.
- Ramalina pollinaria* (Westr.) Ach. - Krempelhuber 1865 (?); -, -.
- Ramalina pusilla* Le Prévost - Krempelhuber 1865 (V), Litterski & Mayrhofer 1998 (I); pl/co; Jn
- Ramalina roesleri* (Schaer.) Hue - Güvenc & Öztürk 1994 (V); pl/co, submo; Ce, Ps.
- Ramalina subfarinacea* (Cromb.) Nyl. - Güvenc & Öztürk 1994 (V); submo; Pi.
- Rhizocarpon geographicum* (L.) DC. - Krempelhuber 1865 (III), Aptroot 1996 (III), Litterski 1997 (III); submo, mo, hmo; ro, si.
- Rhizocarpon lecanorinum* Anders - Litterski 1997 (III); mo; si.
- Rhizocarpon macrosporum* Räsänen - Runemark 1956 (III); [mo]; -.
- Rhizocarpon obscuratum* (Ach.) A. Massal. - Aptroot 1996 (III); [mo]; ro.
- Rhizocarpon polycarpum* (Hepp) Th. Fr. - Litterski 1997 (III); submo; si.
- Rhizocarpon umbilicatum* (Ramond) Flagey - Aptroot 1996 (III); [mo]; ro.
- Rhizocarpon viridiatrum* (Wulfen) Körb. - Litterski & Mayrhofer 1998 (III); pl/co; sa.
- Rhizoplaca chrysouleuca* (Sm.) Zopf - Güvenc & Öztürk 1995 (V); pl/co; ca.
- Rimularia insularis* (Nyl.) Rambold & Hertel - Litterski & Mayrhofer 1998 (III); mo; si, on *Lecanora rupicola*.
- Rinodina bischoffii* (Hepp) A. Massal. - Litterski & Mayrhofer 1998 (I, IV); pl/co; ca.
- Rinodina capensis* Hampe - Giralt & Mayrhofer 1994 (III); mo; Cd.
- Rinodina dalmatica* Zahlbr. - Litterski & Mayrhofer 1998 (I); pl/co; Pi.
- Rinodina dubyana* (Hepp) J. Steiner - Litterski & Mayrhofer 1998 (I); pl/co; ca.
- Rinodina exigua* (Ach.) Gray - Güvenc & Öztürk 1995 (IV), Aptroot 1996 (III), Litterski 1997 (III); pl/co, submo, mo; Cd, Cu, Pi, Ps, Qu.
- Rinodina gennarii* Bagl. - Litterski & Mayrhofer 1998 (III); pl/co; sa.
- Rinodina guzzinii* Jatta - Litterski & Mayrhofer 1998 (III); pl/co; sa.
- Rinodina llimonae* Giralt & Etayo - Giralt & Mayrhofer 1995 (III); salp; Jn, Pi.
- Rinodina oxydata* (A. Massal.) A. Massal. - Aptroot 1996 (III); [mo]; ro.
- Rinodina pyrina* (Ach.) Arnold - Giralt & Mayrhofer 1995 (IV), Güvenc & Öztürk 1995 (IV), Aptroot 1996 (III), Litterski 1997 (III); pl/co, mo; Ce, Pi, Qu, Vi.
- Rinodina septentrionalis* Malm - Giralt & Mayrhofer 1995 (III); mo, salp; Pi, Qu.
- Rinodina sophodes* (Ach.) A. Massal. - Giralt & Mayrhofer 1995 (III), Aptroot 1996 (III), Litterski 1997 (III); mo, salp; Cd, Qu.
- Rinodina teichophila* (Nyl.) Arnold - Litterski & Mayrhofer 1998 (I); pl/co; sa.
- Rinodina trachytica* (A. Massal.) Bagl. & Carestia - Mayrhofer *et al.* 1992 (III); [submo]; si.
- Roccella phycopsis* Ach. - Güvenc & Öztürk 1994 (V), Litterski & Mayrhofer 1998 (I); pl/co; ca, Pi, sa.
- Sarcogyne regularis* Körb. - Aptroot 1996 (II), Litterski & Mayrhofer 1998 (III, IV); pl/co; ca, ro, si.
- Schismatomma decolorans* (Sm.) Clauzade & Vězda - Litterski & Mayrhofer 1998 (I); pl/co; Ce.
- Solenopsora candicans* (Dicks.) J. Steiner - Güvenc & Öztürk 1995 (V); submo; ca.
- Solenopsora cesatii* A. Massal. var. *cesatii* - Litterski & Mayrhofer 1998 (I); pl/co; ca.
- Solenopsora cesatii* A. Massal. var. *grisea* (Bagl.) Nimis - Güvenc & Öztürk 1995 (V, *S. bagliettoana*), Litterski & Mayrhofer 1998 (I); pl/co; ca, mo.
- Solenopsora olivacea* (Fr.) H. Kiliass - Krempelhuber 1865 (V, *Biatorina Micheletiana*); [pl/co]; -.
- Squamarina cartilaginea* (With.) P. James - Flotow 1843 (*Lecanora crassa*, *Lecanora crassa* f. *macra*), Krempelhuber 1865 (IV, *Psoroma crassum* β. *caespitosum*), Güvenc & Öztürk 1995 (IV, V), Aptroot 1996 (II), Litterski 1997 (I, III); pl/co; ca, mo, so.
- Squamarina concrescens* (Müll. Arg.) Poelt - Litterski 1997 (III), Litterski & Mayrhofer 1998 (I); pl/co, submo; so.
- Squamarina lentigera* (Weber) Poelt - Flotow 1843 (*Lecanora l.*), Krempelhuber 1865 (IV, *Psoroma lentigerum*), Güvenc & Öztürk 1995 (IV), Aptroot 1996 (II), Litterski 1997 (III), Litterski & Mayrhofer 1998 (I, III); pl/co, submo; ca, so.

- Staurothele rufa* (A. Massal.) Zschacke - Aptroot 1996 (II); [pl/co]; ro.
- Tephromela atra* (Huds.) Hafellner - Flotow 1843 (*Lecanora a.*), Krempelhuber 1865 (III, IV, *Lecanora a.*), Litterski 1997 (III), Litterski & Mayrhofer 1998 (I); pl/co, submo, mo; sa, si, Ac.
- Tephromela atra* (Huds.) Hafellner var. *cypria* (Körb.) Nimis - Koerber 1862 (*Lecanora cypria*); -, ca.
- Thelopsis isiaca* Stizenb. - Litterski 1997 (I); pl/co; Ce.
- Toninia aromatica* (Sm.) A. Massal. - Timdal 1991; -, -.
- Toninia cinereovirens* (Schaer.) A. Massal. - Timdal 1991; -, -.
- Toninia diffracta* (A. Massal.) Zahlbr. - Timdal 1991; -, -.
- Toninia episema* (Nyl.) Timdal - Timdal 1991; -, -.
- Toninia massata* (Tuck.) Herre - Timdal 1991; -, -.
- Toninia plumbina* (Anzi) Hafellner & Timdal - Timdal 1991; -, -.
- Toninia sedifolia* (Scop.) Timdal - Krempelhuber 1865 (III, *Thalloidima vesicularis*), Timdal 1991, Güvenc & Öztürk 1995 (V, *T. caeruleonigricans*), Aptroot 1996 (II), Litterski 1997 (II, III), Litterski & Mayrhofer 1998 (I); pl/co, submo; ca, so.
- Toninia toepfferi* (Stein) Navas - Litterski 1997 (III); submo; so.
- Toninia tristis* (Th. Fr.) Th. Fr. - Vězda 1988c (IV, No. 2262); pl/co; so.
- Toninia tristis* (Th. Fr.) Th. Fr. ssp. *pseudotabacina* Timdal - Timdal 1991; -, -.
- Topelia heterospora* (Zahlbr.) P. M. Jørg. & Vězda - Litterski & Mayrhofer 1998 (I); pl/co; ca.
- Tornabea scutellifera* (With.) J. R. Laundon - Güvenc & Öztürk 1994 (V, *Anaptychia intricata*), Litterski & Mayrhofer 1998 (III); submo, mo; br, Cu.
- Trapelia coarctata* (Sm.) M. Choisy - Aptroot 1996 (III), Litterski & Mayrhofer 1998 (III); pl/co, [mo]; ro, so.
- Usnea ceratina* Ach. - Krempelhuber 1865 (III); [mo, hmo]; -.
- Usnea scabrata* Nyl. - Litterski 1997 (III); mo; Qu.
- Usnea subfloridana* Stirt. - Litterski 1997 (III); mo; Pi.
- Verrucaria calciseda* DC. - Krempelhuber 1865 (II), Aptroot 1996 (II), Litterski 1997 (I, II); pl/co; ca, li, ro.
- Verrucaria fuscella* (Turner) Winch - Krempelhuber 1865 (IV); [pl/co]; -.
- Verrucaria geophila* Zahlbr. - Litterski & Mayrhofer 1998 (I); pl/co; so.
- Verrucaria lecideoides* (A. Massal.) Trevis. - Krempelhuber 1865 (IV); [pl/co]; -.
- Verrucaria macrostoma* Dufour - Aptroot 1996 (II); [pl/co]; ro.
- Verrucaria muralis* Ach. - Aptroot 1996 (II, III), Litterski 1997 (I, III); pl/co, submo, mo; ca, li, ro, sa, si.
- Verrucaria nigrescens* Pers. - Krempelhuber 1865 (II), Litterski 1997 (I, II); pl/co; ca, li, sa.
- Xanthoparmelia conspersa* (Ach.) Hale - Krempelhuber 1865 (III, *Parmelia c.*), Litterski 1997 (III); mo; si.
- Xanthoria aphrodites* Kalb, Poelt & S. Kondratyuk - Kondratyuk & Poelt 1997 (III); mo; Qu.
- Xanthoria calcicola* Oxner - Krempelhuber 1865 (?), *Physcia parietina* var. *aureola*), Güvenc & Öztürk 1994 (V, *X. aureola*), Litterski 1997 (I); pl/co, submo; ca, si.
- Xanthoria candelaria* (L.) Th. Fr. - Litterski 1997 (III), Kondratyuk & Poelt 1997; mo, salp; Pi, Qu.
- Xanthoria ectaneoides* (Nyl.) Zahlbr. - Krempelhuber 1865 (V, *Physcia parietina* var. *ectanea*); [pl/co]; -.
- Xanthoria elegans* (Link) Th. Fr. - Güvenc & Öztürk 1994 (V); pl/co; si.
- Xanthoria* cf. *fulva* (Hoffm.) Poelt & Petutschnig - Kondratyuk & Poelt 1997 (III, poorly developed); mo; Qu.
- Xanthoria mediterranea* Giralt, Nimis & Poelt - Giralt *et al.* 1993 (I), Litterski 1997 (I); pl/co; ca, li.
- Xanthoria parietina* (L.) Th. Fr. - Krempelhuber 1865 (IV, V, *Physcia p.*), Güvenc & Öztürk 1994 (IV, V), Aptroot 1996 (III), Litterski 1997 (I, III), Kondratyuk & Poelt 1997 (III), Litterski & Mayrhofer 1998 (I); pl/co, submo, mo; Be, Ce, Cr, Cu, Ol, Pi, Ps, Qu, Vi, br.
- Xanthoria stiligera* Giralt, Nimis & Poelt - Giralt *et al.* 1993 (I, IV); pl/co; ca.

Taxonomically not clear records

- Placodium fusco-pallens* Kremp. - Krempelhuber 1865 (?), typus in M; -, sa.
- Porpidia stipata* Kremp. - Krempelhuber 1865 (III), typus in M; [hmo]; wo.

Dubious records

- Lecanora subfusca* (L.) Ach. - Güvenc & Öztürk 1995 (V); [pl/co; Ce].
Parmelia olivacea L. - Krempelhuber 1865 (?; III); [pl/co, mo]; -; a species of *Melanelia*.
Physcia parietina L. var. *granulata* Schaer. - Krempelhuber 1865 (IV); [pl/co]; -; probably *Xanthoria stiligera* or *Xanthoria mediterranea*.
Roccella tinctoria Ach. - Krempelhuber 1865 (IV, V); [pl/co]; -; probably *Roccella phycopsis*.
Zeora sordida Pers. var. *pallide-flava* Kremp. - Krempelhuber 1865 (?); -; -; probably *Lecanora rupicola*.

Index of synonyms

- Alectoria jubata* = *Bryoria fuscescens*
Anaptychia intricata = *Tornabea scutellifera*
Anaptychia obscura var. *orbicularis* = *Phaeophyscia orbicularis*
Anaptychia pulchella var. *dubia* = *Physcia dubia*
Anaptychia pulverulenta var. *angustata* = *Physconia muscigena*
Anaptychia pulverulenta var. *grisea* = *Physconia grisea*
Anaptychia pulverulenta var. *musciicola* = *Physconia distorta*
Anaptychia pulverulenta var. *venusta* = *Physconia venusta*
Anaptychia stellaris v. *aipolia* = *Physcia aipolia*
Anaptychia stellaris v. *ambigua* = *Physcia stellaris*
Anaptychia stellaris v. *hispida* = *Physcia adscendens*
Arthopyrenia punctiformis = *Naetrocymbe punctiformis*
Aspicilia mutabilis = *Megaspora verrucosa* var. *mutabilis*
Aspicilia praeradiosa = *Lobothallia praeradiosa*
Aspicilia viridescens = *Aspicilia contorta*
Biatora decipiens = *Psora decipiens*
Biatora triptophylla var. *corallinoides* = *Parmeliella triptophylla*
Biatorina michelettiana = *Solenopsora olivacea* ssp. *olivacea*
Blastenia lallavei = *Caloplaca erythrocarpa*
Buellia punctata = *Amandinea punctata*
Buellia venusta = *Diplotomma venustum*
Calloppisma aurantiacum B. *flavovirescens* = *Caloplaca flavovirescens*
Calloppisma cerinum = *Caloplaca cerina*
Catapyrenium adami-borosi = *Placidium adami-borosi*

- Catapyrenium michelii* = *Placidium michelii*
Catapyrenium semaforonense = *Placidium semaforonense*
Catapyrenium tenellum = *Placidium tenellum*
Cetraria glauca = *Platismatia glauca*
Cladonia chlorophaea see *Cladonia pyxidata*
Cladonia endiviaefolia = *Cladonia convoluta*
Cladonia endivifolia = *Cladonia convoluta*
Cladonia pyxidata c. *pocillum* = *Cladonia pocillum*
Collema albo-ciliatum = *Leptochidium albociliatum*
Collema turgida = *Leptogium turgidum*
Diploschistes calcarius = *Diploschistes candidissima*
Diplotomma epipolium = *Diplotomma alboatrum*
Dirina repanda see *Dirina massiliensis* f. *massiliensis*
Dirina stenhammari = *Dirina massiliensis* f. *sorediata*
Endocarpon miniatum = *Dermatocarpon miniatum*
Endocarpon nodulosum = *Dermatocarpon nodulosum*
Evernia furfuracea = *Pseudevernia furfuracea*
Gyalolechia bracteata = *Fulgensia bracteata*
Lecanactis grumulosa = *Lecanographa grumulosa*
Lecanora atra = *Tephromela atra*
Lecanora chlarona = *Lecanora chlarotera*
Lecanora crassa = *Squamarina cartilaginea*
Lecanora crassa f. *macra* = *Squamarina cartilaginea*
Lecanora cypria = *Tephromela atra* var. *cypria*
Lecanora lentigera = *Squamarina lentigera*
Lecanora murorum y. *calloppisma* = *Caloplaca saxicola*
Lecanora sienae = *Lecanora horiza*
Lecanora subfusca var. *distans* = *Lecanora allophana*
Lecidea conformis = *Immersaria athrocarpa*
Lecidea elata = *Lecanora marginata*
Lecidea enteroleuca var. *rugulosa* = *Lecidella elaeochroma*
Lecidea fumosa = *Lecidea fuscoatra*
Lecidea polycarpa = *Lecidea lapicida* var. *pantherina*
Lecidea sabuletorum var. *euphorea* = *Lecidella euphorea*
Lecidea sabuletorum, *pilularis* = *Mycobilimbia sphaeroides*
Leptogium atrocaeruleum = *Leptogium lichenoides*
Limboria candidissima = *Diploschistes candidissimus*
Parmelia acetabulum = *Pleurosticta acetabulum*
Parmelia ceratophylla var. *tubulosa* = *Hypogymnia*

tubulosa
Parmelia aspera = *Melanelia exasperata*
Parmelia carporrhizans = *Parmelina carporrhizans*
Parmelia conspersa = *Xanthoparmelia conspersa*
Parmelia exasperata = *Melanelia exasperata*
Parmelia luteonotata = *Neofuscelia luteonotata*
Parmelia pulla = *Neofuscelia pulla*
Parmelia quercifolia a *tiliacea* = *Parmelina tiliacea*
Parmelia tiliacea = *Parmelina tiliacea*
Peltigera polydactyla see *Peltigera neckeri*
Physcia callopisma = *Caloplaca flavescens*
Physcia parietina = *Xanthoria parietina*
Physcia parietina var. *aureola* = *Xanthoria calcicola*
Physcia parietina var. *ectanea* = *Xanthoria ectaneoides*
Physma compactum, *Collema chalazanum* = *Lempholemma chalazanum*
Placodium montagnei var. *calcarea* = *Protoparmelia montagnei*
Placodium radiosum = *Aspicilia radiosa*
Placodium saxicolum = *Lecanora muralis*
Placodium saxicolum var. *diffractum* = *Lecanora bolcana*
Psora lurida = *Lecidea lurida*
Psora ostreata = *Hypocenomyce scalaris*
Psoroma crassum β. *caespitosum* = *Squamarina cartilaginea*
Psoroma fulgens = *Fulgensia fulgens*
Psoroma lentigerum = *Squamarina lentigera*
Thalloidima vesicularis = *Toninia sedifolia*
Toninia caeruleonigricans = *Toninia sedifolia*
Urceolaria calcarea = *Aspicilia calcarea*
Urceolaria ocellata = *Diploschistes ocellatus*
Urceolaria scruposa var. *gypsacea*, var. *cretacea* = *Diploschistes gypsaceus*
Urceolaria scruposa var. *bryophila*, var. *arenaria* = *Diploschistes muscorum*
Urceolaria villarsii = *Diploschistes ocellatus*
Urceolaria villarsii var. *macrior* = *Diploschistes muscorum*
Xanthoria aureola = *Xanthoria calcicola*

Acknowledgments

The authors are most grateful to Dr. A. Aptroot, Dr. O. Breuß, Prof. H. Hertel, Dr. V. John, Prof. P. L. Nimis, Dr. H. Sipman, Dr. A. Vězda for their help and information in the preparation of the checklist, and to Prof. T. Ahti, Dr. A. Aptroot, Dr. O. Breuß, Prof. A. R. Burgaz, Dr. Ph. Clerc, Dr. P. Diederich, Prof. J. M. Egea, Prof. T. Esslinger, Dr. M. Giralt, Dr. M. Grube, Prof. P. M. Jørgensen, Dr. T. Lumbsch, Dr. R. Moberg, Dr. P. Navarro-Rosinés, Dr. W. Obermayer, the late Prof. J. Poelt, Dr. H.

Sipman, Dr. L. Tibell and Dr. A. Vězda for their identification of species identified in earlier papers. We thank Dr. B. Coppins and T. Chester, who checked the English translation.

References

- Aptroot A., 1996. *Contribution to the lichen flora of Cyprus*. Cryptog., Bryol. Lichénol., 17: 313-316.
 Breuß O., 1990. *Die Flechtengattung Cata-pyrenium (Verrucariaceae) in Europa*. Stapfia, 23: 1-153.
 Degelius G., 1954. *The lichen genus Collema in Europe*. Symb. Bot. Upsal., 13: 1-499.
 Esslinger T. L., 1977. *A chemosystematic revision of the brown Parmeliae*. J. Hattori Bot. Lab., 42: 1-211.
 Flotow J. v., 1843. *Lichenen des Reisevereins und des Hochstetterschen Herbarii*. Linnaea: 15 - 30. (II. Lichenes Cyprici, p. 18-20).
 Follmann G., 1993. *Vorarbeiten zu einer Monographie der Flechtenfamilie Roccellaceae (Opegraphales, Ascolichenes) IX. Bestandsaufnahme und Verbreitung, Haushalt und Vergesellschaftung, Gefährdung und Schutz der makaronesischen Sippen*. Courier Forschungs-inst. Senckenberg, 159: 175-193.
 Gams H., 1955. *Das Rätsel der Verbreitung von Letharia vulpina*. Svensk Bot. Tidskr., 49: 29-34.
 Geological Survey Department, 1995. *Geological map of Cyprus*. Revised Edition, Ministry of Agriculture, Natural Resources and Environment, Geological Survey Department, Cyprus.
 Giralt M. & Mayrhofer H., 1994. *Four corticolous species of the genus Rinodina (lichenized Ascomycetes, Physciaceae) containing atranorin in southern Europe and adjacent regions*. Nova Hedw., 59: 129-142.
 Giralt M. & Mayrhofer H., 1995. *Some corticolous and lignicolous species of the genus Rinodina (lichenized Ascomycetes, Physciaceae) lacking secondary compounds and vegetative propagules in Southern Europe and adjacent regions*. Bibl. Lichenol., 57: 127-160.
 Giralt M., Nimis P.L. & Poelt J., 1992. *Studien über den Formenkreis von Caloplaca flavorubescens in Europa*. Cryptog., Bryol. Lichénol., 13: 261-273.
 Giralt M., Nimis P.L. & Poelt J., 1993. *Studien über einige Arten der Flechtengattung Xanthoria mit isidiiformen vegetativen Diasporen*. J. Hattori Bot. Lab., 74: 271-285.
 Güvenc S. & Öztürk S., 1994. *Kuzey Kıbrıs Cumhuriyeti Batı Bölgesi Makrolikenleri zerine aksonomik Incelemeler*. Turk. J. Bot., 18: 187-191.
 Güvenc S. & Öztürk S., 1995. *Crustose Lichens of the Western Part of the Turkish Republic of Northern Cyprus*. Turk. J. Bot., 19: 555-558.
 Hafellner J., 1987. *Studien über lichenicole Pilze und Flechten V. Über die Gattung Phacopsis Tul. emend. Körber (Lecanorales)*. Herzogia, 7: 343-352.
 Hermann S., Leuckert C. & Poelt J., 1973. *Zur Kenntnis der Flechtengruppe Lecanora radiosa s. ampliss.* Willdenowia, 7: 9-30.
 Koerber G.W., 1862. *Reliquiae Hochstetterianae*. Abh. Schles. Ges. Vaterl. Cult., Abth. Naturwiss., 1862: 30-34.
 Kondratyuk S. & Poelt J., 1997. *Two new asian Xanthoria species (Teloschistaceae, lichenized Ascomycotina)*. Lichenologist, 29: 173-190.
 Krempelhuber A. v., 1865. *Lichenes*. In F. Unger & T. Kotschy, *Die Insel Cypern*. Braumüller, Wien, p. 158-166.

- Lettau G., 1932 - 1937. *Monographische Bearbeitung einiger Flechtenfamilien*. Feddes Repert. Beih., 69 (1 - 2): 1-250 + Tab.
- Litterski B., 1997. *Lichen observations on the island of Cyprus*. Feddes Repert., 108: 463-473.
- Litterski B. & Mayrhofer H., 1998. *Additional records to the lichen flora of Cyprus*. Cryptog., Bryol. Lichénol., in press
- Lumbsch H.T., 1989. *Die holarktischen Vertreter der Flechtengattung Diploschistes (Thelotremataceae)*. J. Hattori Bot. Lab., 66: 133-196.
- Mayrhofer H., Scheidegger C. & Sheard J.W., 1992. *On the taxonomy of five saxicolous species of the genus Rinodina (lichenized Ascomycetes)*. Nordic J. Bot., 12: 451-459.
- Navarro-Rosinés P. & Hafellner J., 1996. *Lichenostigma elongata spec. nov. (Dothideales), a lichenicolous ascomycete on Lobothallia and Aspicilia species*. Mycotaxon, 57: 211-225.
- Navarro-Rosinés P. & Roux C., 1993. *Caloplaca tavaresiana Nav.-Ros. et Roux sp. nov., espèce nouvelle de lichen du littoral de la région méditerranéenne*. Nova Hedw., 57: 169-177.
- Navarro-Rosinés P. & Roux C., 1994. *Le Caloplacetum tavaresianae Roux et Nav.-Ros. ass. nov., une association lichénique saxicole-calcicole, halophile*. Nova Hedw., 59: 255-264.
- Navarro-Rosinés P. & Roux C., 1995. *Caloplaca navasiana Nav.-Ros. et Roux sp. nov. espèce nouvelle de lichen du littoral Méditerranéen*. Cryptog., Bryol. Lichénol., 16: 89-97.
- Nimis P.L. 1996. *Towards a checklist of Mediterranean lichens*. Bocconea, 6: 5-17
- Øvstedal D.O., 1987. *Noen lav fra toppen av Khionistra, Kypros*. Graphis Scripta, 1: 71.
- Petrak F., Reehinger K.H., Szatala Ö. & Baumgärtner J., 1941. *Schedae ad "Cryptogamas exsiccatas"*. Cent. XXXV. Ann. Naturhist. Mus. Wien, 52: 279-300.
- Roux C., 1991. *Phytogéographie des lichens saxicoles-calcicoles d'Europe méditerranéenne*. Bot. Chron., 10: 163-178.
- Roux C. & Egea J.M., 1992. *Opegraphetum durieui Egea et Roux, ass. nov., une association lichénique saxicole-calcicole, halophile*. Cryptog., Bryol. Lichénol., 13: 105-115.
- Roux C. & Navarro-Rosinés P., 1992. *Caloplaca egeana Roux et Nav.-Ros. sp. nov. kaj Caloplaca veneris Roux et Nav.-Ros. sp. nov., du novaj likenspecioj de la mediteranea marbordo*. Bull. Soc. linn. Provence, 43: 97-103.
- Runemark H., 1956. *Studies in Rhizocarpon. II. Distribution and ecology of the yellow species in Europe*. Opera Bot., 2: 1-150.
- Schade A., 1954. *Über Letharia vulpina (L.) Vain. und ihre Vorkommen in der Alten Welt*. Ber. Bayer. Bot. Ges., 30: 108-126.
- Timdal E., 1991. *A monograph of the genus Toninia (Lecideaceae, Ascomycetes)*. Opera Bot., 110: 1-137.
- Tehler A., 1983. *The genera Dirina and Roccellina (Roccellaceae)*. Opera Bot., 70: 1-86.
- Vězda A., 1987. *Lichenes Selecti Exsiccati*. Fasc. LXXXVII (No. 2151 - 2175). Pruhonice.
- Vězda A., 1988a. *Lichenes Selecti Exsiccati*. Fasc. LXXXVIII (No. 2176 - 2200). Pruhonice.
- Vězda A., 1988b. *Lichenes Selecti Exsiccati*. Fasc. XC (No. 2226 - 2250). Pruhonice.
- Vězda A., 1988c. *Lichenes Selecti Exsiccati*. Fasc. XCI (No. 2251 - 2275). Pruhonice.

Received February 2, 1998

Accepted July 16, 1998

CARACTERIZACIÓN ECOLÓGICA DE LOS ALCORNOCALES IBEROLEVANTINOS (ESTE DE ESPAÑA) BASADA EN BIOINDICADORES LIQUÉNICOS

Simón FOS y Eva BARRENO

Departamento de Biología Vegetal, Botánica, Facultat de Ciències Biològiques, Universitat de València, Dr. Moliner, 50, E 46100 Burjassot, España.

Keywords: Bioclimatology, Bioindicators, Cork-oak, Ecology, Epiphytes, Lichens, *Quercus suber*, Spain.

Abstract: ECOLOGICAL CHARACTERIZATION OF THE IBERO-LEVANTINIAN CORK-OAK FORESTS (EASTERN SPAIN) USING EPIPHYTIC LICHENS AS BIOINDICATORS. This study compares the epiphytic lichen floras of cork-oak forests from eastern Spain. The methodology was designed in such a way as to avoid subjectivity at all stages, from the sampling protocol to data analysis. Field work was carried out in 15 Catalan localities, in mature stands of *Carici-Quercus suberis sigmetum*, and in 9 localities of the Valencian Community, in mature stands of *Asplenio-Quercus suberis sigmetum*. In each locality, ten trees were selected, and the lichens growing on virgin bark from 50 to 200 cm above the ground were identified. All together, 153 species were encountered, with 69 species present in both territories, and 33 found in at least three samplig plots. Multivariate methods of classification and ordination were used to analyze a matrix of stations and species based on presence/absence data. The classification clearly separated two main groups of stations, Catalanian and Valencian, with the exception of two Catalanian localities that were more related to the Valencian ones. This accords well with the climate of the two regions: the Catalanian cork-oak forests are subjected to a strong maritime influence, experiencing frequent fog episodes, whereas the Valencian ones are more continental and drier. The higher incidence of nitrophytic lichens in the Valencian cork-oak forests is probably due to dust accumulation on the trunks. Within the two groups of stations the lichen flora is diversified according to differences in temperature, rainfall, and relative humidity regimes. The results support the assumption that epiphytic lichens can provide important information on the phytoclimatical features of a territory.

Introducción

La sensibilidad de los líquenes y sus comunidades a diferentes parámetros ambientales está plenamente reconocida en la literatura. Esto los convierte en excelentes bioindicadores de factores climáticos relevantes para las plantas (temperatura, la humedad relativa del aire, precipitación, etc.), ya que suministran información de gran valor en la caracterización bioclimática de los territorios (Nimis 1981, 1993; Nimis & Loi 1982, 1984; Sergio *et al.* 1990; Nimis & Tretiach 1995; Fos & Barreno 1994a; Fos 1997; Barreno 1997, 1998). El planteamiento de trabajos comparativos entre áreas climáticamente heterogéneas, manteniendo el mismo sustrato, permite extraer conclusiones sobre las características generales del clima en cada territorio y, simultáneamente, sobre la incidencia de los pará-

metros climáticos que intervienen en la selección de las especies presentes en cada área (James *et al.* 1977; Hawksworth & Hill 1984).

En la Península Ibérica, el alcornoque (*Quercus suber* L.) tiene su óptimo en el cuadrante suroccidental. En el área levantina, forma bosques más o menos extensos en Cataluña (*Carici depressae-Quercetum suberis* (O. Bolós) Rivas-Martínez 1987 = *Quercetum ilicis galloprovinciale suberetosum* Br.-Bl. 1936) y en la Comunidad Valenciana (*Asplenio onpteridis-Quercetum suberis* Costa *et al.* 1985), sometidos a condiciones termo- y ombroclimáticas diferentes (Folch 1981; Costa *et al.* 1985; Costa 1986; Rivas-Martínez 1987; Vilar *et al.* 1989, 1994).

El objetivo de este estudio es comparar la flora

liquénica epífita sobre *Q. suber* en estas dos áreas. La presencia-ausencia de determinados conjuntos de flora y la aplicación de técnicas de análisis multivariante han permitido poner de manifiesto los factores más relevantes en la selección de los táxones en cada territorio y se han obtenido conjuntos de localidades florísticamente homogéneas que delimitan regiones climáticas uniformes.

Área de estudio

El área de estudio abarca los principales núcleos de alcornocales del Levante español (alcornocales iberolevantineos): Cataluña y Comunidad Valenciana (Fig. 1). Los alcornocales catalanes (Fig. 1A) se localizan en dos áreas de desigual importancia, una más extensa que abarca buena parte del sector nororiental de la Cordillera Catalana y otra que ocupa la vertiente meridional del extremo oriental de los Pirineos. Los valencianos (Fig. 1B) tienen una extensión más reducida y quedan limitados a las Sierras de Espadán y Calderona y algunos enclaves del Desierto de las Palmas. Las características climáticas — parámetros climáticos e índices bioclimáticos (Rivas-Martínez *et al.* 1991; Fos 1997) — de cada área se resumen en los diagramas ombroclimáticos de la Fig. 2. En general, los alcornocales catalanes reciben mayor precipitación y muestran un menor período de aridez estival ($P = 685-804$ mm; $Imv = 2.8-4.2$) que los valencianos ($P = 503-637$ mm; $Imv = 4.7-7$). La influencia litoral es más notable en los primeros, tanto por la amortiguación de la oscilación térmica como por la frecuencia de los episodios de nieblas o brumas. Además, en los valencianos, son relativamente frecuentes los episodios de vientos de poniente, muy cálidos y secos, que impiden la formación del régimen de brisas marítimas.

Las localidades muestreadas en cada territorios (Fig. 1) son las siguientes:

Localidades catalanas: 1) AGULLANA, Pla de Pujalts, 300 m, 31TDG8594; 2) DARNÍUS, Serrat del Sentinela, 220 m, 31TDG8590; 3) CAPMANY, Ctra. a Capmany, km 2, 120 m, 31TDG9390; 4) BEGUR, Puig Malaret, 180 m, 31TEG1846; 5) Begur, SA TUNA, 250 m, 31TEG1845; 6) PALAFRUGELL, Cap Roig, 60 m, 31TEG1436; 7) SANT SADURNÍ, Puig dels Cristians, 350 m, 31TDG9239; 8) SANTA CRISTINA d'Aro, Les Taules, 140 m, 31TEG0129; 9) Caldes de Malavella, Riera de RECLÀ, 120 m, 31TDG8627; 10) Caldes de MALAVELLA, Llagostera-Caldes de Malavella, km 5, 100 m, 31 TDG -




Fig. 1 - Áreas potenciales de los alcornocales iberolevantineos (zonas punteadas; tomado de Rivas-Martínez 1987). Situación geográfica de las localidades estudiadas. Retículo UTM de 20 km de lado. Los triángulos indican la localización de las estaciones meteorológicas (ver Fig. 2)

Potential areas of the ibero-levantinian cork-oak forests (dotted areas; according to Rivas-Martínez 1987) and location of the sampling stations. UTM lattice side is 20 km. Triangles: meteorological stations (see Fig. 2).




Fig. 2 - Diagramas ombroclimáticos de algunas localidades catalanas (Palafrugell, triángulo negro, y Castanyet, triángulo blanco en Fig. 1A) y valencianas (Eslida, triángulo negro, y Vall d'Uixó, triángulo blanco en Fig. 1B)(Rivas-Martínez *et al.* 1991).

Climatic diagrams of some Catalanian (Palafrugell, full triangle in Fig. 1A, and Castanyet, hollow triangle) and Valencian (Eslida, full triangle in Fig. 1B, and Vall d'Uixó, hollow triangle) localities, according to Rivas-Martínez *et al.* 1991.

8831; 11) LLORET de mar, Los Pinares, 180 m, 31TDG8520; 12) SANT CELONI, Serra del Solà, Torrent de Olzinelles, 220 m, 31TDG5814; 13) Fogàs de MONCLÚS, La Costa-Mosqueroles, km 6, 580 m, 31TDG5121; 14) BRUNYOLA, Anglès-Sta Coloma de Farnés, km 5, 220 m, 31TDG7138; 15) SANTA COLOMA de Farnés, Serrat del Corb, 260 m, 31TDG6736.

Localidades valenciano-castellonenses: 16) Serra, Serra Calderona, Bco. SARAGUTILLO (I), 550 m, 30SYJ1799; 17) Bco. SARAGUTILLO (II), 580 m, 30SYJ1698; 18) Serra, Serra Calderona, Font del BERRO, 550 m, 30SYJ1699; 19) CHÓVAR, Serra d'Espadà, Bco. del Carbón, 450 m, 30SYK2915; 20) ARTANA, Serra d'Espadà, El Hondo, 280 m, 30SYK3418; 21) Azuebar, Serra d'Espadà, MOSQUERA, 700 m, 30SYK2517; 22) AHÍN, Serra d'Espadà, Los Noguerals, 725 m, 30SYK2718; 23) Algimia de Almonacid, Serra d'Espadà, Bco. de AGUA NEGRA, 800 m, 30SYK2320; 24) Alcudia de Veo, Serra d'Espadà, Emb. de BENITANDÚS, 350 m, 30SYK2724.

Datos y Métodos

Se ha elaborado un catálogo florístico (matriz de presencia-ausencia de especies por localidades) destinado a poner de manifiesto las semejanzas y diferencias florísticas entre los alcornocales iberolevanticos. Para la recolección del material líquénico, en cada punto de muestreo, se han considerado 10 árboles que cumplieran una serie de requisitos: pertenecer a las clases diamétricas medias (diámetro sobre corcho a 1.3 m del suelo entre 90-130 cm), inclinación del tronco inferior a 20° respecto a la perpendicular y, para que fuera posible la recolección de líquenes, altura de extracción del corcho inferior 1.50 m.

Con los inventarios obtenidos, se confeccionó una matriz de datos que sólo incluye las especies recolectadas sobre corcho virgen (bornizo) entre 50 cm y 2 m de altura, en todas las orientaciones. Esto no implica que toda la superficie se encuentre homogéneamente cubierta por la corteza original del árbol, sino que la recolección del material líquénico se ha limitado al bornizo. Esta delimitación del área de muestreo permite evitar la introducción de asimetrías debidas a parámetros diferentes de los que se pretende valorar. Así, quedan excluidas las especies de óptimo terrícola o saxícola silicícola que colonizan las bases de los troncos y las raíces superficiales (*Cladonia* sp. pl., *Diploschistes scrupulo-*

sus, *Lecidella carpathica*, *Parmelia pulla*, *Parmelia loxodes*, *Parmelia verrucigera*, etc.).

También, se han excluido aquellas especies encontradas únicamente en una localidad y sólo representadas por un individuo (*Arthonia galactites*, *A. pinastri*, *Arthothelium ilicinum*, *Bacidia hegetschweileri*, *Collema occultatum*, *Opegrapha vermicellifera*, *Ramalina obtusata*, etc.). En la Tab. 1 sólo se presentan las especies consideradas en los análisis estadísticos, lo que introduce diferencias cuantitativas respecto a los catálogos completos (Fos 1997).

La flora líquénica identificada en cada localidad ha sido analizada utilizando técnicas de análisis multivariantes (clasificación y ordenación), con objeto de obtener grupos de localidades con composición florística semejante. Para la clasificación numérica se ha aplicado la técnica conocida como Análisis Cluster, utilizando la distancia euclídea entre pares de localidades como medida de semejanza y el ligamiento de mínima varianza como algoritmo para la representación del dendrograma. Para la ordenación, se ha utilizado el Análisis de Componentes Principales (ACP), después de la transformación de los datos para análisis recíprocos. Todos los cálculos se han realizado utilizando el programa MULVA para estudio de la vegetación (Wildi & Orlóci 1983).

La caracterización ecológica de las especies está basada, fundamentalmente, en los trabajos de Barkman (1958), Wirth (1980) y Nimis (1993); además se ha tenido en cuenta toda la información recopilada en las numerosas publicaciones líquenológicas que aportan información en este sentido.

Resultados

El estudio de la flora líquénica epífita en los alcornocales iberolevanticos ha permitido obtener una matriz (Tab. 1) de 24 localidades (15 en Cataluña y 9 en la Comunidad Valenciana) y 153 especies (122 en Cataluña y 106 en la Comunidad Valenciana). Ambos territorios comparten 69 táxones, pero de ellos, sólo 33 están presentes en, al menos, tres localidades de muestreo. Estas asimetrías sugieren diferencias florísticas importantes entre los bosques de alcornoques del Este de España.

El análisis cluster (Fig. 3) revela aspectos interesantes sobre la diversidad de la flora líquénica epífita de *Quercus suber* en los alcornocales

Presence of the species in the sampling stations.

ESPECIE	4	5	6	8	7	10	9	15	12	14	1	11	13	17	16	22	21	3	2	20	19	24	23	18	ESPECIE	4	5	6	8	7	10	9	15	12	14	1	11	13	17	16	22	21	3	2	20	19	24	23	18		
Agonimia opuntella	.	.	.	.	.	.	.	.	.	.	1	.	.	1	.	.	.	.	.	.	.	.	.	.	.	Gyalecta derivata	.	1	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	
Amandinea punctata	.	1	1	.	1	1	1	.	.	1	1	.	1	.	1	1	1	1	1	1	.	.	1	1	1	Gyalecta liguriensis	.	.	.	.	1	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.		
Anaptychia ciliaris	.	.	.	.	.	.	.	.	.	.	.	.	1	1	1	1	.	.	.	.	.	.	.	.	Hyperphyscia adglutinata	1	1	1	1	1	1	1	1	1	1	1	.	1	1	1	1	1	1	1	1	1	1	1	1		
Arthonia granosa	.	1	.	.	.	1	.	.	.	.	.	1	.	.	.	.	.	.	.	.	.	.	.	1	.	Hypogymnia physodes	.	.	.	.	1	.	1	.	1	.	1	.	1	1	1	1	1	1	1	1	1	1	1		
Arthonia pruinata	.	.	.	.	1	1	1	.	.	.	.	1	.	.	.	.	.	.	.	.	.	.	.	.	.	Hypogymnia tubulosa	.	.	.	.	.	.	.	.	1	.	.	.	1	.	1	.	1	.	1	.	1	.	1	.	
Arthonia radiata	.	.	.	.	.	.	.	.	.	1	.	.	1	1	1	1	.	1	1	1	1	1	1	1	.	Lecania naegelii	1	1	1	1	1	.	.	1	.	.	1	1	1	1	1	1	1	1	1	1	1	1	1	1	
Arthothelium crozalsianum	.	.	.	.	.	1	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	1	.	.	Lecania cyrtella	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	1	.	.	1	.		
Arthothelium sardoum	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	1	.	.	.	.	.	Lecania fuscella	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	1	.		
Arthothelium spectabile	.	.	.	.	1	1	1	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	Lecanographa lyncea	.	.	.	1	.	1	1	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	
Bacidia rosella	.	.	.	.	1	.	.	.	1	1	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	Lecanora argentata	.	.	.	.	.	.	.	.	.	1	.	.	.	1	.	1	.	1	.	1	.	1	.		
Bactrospora patellarioides	1	1	1	1	.	.	.	.	.	.	.	.	1	.	.	.	1	.	.	.	.	.	.	.	.	Lecanora carpineae	.	.	.	.	.	.	.	.	.	.	1	.	.	.	.	.	.	.	.	.	.	.	.		
Buellia disciformis	.	.	1	.	1	.	.	.	1	.	.	.	.	.	.	1	.	.	.	.	.	.	1	.	.	Lecanora circumborealis	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	1	.	1	.	
Buellia ericaria	.	.	1	1	.	.	.	.	.	.	.	.	1	.	.	.	.	.	.	.	.	.	.	.	.	Lecanora conizaeoides	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	1	1	.	1	.
Buellia erubescens	.	.	.	.	.	.	.	.	1	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	Lecanora chlarotera	.	.	.	1	.	.	.	.	.	.	.	.	.	.	1	.	1	.	.	.	.	.	.	.	
Buellia schaeeri	.	.	.	.	1	1	.	.	.	.	.	.	1	.	.	.	.	.	.	.	.	.	.	.	.	Lecanora expallens	.	.	.	.	1	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	
Caloplaca aegatica	.	1	1	1	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	Lecanora horiza	.	.	.	1	1	.	.	.	.	.	.	1	1	1	1	1	1	.	1	.	.	.	1	.	
Caloplaca cerina	.	.	1	.	.	.	.	.	.	.	.	.	1	.	.	.	1	.	.	.	.	.	.	.	.	Lecanora livido-cinerea	.	1	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	
Caloplaca cerinella	.	.	.	.	.	.	.	.	.	.	.	.	1	.	.	.	1	1	1	.	.	1	1	1	.	Lecanora pallida	.	.	.	1	.	1	1	1	1	1	1	.	.	.	.	.	.	.	.	.	.	.	.		
Caloplaca cerineloides	.	.	.	.	.	.	.	.	.	.	.	.	1	.	.	.	.	.	.	.	.	.	.	.	.	Lecanora pulcaris	1	1	1	1	1	.	1	.	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	
Caloplaca ferruginea	.	.	1	.	1	.	.	1	.	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	.	Lecanora rubicunda	.	1	1	1	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	
Caloplaca flavorubescens	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	1	.	.	.	.	.	.	.	.	Lecanora strobilina	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	
Caloplaca herbidella	.	.	.	.	.	.	.	.	.	.	.	.	1	.	.	.	.	.	.	.	.	.	.	.	.	Lecanora sylvestris	.	.	.	1	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	
Caloplaca holocarpa	.	.	1	.	.	.	.	.	.	.	.	.	.	.	.	.	.	1	.	.	.	.	1	1	.	Lecanora symmicta	.	.	.	1	1	.	.	.	1	.	1	1	.	.	.	1	1	.	1	1	.	1	.	1	.
Caloplaca hungarica	.	.	.	.	.	.	.	.	.	.	.	.	1	.	.	.	.	.	.	.	.	.	.	.	.	Lecidella achrostotera	.	.	.	1	1	.	1	1	.	1	.	1	1	1	1	1	.	1	1	.	1	1	.	1	.
Caloplaca lucifuga	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	1	.	.	.	.	.	.	1	1	.	Lecidella elaeochroma	.	.	.	.	.	.	.	.	.	1	.	.	.	.	.	.	.	1	.	.	.	1	.	.	
Caloplaca obscurella	.	1	.	.	1	1	.	.	.	.	.	.	1	.	1	1	1	1	1	1	1	1	1	1	.	Lecidella euphorea	.	.	.	.	.	.	.	.	.	1	1	1	.	1	1	.	1	1	.	1	1	.	1	1	
Caloplaca pollinii	.	.	.	1	1	1	.	.	1	1	.	1	1	1	1	1	1	1	1	1	1	1	1	1	.	Leptogium cyanescens	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.
Caloplaca sarcopisoides	.	.	.	1	.	.	.	.	.	.	.	.	.	.	.	1	.	.	.	.	.	.	.	.	.	Leptogium teretiusculum	.	1	.	.	1	.	.	.	.	1	.	.	1	1	1	1	1	.	.	.	1	1	1	1	1
Candelaria concolor	.	.	1	.	.	1	.	.	1	1	.	1	.	.	.	.	.	1	1	1	1	1	1	1	.	Maronea constans	.	.	.	.	.	.	.	1	.	.	1	.	.	.	.	.	.	.	.	.	.	.	.	.	
Candelariella vitellina	.	.	1	.	.	.	.	.	.	.	.	.	1	.	1	1	1	1	1	1	1	1	1	1	.	Ochrolechia pallescens	1	.	.	.	.	.	.	.	.	1	.	.	1	.	.	.	.	.	.	.	.	.	.	.	
Candelariella xanthostigma	.	.	.	.	.	.	.	.	.	.	.	.	1	.	.	.	.	.	.	.	.	1	.	.	.	Ochrolechia turneri	.	.	.	.	.	.	.	.	.	1	1	.	1	.	.	.	.	.	.	.	.	1	.		
Catillaria chalybeia	.	.	.	.	.	.	.	.	1	.	.	.	1	1	1	1	1	1	1	1	1	1	1	1	.	Opegrapha atra	1	1	1	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.
Catillaria nigroclavata	.	1	.	1	.	.	.	.	1	.	.	1	1	1	1	1	1	1	1	1	1	1	1	1	.	Opegrapha niveoatra	1	.	.	1	1	1	1	1	.	1	.	1	1	1	1	1	1	1	1	1	1	1	1	1	
Collema furfuraceum	.	.	.	.	.	.	.	.	.	.	.	.	1	1	.	1	.	.	.	.	1	.	1	.	.	Opegrapha varia	1	1	1	.	1	1	1	.	.	.	.	.	.	.	1	1	.	.	.	.	.	.	.	.	
Collema subnigrescens	.	.	.	.	.	.	.	.	.	.	.	.	1	.	.	.	1	.	.	.	.	.	.	.	.	Opegrapha vulgata	1	1	1	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.
Chrysothrix candelaris	1	1	1	1	1	1	1	1	1	1	1	1	.	.	.	1	1	1	1	1	1	1	1	1	.	Parmelia caperata	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	
Dimerella pineti	.	.	.	.	.	1	.	.	.	.	.	1	.	.	.	.	.	.	.	.	.	.	.	.	.	Parmelia chinense	1	1	.	.	1	.	.	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	
Diploicia canescens	1	1	1	1	.	1	.	.	.	.	1	.	.	.	.	.	.	.	1	.	.	.	.	.	.	Parmelia crozalsiana	.	.	.	.	.	.	.	.	.	1	.	.	.	.	1	.	.	.	.	.	.	.	.	.	
Eopyprenula leucoplaea	.	.	.	.	.	.	.	.	.	.	.	.	1	1	.	.	.	.	.	.	.	.	1	.	.	Parmelia glabra	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	1	.	.	.	.	.	.	.	.	
Evernia prunastri	.	.	.	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	.	Parmelia glabratula	.	.	.	1	.	.	1	.	.	1	.	1	1	1	1	.	.	.	1	.	.	1	.	.	.

ESPECIE	4	5	6	8	7	10	9	15	12	14	1	11	13	17	16	22	21	3	2	20	19	24	23	18	ESPECIE	4	5	6	8	7	10	9	15	12	14	1	11	13	17	16	22	21	3	2	20	19	24	23	18				
Parmelia hypoleucina	1		1	1								1														Physconia servitii																								1			
Parmelia laciniatula																		1								Physconia venusta																											
Parmelia quercina					1													1	1						1	Polyblastiopsis subericola	1		1		1	1	1			1	1		1	1													
Parmelia reticulata	1	1	1	1	1	1	1	1	1	1	1	1	1						1	1	1					Porina aenea	1	1			1		1	1																1			
Parmelia soredians	1	1	1	1	1	1	1	1	1	1	1	1	1					1	1	1	1	1	1	1	1	Pseudevernia furfuracea				1							1																
Parmelia subaurifera								1	1	1	1	1	1	1					1	1	1				1	Pyrenula chlorospila	1																										
Parmelia subrudecta	1	1		1	1			1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	Pyrrhospora quernea					1							1															
Parmelia sulcata					1	1		1		1	1	1	1	1					1	1	1				1	Ramalina canariensis				1																					1		
Parmelia tiliacea					1			1	1		1		1	1	1	1	1	1	1	1	1	1	1	1	1	Ramalina farinacea				1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1		
Pertusaria albescens					1							1	1					1	1	1	1				1	Ramalina fastigiata	1		1			1						1	1	1	1			1	1	1	1	1	1	1	1	1	
Pertusaria amara	1	1		1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1					1	Ramalina fraxinea																										1	
Pertusaria flavida						1						1						1	1	1	1					Ramalina pusilla	1																										
Pertusaria hemisphaerica					1				1	1								1	1	1					1	Rinodina capensis		1									1		1	1	1									1	1	1	
Pertusaria heterochroa	1	1	1	1		1	1				1	1														Rinodina euskadiensis																											
Pertusaria hymenea																										Rinodina plana																											
Pertusaria leucostoma				1	1														1	1	1	1			1	Rinodina aff. poeltiana								1			1															1	
Pertusaria pertusa																										Schismatomma decolorans	1	1	1	1	1		1	1	1	1	1		1														
Pertusaria pustulata	1	1	1	1	1	1	1				1	1														Scoliciosporum chlorococcum											1		1														
Phaeophyscia cernohorskyi																										Scoliciosporum sarothamni		1																									
Phaeophyscia cloantha					1																					Scoliciosporum umbrinum																											
Phaeophyscia hirsuta					1																					Teloschistes chrysophthalmus		1		1										1	1	1	1	1	1			1	1	1	1	1	
Phaeophyscia orbicularis																										Thelenella modesta				1												1											
Phlyctis agelaea					1																					Trapeliopsis flexuosa					1							1															
Physcia adscendens	1	1		1	1	1	1	1	1	1	1	1														Trapeliopsis granulosa																											
Physcia aipolia																										Usnea ceratina					1																						
Physcia caesia																										Usnea cornuta																											
Physcia clementei					1																					Usnea dasaea						1																					
Physcia dubia																										Usnea esperantiana							1																				
Physcia leptalea				1																						Usnea glabrata																											
Physcia stellaris																										Usnea hirta																											
Physcia tenella	1		1	1	1	1		1		1	1	1	1	1	1	1	1	1	1	1	1	1	1	1		Usnea mutabilis																											
Physcia tribacia																										Usnea rubicunda		1		1	1	1	1	1	1	1	1		1														
Physcia tribacioides																										Usnea subscabrosa					1																						
Physcia vitti																										Usnea wasmuthii																											




Fig. 3 - Clasificación de las localidades basada en los datos de la Tab. 1. La medida de semejanza es la Distancia Euclídea. La numeración de las localidades se indica en la figura 1.

Classification of sampling stations based on the data of Tab. 1. Resemblance measure: Euclidean Distance. Numbers as in Fig. 1.

levantinos. Esta diversidad parece responder a las condiciones bioclimáticas de cada territorio, al área biogeográfica y a la serie de vegetación. Las localidades se agrupan en dos conjuntos bien diferenciados (98.1 de distancia euclídea) que se corresponden con cada territorio estudiado, a excepción de las localidades muestreadas en la comarca del Alto Ampurdán gerundense, Darnius y Capmany, que se agrupan con las valenciano-castellonenses.

Los alcornocales catalanes (cluster 1) se subdividen inicialmente en dos grupos (32.4 de distancia euclídea): el cluster 1a (CAT.LIT. en Tab. 2) agrupa las estaciones litorales de Begur (Loc. 4), Sa Tuna (Loc. 5.), Palafrugell (Loc. 6) y Santa Cristina d'Aro (Loc. 8). Las diferencias responden de manera clara al grado de influencia marítima que soportan. En estas localidades hay una mayor

diversidad, frecuencia y cobertura de los líquenes con *Trentepohlia* como fotobionte (*Arthonia granosa*, *A. radiata*, *A. pruinata*, *Opegrapha atra*, *O. niveoatra*, *O. varia*, *O. vermicellifera*, *O. vulgata*, *Pyrenula chlorospila*, etc.), indicadoras de una notable incidencia del litoral. A todo este conjunto de especies, se unen otras de carácter termófilo y/o de óptimo litoral (Tab. 2) como *Caloplaca aegatica*, *Diploicia canescens*, *Lecanora lividocinerea*, *L. rubicunda*, *L. sylvestris*, *Parmelia hypoleucina*, *Ramalina pusilla*, *R. canariensis*, que determinan la separación de este grupo y revelan las condiciones en las que se desarrollan estos alcornocales.

Las restantes estaciones se separan igualmente en dos grupos (24.2 de distancia euclídea). La distinción más clara es la de Monclús (Loc. 13): la localidad más alejada de la costa y situada a mayor altitud (cluster 1c; CAT.CONT.). La influencia

litoral se deja sentir bastantes kilómetros hacia el interior, pero se ve atenuada con el desplazamiento perpendicular a la línea de costa, siendo otros los parámetros que adquieren mayor importancia en la selección de las especies. Algunas de ellas todavía revelan su incidencia (*Bactrospora patellarioides*, *Caloplaca pollinii*, *Opegrapha niveoatra*, *Parmelia reticulata*, *Schismatomma decolorans*, etc.), pero ésta resulta insuficiente para anular la mayor continentalidad que impone su localización. Este contraste térmico es aprovechado por especies como *Hypogymnia tubulosa*, *Parmelia quercina*, *P. sulcata*, *Pseudevernia furfuracea* y *Usnea hirta*, indicadoras de una mayor amplitud térmica. La convergencia de diversas influencias, unido al buen estado de conservación del bosque son, probablemente, las que determinan que esta localidad tenga la mayor riqueza florística del territorio (54 especies; Fig. 4).

En el otro grupo (cluster 1b) se reúnen, por un lado, las situadas entre la línea de costa y los territorios potenciales de la serie del quejigo africano (*Carici-Quercus canariensis sigmetum*) — Sant Sadurní (Loc. 7), Reclà (Loc. 9) y Malavella (Loc. 10) — las cuales, según demuestran los datos climáticos, los líquenes (Fos 1997) y la flora vascular que acompaña al alcornoque (Vilar *et al.* 1989; 1994), gozan de un ombroclima más húmedo. El otro grupo está formado por un conjunto más heterogéneo desde la óptica de su situación geográfica, pero que, evidentemente, comparte unas semejanzas florísticas resultantes de su proximidad ecológica. En éste grupo se aglutinan, por un lado, las estaciones que se localizan también próximas a estos territorios, pero más hacia el interior (Locs. 14. Brunyola y 15. Santa Coloma) y San Celoni (Loc. 12). Por otro lado, se aproximan las localidades de Agullana (Loc. 1) y Lloret (Loc. 11). En general, todas ellas corresponden a bosques maduros, poco alterados por las intervenciones selvícolas, y húmedos, aunque se observa un incremento de las especies fotófilas y una reducción de las termófilas y/o de tendencia litoral, que mantenían valores elevados en las localidades más húmedas (Tab. 2). Estas dos características ambientales parecen ser las que tienen mayor significación para explicar los conjuntos que se obtienen aunque, el grupo formado por Lloret y Agullana requeriría de estudios más profundos con objeto de poder descartar si esta afinidad se debe más a sus diferencias respecto a las restantes localidades catalanas que a las afinidades existentes entre ellas. En este sentido hay que recordar dos cuestiones: Lloret (Loc. 11) está situada




Fig. 4 - Riqueza de especies en las localidades estudiadas.
Species number in the sampling stations.

en una posición intermedia entre las localidades litorales y las más húmedas (Fig. 1A) y, sin embargo, prácticamente carece de elementos indicadores de termicidad o litoralidad. Por su parte, Agullana (Loc. 1) es la única localidad alto ampurdanesa que se mantiene próxima a las localidades catalanas. Probablemente, la mejor estructuración de este bosque frente a los aclarados y desprovistos del matorral que observamos en Darnús (Loc. 2) y Capmany (Loc. 3), sea la principal responsable de esta proximidad florística y de las diferencias con estas últimas. Lamentablemente, no existen datos climáticos de esta zona, que serían necesarios para poder establecer si las condiciones de humedad son tan limitantes y, al mismo tiempo, explicar las repercusiones que parecen tener las intervenciones realizadas en el bosque.

Las localidades valenciano-castellonenses se agrupan en el cluster 2 (Fig. 3), separadas en dos subgrupos bastante próximos (32.4 de distancia euclídea). El grupo 2a (VAL.STR. en Tab. 2) reúne las localidades Saraguttillo (Locs. 16 y 17), Mosquera (Loc. 21) y Ahín (Loc. 22) que, por su estado de conservación, representan muestras excelentes de estos alcornoques (Costa *et al.* 1985; Costa 1986). Estos bosques bien estructurados ofrecen un ambiente más húmedo (Barkman, 1958) que favorece la entrada de todo un conjunto de especies (*Anaptychia ciliaris*, *Collema subnigrescens*, *Hypogymnia physodes*, *Parmelia crozalsiana*, *P. glabra*, *P. glabratula*, *Pertusaria flavida*, *P. hemisphaerica*, *P. hymenea*, *P. pertusa*, *Physconia venusta*) que faltan en las restantes estaciones, y determinan las mayores riquezas florísticas registradas en estos territorios (Fig. 4; Tab. 1). En estas localidades es donde se concentran las especies higrofilas (Tab. 2); ocasionalmente, pueden aparecer en otras, pero, en estos casos, siempre refugiadas en

Tab. 2 - Preferencias ecológicas de la flora líquénica epífita en los territorios estudiados, distinguiendo los grupos establecidos en el dendrograma de la Fig. 3. Xer = Xerófilo; Hig = Higrófilo; Esc = Esciófilo; Fot = Fotófito; Nitr = Nitrófilo; Anit = Anitrófilo; Oce = Oceánico; Suboc = Suboceánico; Cont = Continental; Term = Termófilo; Lit = Litoral. Caracterización ecológica basada en Barkman (1958), Wirth (1980), Nimis (1993), etc.

Ecological preferences of the epiphytic lichen flora in the two survey areas. Independent groups formed according to the dendrogram of Fig. 3 are distinguished. Xer = Xerophytic; Hig = Higrrophytic; Esc = Skiophytic; Fot = Photophytic; Nitr = Nitrophytic; Anit = Anitrophytic; Oce = Oceanic; Suboc = Suboceanic; Cont = Continental; Term = Thermophytic; Lit = Maritime. Data from Barkman (1958), Wirth (1980), Nimis (1993), etc. (further explanation in the main text).

	Xer	Hig	Esc	Fot	Nitr	Anit	Oce	Suboc	Cont	Term	Lit
ALC. CATALANES	29	42	20	53	44	18	15	22	9	17	21
CAT.LIT. (Locs. 4, 5, 6, 8)	13	25	11	28	24	5	6	14	1	15	17
CAT.HD. (Locs. 7, 9, 10)	12	25	11	29	25	11	8	15	4	12	11
CAT.MED. (Locs. 1, 11, 12, 14, 15)	11	24	13	34	22	11	7	12	4	9	6
CAT.VAL. (Locs. 2, 3)	16	7	1	31	31	1	1	7	0	7	2
CAT.CONT. (Loc. 13)	15	16	2	33	26	6	2	7	4	5	3
ALC. VALENCIANOS	35	27	9	63	60	9	5	16	7	14	12
VAL.STR. (Locs. 16, 17, 21, 22)	26	26	9	53	47	9	3	12	5	10	8
VAL.TER. (Locs. 19, 20)	19	9	1	35	39	2	1	6	1	6	3
VAL.MED. (Locs. 23, 24)	20	11	3	38	38	2	2	7	4	7	4
VAL.XER. (Loc. 18)	19	12	3	35	32	1	1	6	2	8	4

microhábitats esciófilos y húmedos.

Las restantes localidades se han visto más afectadas como consecuencia de los incendios y por la deforestación. El cluster 2c (VAL.XER.) separa, primeramente, la localidad Font del Berro (Loc. 18), que junto con las estudiadas en el Barranco Saragutillo (Locs. 16 y 17), constituyen los núcleos de alcornocal termomediterráneo seco estudiados en la Sierra Calderona. Sin embargo, las diferencias estructurales y microambientales son suficientes para separarla florísticamente de éstas y de las demás localidades que forman este grupo. Las modificaciones introducidas por la alteración severa del bosque han provocado un empobrecimiento de la riqueza florística, con una disminución de los táxones de preferencias higrófilas, esciófilas y/o anitrófilas. Esta tendencia favorece el incremento de cobertura de las especies más tolerantes, sin que se observen asimetrías en las proporciones de especies favorecidas por las condiciones dominantes. Un hecho que no se refleja en los inventarios de

presencia-ausencia, pero que marca una clara diferencia con las restantes estaciones de este área, es la cobertura mínima del estrato foliáceo. En Font del Berro (Loc. 18), los especies crustáceas son las que muestran coberturas más elevadas, pudiendo llegar a colonizar la mayor parte de la superficie. Sobre este estrato primario aparecen, de forma muy dispersa o concentrados a las posiciones más húmedas, los talos de las especies foliáceas que, por lo general, muestran un escaso desarrollo. Además, las especies fruticulosas están representadas por talos de 1-2 cm y no siempre han podido ser identificadas. Las limitaciones encontradas para las especies con mayores demandas hídricas están poniendo de manifiesto las condiciones de xericidad que afectan a este enclave y, al mismo tiempo, las razones que explican su independencia florística dentro del conjunto de localidades valenciano-castellonenses más afectadas por las intervenciones humanas.

El cluster 2c se completa con otro grupo que

sustitución de conjuntos florísticos que son desplazados por especies banales de amplia valencia ecológica (Boreham 1993; Canters *et al.* 1991; Hyvärinen *et al.* 1992; Kuusinen 1994a,b, 1996) que, aprovechando la nueva situación, avanzan desde la orla del bosque o desde sus posiciones de refugio para ocupar el espacio que ha quedado disponible. Las observaciones realizadas en los alcornocales iberolevantineos confirman este enunciado: en ambos territorios las estaciones más ricas corresponden a los bosques mejor conservados, aunque no siempre la menor diversidad coincide con la mayor alteración. Las características topográficas del bornizo, con sus profundas grietas, cavidades e irregularidades diversas, ofrece ambientes adecuados para que determinados elementos, algo más exigentes, puedan encontrar condiciones aptas, cuanto menos, para sobrevivir. Esta correspondencia entre elevada riqueza florística y estado de conservación del bosque, está generalizada para todos los alcornocales ibéricos (Fos 1997) y es un buen indicador de cómo estos ambientes favorecen comunidades epífitas diversificadas y bien estructuradas en sus tres estratos: crustáceo, foliáceo y fruticuloso (Barreno *et al.* 1988; Crespo *et al.* 1983; Atienza 1990; Etayo & Gómez-Bolea 1992; Fos & Barreno 1994a; Holien 1996; Kuusinen 1994a,b; Rose 1976).

En resumen, las localidades catalanas corresponden a bosques térmicos y húmedos, afectados por una influencia marítima u oceánica que ejerce un efecto atemperador de la amplitud térmica y, simultáneamente, representa un aporte suplementario de agua, incluso durante los meses de sequía estival. Estas condiciones se traducen en una mayor riqueza en especies higrófilas (59%), suboceánicas (20%) y oceánicas (14%) (Tab. 2). A este respecto, sorprende que con precipitaciones totales anuales entre 685 y 804 mm (Fig. 2), *Leptogium teretiusculum* sea el único cianolíquen encontrado en los bosques catalanes.

Los alcornocales valenciano-castellonenses presentan una flora líquénica epífita algo más continental y xérica, soportan un prolongado período de aridez estival que coincide con las máximas temperaturas y el aporte de agua por criptoprecipitaciones parece ser claramente inferior. Estas condiciones marcan diferencias con los bosques catalanes: disminuyen las especies higrófilas (44%), así como las de carácter suboceánico (15%) y oceánico (5%). Por otra parte, la menor cuantía y la distribución anual de las precipitaciones (Fig. 2) favorecen los procesos de

acumulación de polvo y nutrientes. Con carácter general, y como corresponde a territorios mediterráneos, con largos períodos de depósito, la mayoría de las especies tienen preferencia por los ambientes ricos en sustancias nitrogenadas (87%).

Esta caracterización bioclimática contrasta con la información suministrada por otros estudios de flora y vegetación en estas mismas sierras. La flora vascular (Costa *et al.* 1985; Crespo-Villalva 1989; García-Fayos 1982, 1991) y la flora líquénica saxícola silicícola (Calatayud 1991; Calatayud & Barreno 1994) evidencian un carácter térmico y litoral que no encuentra correspondencia en la flora líquénica epífita. En la Tab. 2 puede comprobarse la escasa representación de especies indicadoras de estas condiciones, especialmente fuera del subconjunto formado por las localidades mejor conservadas (VAL.STR). Es evidente que estas asimetrías son el resultado de la mayor o menor incidencia de las condiciones dominantes en los diferentes tipos de hábitats. Los epífitos se encuentran más expuestos a las variaciones del clima mientras que los saxícolas están más favorecidos por la atenuación de la oscilación térmica y y la conservación de la humedad (Barkman 1958; Jörgensen, 1978). Además, estas zonas están influenciadas también por los vientos de poniente, secos y cálidos, que inhiben el régimen de brisas marinas, creándose un gradiente térmico y de humedad entre la costa y unos pocos centenares de metros hacia el interior (Mediavilla 1994).

La caracterización bioclimática de estos territorios, basada en la flora líquénica epífita, también coincide con los resultados obtenidos mediante la cuantificación del crecimiento radial del corcho de reproducción (Fos & Barreno 1994a,b; Fos 1997). En conclusión, el gradiente oceanidad-continentalidad parece ser el que selecciona principalmente los conjuntos de flora líquénica epífita en territorios mediterráneos.

Referencias

- Atienza V., 1990. *Flora y vegetación líquénica epífita de las Comarcas de Els Ports y Baix Maestrat (Castellón)*. Tesis Doctoral. Universitat de València.
- Barkman J.J., 1958. *Phytosociology and ecology of Cryptogamic Epiphytes*. Van Gorcum, Assen, 644 pp.
- Barreno E., 1997. *Hongos simbiosis: Líquenes, micoficobiosis y micorrizas*. In: Izco J., Barreno E., Brugués M., Costa M., Devesa J., Fernández F., Gallardo T., Llimona X., Salvo E., Talavera S. & Valdés B., *Botánica*. McGraw-Hill-Interamericana, Madrid, 309-340.
- Barreno E., 1998. *Líquenes en los castaños*. En: Berrocal M., Gallardo J.F. & Cardeñoso J.M., *El Castaño*. Mundi-Prensa,

- Barcelona, 59-96.
- Barreno E., Sanz M.J., Atienza V. & Muñoz A., 1988. *Biogeografía y ecología comparadas de líquenes epífitos de alcornocales ibéricos y sardos*. Actes del Simposi Internacional de Botànica "Pius Font i Quer", Vol. I, Criptogàmia: 179-185.
- Boreham S., 1993. *Changes in the lichen flora on birch Betula pendula in Northern Epping Forest*. London Naturalist, 72: 25-30.
- Burgaz A.R., Fuertes E. & Escudero A., 1994a. *Ecology of cryptogamic epiphytes and their communities in deciduous forests in the mediterranean Spain*. Vegetatio, 112: 73-86.
- Burgaz A.R., Fuertes E. & Escudero A., 1994b. *Climax epiphytic communities in Mediterranean Spain*. Botanical Journal of the Linnean Society, 115: 35-47.
- Calatayud V., 1991. *Líquenes saxícolas de las rocas silíceas de la Sierra de Espadán*. Tesis, Universitat de València.
- Calatayud V. & Barreno E., 1994. *Contribution to the lichen floristics of Eastern Spain. I. Siliceous lichens and their lichenicolous fungi of Serra d'Espadà (Castelló)*. Crypt., Bryol. Lichénol., 15: 23-41.
- Canters K.J., Schöller H., Ott S. & Jahns H.M., 1991. *Microclimatic influence on lichen distribution and community development*. Lichenologist, 23: 237-252.
- Costa M., 1986. *La vegetación en el País Valenciano*. Cultura Universitaria Popular 5, Universitat de València, 246 pp.
- Costa M., Peris J.B., Figuerola R. & Stübing G., 1985. *Los alcornocales valencianos*. Doc. Phytosoc., 9: 301-318.
- Crespo A., Barreno E. & Sancho L.G., 1983. *Esbozo de la vegetación líquénica de algunas localidades de los Valles del Tambre y Ulla (La Coruña, España)*. Trab. Compost. Biol., 10: 97-108.
- Crespo-Villalba M.B., 1989. *Contribución al estudio florístico, fitosociológico y fitogeográfico de la Sierra Calderona (Valencia Castellón)*. Tesis Doctoral. Universidad de Valencia.
- Etayo J. & Gómez-Bolea A., 1992. *Estabilidad ecológica por medio de bioindicadores líquénicos en robledales de los Pirineos atlánticos*. Folia Bot. Misc., 8: 61-75.
- Folch R., 1981. *La vegetació dels Països Catalans*. Mem. Inst. Catalana Hist. Nat., 10: 3-513.
- Fos S., 1997. *Líquenes epífitos de los alcornocales ibéricos. Correlaciones bioclimáticas, anatómicas y densimétricas con el corcho de reproducción*. Tesis Doctoral. Universitat de València.
- Fos S. & Barreno E., 1994a. *Epiphytic lichens on Quercus suber and their relation to the quality of cork*. Crypt. Bot., 4: 156-165.
- Fos S. & Barreno E., 1994b. *Crecimiento radial del corcho de reproducción en los alcornocales catalanes y valencianos*. Scientia gerund., 20: 5-15.
- García-Fayos P., 1982. *Estudio sobre la vegetación de los alcornocales de la Sierra Calderona*. Tesis, Universidad de Valencia.
- García-Fayos P., 1991. *La vegetación silicícola de la Sierra Calderona (Comunidad Valenciana)*. Lazaroa, 12: 317-332.
- Giralt M., 1996. *Líquens epífitos i contaminació atmosfèrica a la plana i les serralades litorals tarragonines*. Institut d'Estudis Catalans. ASC 113, Barcelona, 525 pp.
- Hawksworth D.L. & Hill D.J., 1984. *The Lichen-Forming fungi*. Chapman & Hall, New York, 158 pp.
- Holien H., 1996. *Influence of site and stand factors on the distribution of crustose lichens of the Caliciales in a suboceanic spruce forest area in central Norway*. Lichenologist, 28: 315-330.
- Hyvärinen M., Halonen P. & Kauppi M., 1992. *Influence of stand age and structure on the epiphytic lichen vegetation in the middle-boreal forest of Finland*. Lichenologist, 24: 165-180.
- James P.W., Hawksworth D.L. & Rose F., 1977. *Lichen communities in the British Isles: a preliminary conspectus*. In: Seaward, M.R.D. (ed.). *Lichen Ecology*. Academic Press, London, 295-413.
- Jørgensen P.M., 1978. *The lichen family Panariaceae in Europe*. Opera Bot., 45: 1-124.
- Kuusinen M., 1994a. *Epiphytic lichen diversity on Populus tremula in old-growth and managed forest of southern and middle boreal Finland*. Ann. Bot. Fennici, 31: 77-92.
- Kuusinen M., 1994b. *Epiphytic lichen diversity on Salix caprea in old-growth southern and middle boreal forest of Finland*. Ann. Bot. Fennici, 31: 245-260.
- Kuusinen M., 1996. *Epiphyte flora and diversity on basal trunks of six old-growth forest tree species in southern and middle boreal Finland*. Lichenologist, 28: 443-463.
- Mediavilla J., 1994. *Ponientes y olas de calor*. In: Pérez Cueva A.J. (Coord.), *Atlas climático de la Comunidad Valenciana*. Conselleria de d'Obres Públiques, Urbanisme i Transports, Generalitat Valenciana, Valencia, 62-63.
- Nimis P.L., 1981. *Epiphytic lichen vegetation in the Lumiei-Valley (Carnian Alps)*. Gortania, 3: 123-142.
- Nimis P.L., 1993. *The Lichens of Italy. An annotated catalogue*. Museo Regionale di Scienze Naturali, Torino, 897 pp.
- Nimis P.L. & Loi E., 1982. *I Licheni epifiti della provincia di Trieste*. Gortania, 3: 101-122.
- Nimis P.L. & Loi E., 1984. *I Licheni della dolina di Percedol (Carso Triestino)*. Studio fitogeografico. Atti Mus. Civ. Stor. Nat. Trieste, 36: 1-12.
- Nimis P.L. & Poelt J., 1987. *The lichens and lichenicolous fungi of Sardinia (Italy)*. Studia Geobot., 7, suppl. 1: 1-269.
- Nimis P.L. & Tretiach M., 1995. *The lichens of Italy. A phytoclimatical outline*. Crypt. Bot., 5: 199-208.
- Rivas-Martínez S., 1987. *Memoria del mapa de series de vegetación 1:400.000*. Ministerio de Agricultura, Pesca y Alimentación, Serie técnica, Madrid, 268 pp.
- Rivas-Martínez S., Bascones J.C., Díaz T.E., Fernández F. & Loidi J., 1991. *Vegetación del Pirineo Occidental y Navarra*. Itin. Geobot., 5: 5-456.
- Rose F., 1976. *Lichenological indicator of age and environmental continuity in woodland*. In: Brown D.H.; Hawksworth D.L. & Bailey R.H. (eds.), *Lichenology: Progress & Problems*. Academic Press, London & New York, 279-307.
- Sergio C., Sim-Sim M. & Santos-Silva C., 1990. *Briófitos epífitos como indicadores de dominios bioclimáticos em Portugal. Tratamento estadístico de áreas seleccionadas*. Anales Jard. Bot. Madrid, 46: 457-467.
- Vilar L., Polo L. & Domínguez Planella A., 1989. *Los alcornocales de la provincia de Girona*. Scientia gerund., 15: 143-151.
- Vilar L., Domínguez-Planella A. & Polo L., 1994. *Aplicació dels ventarís fitosociològics a l'estudi de les suredes catalanes*. Scientia gerund., 20: 35-43.
- Wildi O. & Orlóci L., 1983. *Management and multivariate analysis of vegetation data*. Swiss Federal Institute of Forestry Research, Report 215, Birmensdorf, 68 pp.
- Wirth V., 1980. *Flechtenflora: Ökologische Kennzeichnung und Bestimmung der Flechten Südwestdeutschlands und angrenzender Gebiete*. Ulmer, Stuttgart, 552 pp.

Received February 2, 1998
Accepted September 21, 1998

Finito di stampare
il 20 novembre 1998
dalla Tipografia Villaggio del Fanciullo
Opicina (Trieste)

Studia Geobotanica Vol. 16 1998

INDEX

- A. Pirola. Un ricordo di Duilio Lausi (1923-1997). 3-4
- L. Poldini, G. Oriolo and G. Mazzolini. The segetal vegetation of vineyards and crops fields in Friuli – Venezia Giulia (NE Italy). 5-32
- P. Bianco, S. Brullo, P. Minissale, P. Signorello e G. Spampinato. Considerazioni fitosociologiche sui boschi a *Quercus trojana* Webb della Puglia (Italia Meridionale). 33-38
- L. Coassini Lokar and F. Martini. On the presence of phytosterols in some species of *Euphorbia* L. 39-43
- F. Cateni, L. Coassini Lokar and F. Martini. Chemosystematic value of aristolochic acids in the genus *Aristolochia* L. in Friuli – Venezia Giulia (NE Italy). 45-49
- M. Aleffi, M. Privitera and M. Puglisi. Phytosociological observations on the bryological vegetation of Mt. Palon (Mt. Bondone group, Trentino – Alto Adige). 51-56
- B. Litterski and H. Mayrhofer. Catalogue of lichenized and lichenicolous fungi of Cyprus. 57-70
- S. Fos y E. Barreno. Caracterización ecológica de los alcornocales Iberolevantineos (este de España) basada en bioindicadores liquénicos. 71-83